

BOTTOM, STRANGE MESONS ($B = \pm 1$, $S = \mp 1$)

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^* \text{'s}$$

B_s^0

$$I(J^P) = 0(0^-)$$

I , J , P need confirmation. Quantum numbers shown are quark-model predictions.

Mass $m_{B_s^0} = 5366.93 \pm 0.10$ MeV

$m_{B_s^0} - m_B = 87.37 \pm 0.12$ MeV

Mean life $\tau = (1.520 \pm 0.005) \times 10^{-12}$ s

$$c\tau = 455.7 \mu\text{m}$$

$$\Delta\Gamma_{B_s^0} = \Gamma_{B_{sL}^0} - \Gamma_{B_{sH}^0} = (0.083 \pm 0.005) \times 10^{12} \text{ s}^{-1} \quad (S = 1.8)$$

B_s^0 - \bar{B}_s^0 mixing parameters

$$\begin{aligned} \Delta m_{B_s^0} &= m_{B_{sH}^0} - m_{B_{sL}^0} = (17.765 \pm 0.006) \times 10^{12} \hbar \text{ s}^{-1} \\ &= (1.1693 \pm 0.0004) \times 10^{-8} \text{ MeV} \end{aligned}$$

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0} = 26.99 \pm 0.09$$

$$\chi_s (B_s^0\text{-}\bar{B}_s^0 \text{ mixing parameter}) = 0.499318 \pm 0.000005$$

CP violation parameters in B_s^0

$$\text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2) = (-0.15 \pm 0.70) \times 10^{-3}$$

$$C_{KK}(B_s^0 \rightarrow K^+ K^-) = 0.162 \pm 0.035$$

$$S_{KK}(B_s^0 \rightarrow K^+ K^-) = 0.14 \pm 0.05 \quad (S = 1.3)$$

$$r_B(B_s^0 \rightarrow D_s^\mp K^\pm) = 0.37^{+0.10}_{-0.09}$$

$$r_B(B_s^0 \rightarrow D_s^\mp K^\pm \pi^\pm \pi^\mp) = 0.47 \pm 0.08$$

$$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp) = (358 \pm 14)^\circ$$

$$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp \pi^\pm \pi^\mp) = (-6^{+10}_{-13})^\circ$$

$$CP \text{ Violation phase } \beta_s (b \rightarrow c\bar{c}s) = (2.0 \pm 0.8) \times 10^{-2} \text{ rad}$$

$$CP \text{ Violation phase } \beta_s (b \rightarrow s\bar{s}s) = (3.7 \pm 3.5) \times 10^{-2} \text{ rad}$$

$$|\lambda| (B_s^0 \rightarrow J/\psi(1S)\phi) = 0.988 \pm 0.009$$

$$|\lambda| (b \rightarrow c\bar{c}s) = 0.989 \pm 0.008$$

$$A, CP \text{ violation parameter} = -0.79 \pm 0.08$$

$$C, CP \text{ violation parameter} = 0.19 \pm 0.06$$

$$S, CP \text{ violation parameter} = 0.17 \pm 0.06$$

$$A_{CP}^L(B_s \rightarrow J/\psi \bar{K}^*(892)^0) = -0.05 \pm 0.06$$

$$\begin{aligned}
A_{CP}^{\parallel}(B_s \rightarrow J/\psi \bar{K}^*(892)^0) &= 0.17 \pm 0.15 \\
A_{CP}^{\perp}(B_s \rightarrow J/\psi \bar{K}^*(892)^0) &= -0.05 \pm 0.10 \\
\mathbf{A}_{CP}(B_s \rightarrow \pi^+ K^-) &= 0.224 \pm 0.012 \\
A_{CP}(B_s^0 \rightarrow [K^+ K^-]_D \bar{K}^*(892)^0) &= -0.04 \pm 0.07 \\
A_{CP}(B_s^0 \rightarrow [\pi^+ K^-]_D K^*(892)^0) &= -0.01 \pm 0.04 \\
A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0) &= 0.06 \pm 0.13 \\
S(B_s^0 \rightarrow \phi \gamma) &= 0.43 \pm 0.32 \\
C(B_s^0 \rightarrow \phi \gamma) &= 0.11 \pm 0.31 \\
A^{\Delta}(B_s^0 \rightarrow \phi \gamma) &= -0.7 \pm 0.4 \\
\Delta a_{\perp} &< 1.2 \times 10^{-12} \text{ GeV, CL = 95\%} \\
\Delta a_{\parallel} &= (-0.9 \pm 1.5) \times 10^{-14} \text{ GeV} \\
\Delta a_X &= (1.0 \pm 2.2) \times 10^{-14} \text{ GeV} \\
\Delta a_Y &= (-3.8 \pm 2.2) \times 10^{-14} \text{ GeV} \\
\text{Re}(\xi) &= -0.022 \pm 0.033 \\
\text{Im}(\xi) &= 0.004 \pm 0.011
\end{aligned}$$

These branching fractions all scale with $B(\bar{b} \rightarrow B_s^0)$.

The branching fraction $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_{\ell} \text{anything})$ is not a pure measurement since the measured product branching fraction $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_{\ell} \text{anything})$ was used to determine $B(\bar{b} \rightarrow B_s^0)$, as described in the note on “ B^0 - \bar{B}^0 Mixing”

For inclusive branching fractions, e.g., $B \rightarrow D^{\pm}$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

B_s^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
D_s^- anything	(62 \pm 6) %	—	—
D_s^{\pm} anything	(92 \pm 11) %	—	—
D_s^0/\bar{D}^0 anything	(38 \pm 10) %	—	—
$\ell \nu_{\ell} X$	(9.6 \pm 0.8) %	—	—
$e^+ \nu X^-$	(9.1 \pm 0.8) %	—	—
$\mu^+ \nu X^-$	(10.2 \pm 1.0) %	—	—
$D_s^- \ell^+ \nu_{\ell}$ anything	[a] (8.1 \pm 1.3) %	—	—
$D_s^{*-} \ell^+ \nu_{\ell}$ anything	(5.4 \pm 1.1) %	—	—
$D_s^- \mu^+ \nu_{\mu}$	(2.31 \pm 0.21) %	2321	—
$D_s^{*-} \mu^+ \nu_{\mu}$	(5.2 \pm 0.5) %	2266	—
$D_{s1}(2536)^- \mu^+ \nu_{\mu}, D_{s1}^- \rightarrow D^{*-} K_S^0$	(2.7 \pm 0.7) $\times 10^{-3}$	—	—
$D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+$	(4.4 \pm 1.3) $\times 10^{-3}$	—	—
$D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+$	(2.7 \pm 1.0) $\times 10^{-3}$	—	—

$K^- \mu^+ \nu_\mu$	$(1.06 \pm 0.09) \times 10^{-4}$	2660
$D_s^- \pi^+$	$(2.98 \pm 0.14) \times 10^{-3}$	2320
$D_s^- \rho^+$	$(6.8 \pm 1.4) \times 10^{-3}$	2249
$D_s^- \pi^+ \pi^+ \pi^-$	$(6.1 \pm 1.0) \times 10^{-3}$	2301
$D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-$	$(2.4 \pm 0.8) \times 10^{-5}$	—
$D_s^\mp K^\pm$	$(2.25 \pm 0.12) \times 10^{-4}$	2293
$D_{s1}(2536)^\mp K^\pm, D_{s1}^- \rightarrow \bar{D}^*(2007)^0 K^-$	$(2.48 \pm 0.28) \times 10^{-5}$	—
$D_s^- K^+ \pi^+ \pi^-$	$(3.2 \pm 0.6) \times 10^{-4}$	2249
$D_s^+ D_s^-$	$(4.4 \pm 0.5) \times 10^{-3}$	1824
$D_s^- D^+$	$(2.8 \pm 0.5) \times 10^{-4}$	1875
$D^+ D^-$	$(2.2 \pm 0.6) \times 10^{-4}$	1925
$D^{*-} D^{*-}$	$(2.14 \pm 0.32) \times 10^{-4}$	1778
$D^0 \bar{D}^0$	$(1.9 \pm 0.5) \times 10^{-4}$	1930
$D_s^{*-} \pi^+$	$(1.9 \pm 0.5) \times 10^{-3}$	2265
$D_s^{*\mp} K^\pm$	$(1.32 \pm 0.40) \times 10^{-4}$	—
$D_s^{*-} \rho^+$	$(9.5 \pm 2.0) \times 10^{-3}$	2191
$D_s^{*+} D_s^- + D_s^{*-} D_s^+$	$(1.39 \pm 0.17) \%$	1742
$D_s^{*+} D_s^{*-}$	$(1.44 \pm 0.21) \%$	S=1.1 1655
$D_s^{(*)+} D_s^{(*)-}$	$(4.5 \pm 1.4) \%$	—
$D_s^{*-} D_s^+$	$(3.9 \pm 0.8) \times 10^{-4}$	1801
$\bar{D}^{*0} \bar{K}^0$	$(2.8 \pm 1.1) \times 10^{-4}$	2278
$\bar{D}^0 \bar{K}^0$	$(4.3 \pm 0.9) \times 10^{-4}$	2330
$\bar{D}^0 K^- \pi^+$	$(1.04 \pm 0.13) \times 10^{-3}$	2312
$\bar{D}^*(2007)^0 K^- \pi^+$	$(7.3 \pm 2.6) \times 10^{-4}$	2259
$\bar{D}^0 \bar{K}^*(892)^0$	$(4.4 \pm 0.6) \times 10^{-4}$	2264
$\bar{D}^0 \bar{K}^*(1410)$	$(3.9 \pm 3.5) \times 10^{-4}$	2117
$\bar{D}^0 \bar{K}_0^*(1430)$	$(3.0 \pm 0.7) \times 10^{-4}$	2113
$\bar{D}^0 \bar{K}_2^*(1430)$	$(1.1 \pm 0.4) \times 10^{-4}$	2112
$\bar{D}^0 \bar{K}^*(1680)$	$< 7.8 \times 10^{-5}$	CL=90% 1997
$\bar{D}^0 \bar{K}_0^*(1950)$	$< 1.1 \times 10^{-4}$	CL=90% 1884
$\bar{D}^0 \bar{K}_3^*(1780)$	$< 2.6 \times 10^{-5}$	CL=90% 1970
$\bar{D}^0 \bar{K}_4^*(2045)$	$< 3.1 \times 10^{-5}$	CL=90% 1835
$\bar{D}^0 K^- \pi^+ (\text{non-resonant})$	$(2.1 \pm 0.8) \times 10^{-4}$	2312
$D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \bar{D}^0 K^-$	$(2.6 \pm 0.4) \times 10^{-4}$	—
$D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-$	$(1.6 \pm 0.8) \times 10^{-5}$	—
$D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-$	$(5 \pm 4) \times 10^{-5}$	—

$D_{s3}^*(2860)^-\pi^+$, $D_{s3}^* \rightarrow \overline{D}^0 K^-$	$(2.2 \pm 0.6) \times 10^{-5}$	-
$\overline{D}^0 K^+ K^-$	$(5.6 \pm 0.9) \times 10^{-5}$	2243
$\overline{D}^0 f_0(980)$	$< 3.1 \times 10^{-6}$	CL=90% 2242
$\overline{D}^0 \phi$	$(2.30 \pm 0.25) \times 10^{-5}$	2235
$\overline{D}^{*0} \phi$	$(3.2 \pm 0.4) \times 10^{-5}$	2178
$D^{*\mp} \pi^\pm$	$< 6.1 \times 10^{-6}$	CL=90% -
$\eta_c \phi$	$(5.0 \pm 0.9) \times 10^{-4}$	1663
$\eta_c \pi^+ \pi^-$	$(1.8 \pm 0.7) \times 10^{-4}$	1840
$J/\psi(1S) \phi$	$(1.04 \pm 0.04) \times 10^{-3}$	1588
$J/\psi(1S) \phi \phi$	$(1.20 \pm 0.14) \times 10^{-5}$	764
$J/\psi(1S) \pi^0$	$< 1.2 \times 10^{-3}$	CL=90% 1787
$J/\psi(1S) \eta$	$(4.0 \pm 0.7) \times 10^{-4}$	S=1.4 1733
$J/\psi(1S) K_S^0$	$(1.92 \pm 0.14) \times 10^{-5}$	1743
$J/\psi(1S) \overline{K}^*(892)^0$	$(4.1 \pm 0.4) \times 10^{-5}$	1637
$J/\psi(1S) \eta'$	$(3.3 \pm 0.4) \times 10^{-4}$	1612
$J/\psi(1S) \pi^+ \pi^-$	$(2.02 \pm 0.17) \times 10^{-4}$	S=1.7 1775
$J/\psi(1S) f_0(500)$, $f_0 \rightarrow \pi^+ \pi^-$	$< 4 \times 10^{-6}$	CL=90% -
$J/\psi(1S) \rho$, $\rho \rightarrow \pi^+ \pi^-$	$< 3.4 \times 10^{-6}$	CL=90% -
$J/\psi(1S) f_0(980)$, $f_0 \rightarrow \pi^+ \pi^-$	$(1.24 \pm 0.15) \times 10^{-4}$	S=2.1 -
$J/\psi(1S) f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.0 \pm 0.4) \times 10^{-6}$	-
$J/\psi(1S) f_2(1270)_0$, $f_2 \rightarrow \pi^+ \pi^-$	$(7.3 \pm 1.7) \times 10^{-7}$	-
$J/\psi(1S) f_2(1270)_{ }$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.05 \pm 0.33) \times 10^{-6}$	-
$J/\psi(1S) f_2(1270)_{\perp}$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.3 \pm 0.7) \times 10^{-6}$	-
$J/\psi(1S) f_0(1370)$, $f_0 \rightarrow \pi^+ \pi^-$	$(4.4 \pm 0.6) \times 10^{-5}$	-
$J/\psi(1S) f_0(1500)$, $f_0 \rightarrow \pi^+ \pi^-$	$(2.04 \pm 0.32) \times 10^{-5}$	-
$J/\psi(1S) f'_2(1525)_0$, $f'_2 \rightarrow \pi^+ \pi^-$	$(1.03 \pm 0.22) \times 10^{-6}$	-
$J/\psi(1S) f'_2(1525)_{ }$, $f'_2 \rightarrow \pi^+ \pi^-$	$(1.2 \pm 2.6) \times 10^{-7}$	-
$J/\psi(1S) f'_2(1525)_{\perp}$, $f'_2 \rightarrow \pi^+ \pi^-$	$(5 \pm 4) \times 10^{-7}$	-
$J/\psi(1S) f_0(1790)$, $f_0 \rightarrow \pi^+ \pi^-$	$(4.9 \pm 10.0) \times 10^{-6}$	-
$J/\psi(1S) \pi^+ \pi^-$ (nonresonant)	$(1.74 \pm 1.10) \times 10^{-5}$	1775

$J/\psi(1S)\bar{K}^0\pi^+\pi^-$	$< 4.4 \times 10^{-5}$	CL=90%	1675
$J/\psi(1S)K^+K^-$	$(7.9 \pm 0.7) \times 10^{-4}$		1601
$J/\psi(1S)K^0K^-\pi^++\text{c.c.}$	$(9.5 \pm 1.3) \times 10^{-4}$		1538
$J/\psi(1S)\bar{K}^0K^+K^-$	$< 1.2 \times 10^{-5}$	CL=90%	1333
$J/\psi K^*(892)^0\bar{K}^*(892)^0$	$(1.10 \pm 0.09) \times 10^{-4}$		1083
$J/\psi(1S)f'_2(1525)$	$(2.6 \pm 0.6) \times 10^{-4}$		1310
$J/\psi(1S)p\bar{p}$	$(3.6 \pm 0.4) \times 10^{-6}$		982
$J/\psi(1S)\gamma$	$< 7.3 \times 10^{-6}$	CL=90%	1790
$J/\psi\mu^+\mu^-,\ J/\psi \rightarrow \mu^+\mu^-$	$< 2.6 \times 10^{-9}$	CL=95%	—
$J/\psi(1S)\pi^+\pi^-\pi^+\pi^-$	$(7.5 \pm 0.8) \times 10^{-5}$		1731
$J/\psi(1S)f_1(1285)$	$(7.2 \pm 1.4) \times 10^{-5}$		1460
$\psi(2S)\eta$	$(3.3 \pm 0.9) \times 10^{-4}$		1338
$\psi(2S)\eta'$	$(1.29 \pm 0.35) \times 10^{-4}$		1158
$\psi(2S)\pi^+\pi^-$	$(6.9 \pm 1.2) \times 10^{-5}$		1397
$\psi(2S)\phi$	$(5.3 \pm 0.4) \times 10^{-4}$		1120
$\psi(2S)K^0$	$(1.9 \pm 0.5) \times 10^{-5}$		1352
$\psi(2S)K^-\pi^+$	$(3.1 \pm 0.4) \times 10^{-5}$		1310
$\psi(2S)\bar{K}^*(892)^0$	$(3.3 \pm 0.5) \times 10^{-5}$		1196
$\chi_{c1}\phi$	$(1.97 \pm 0.25) \times 10^{-4}$		1275
$\chi_{c1}(3872)\phi$	$(1.22 \pm 0.35) \times 10^{-4}$		936
$\chi_{c1}(3872)(K^+K^-)_{non-\phi}$	$(9.4 \pm 3.4) \times 10^{-5}$		961
$\chi_{c1}(3872)\pi^+\pi^-$	$(4.6 \pm 1.6) \times 10^{-5}$		1264
$\pi^+\pi^-$	$(7.2 \pm 1.0) \times 10^{-7}$		2680
$\pi^0\pi^0$	$< 7.7 \times 10^{-6}$	CL=90%	2680
$\eta\pi^0$	$< 1.0 \times 10^{-3}$	CL=90%	2654
$\eta\eta$	$< 1.43 \times 10^{-4}$	CL=90%	2627
$\rho^0\rho^0$	$< 3.20 \times 10^{-4}$	CL=90%	2569
$\eta'K_S^0$	$< 8.16 \times 10^{-6}$	CL=90%	2573
$\eta'\eta$	$< 6.5 \times 10^{-5}$	CL=90%	2568
$\eta'\eta'$	$(3.3 \pm 0.7) \times 10^{-5}$		2507
$\eta'\phi$	$< 8.2 \times 10^{-7}$	CL=90%	2495
$\phi f_0(980),\ f_0(980) \rightarrow \pi^+\pi^-$	$(1.12 \pm 0.21) \times 10^{-6}$		—
$\phi f_2(1270),\ f_2(1270) \rightarrow \pi^+\pi^-$	$(6.1 \pm 1.8) \times 10^{-7}$		—
$\phi\rho^0$	$(2.7 \pm 0.8) \times 10^{-7}$		2526
$\phi\pi^+\pi^-$	$(3.5 \pm 0.5) \times 10^{-6}$		2579
$\phi\phi$	$(1.85 \pm 0.14) \times 10^{-5}$		2482
$\phi\phi\phi$	$(2.2 \pm 0.6) \times 10^{-6}$		2165
π^+K^-	$(5.9 \pm 0.7) \times 10^{-6}$		2659
K^+K^-	$(2.72 \pm 0.23) \times 10^{-5}$		2638
$K^0\bar{K}^0$	$(1.76 \pm 0.31) \times 10^{-5}$		2637
$K^0\pi^+\pi^-$	$(9.5 \pm 2.1) \times 10^{-6}$		2653
$K^0K^\pm\pi^\mp$	$(8.4 \pm 0.9) \times 10^{-5}$		2622
$K^*(892)^-\pi^+$	$(2.9 \pm 1.1) \times 10^{-6}$		2607

$K^*(892)^{\pm} K^{\mp}$	$(1.9 \pm 0.5) \times 10^{-5}$	2585
$K_0^*(1430)^{\pm} K^{\mp}$	$(3.1 \pm 2.5) \times 10^{-5}$	—
$K_2^*(1430)^{\pm} K^{\mp}$	$(1.0 \pm 1.7) \times 10^{-5}$	—
$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(2.0 \pm 0.6) \times 10^{-5}$	2585
$K_0^*(1430)^0 \bar{K}^0 + \text{c.c.}$	$(3.3 \pm 1.0) \times 10^{-5}$	2468
$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.}$	$(1.7 \pm 2.2) \times 10^{-5}$	2467
$K_S^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$	2585
$K^0 K^+ K^-$	$(1.3 \pm 0.6) \times 10^{-6}$	2568
$\bar{K}^*(892)^0 \rho^0$	$< 7.67 \times 10^{-4}$ CL=90%	2550
$\bar{K}^*(892)^0 K^*(892)^0$	$(1.11 \pm 0.27) \times 10^{-5}$	2531
$\phi K^*(892)^0$	$(1.14 \pm 0.30) \times 10^{-6}$	2507
$p\bar{p}$	$< 4.4 \times 10^{-9}$ CL=90%	2514
$p\bar{p} K^+ K^-$	$(4.5 \pm 0.5) \times 10^{-6}$	2231
$p\bar{p} K^+ \pi^-$	$(1.39 \pm 0.26) \times 10^{-6}$	2355
$p\bar{p} \pi^+ \pi^-$	$(4.3 \pm 2.0) \times 10^{-7}$	2454
$p\bar{p} p\bar{p}$	$(2.3 \pm 1.0) \times 10^{-8}$	1797
$p\Lambda K^- + \text{c.c.}$	$(5.5 \pm 1.0) \times 10^{-6}$	2358
$\Lambda_c^- \Lambda \pi^+$	$(3.6 \pm 1.6) \times 10^{-4}$	1979
$\Lambda_c^- \Lambda_c^+$	$< 8.0 \times 10^{-5}$ CL=95%	1405

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

$\gamma\gamma$	<i>B1</i>	$< 3.1 \times 10^{-6}$	CL=90%	2683
$\phi\gamma$	<i>B1</i>	$(3.4 \pm 0.4) \times 10^{-5}$	—	2587
$\mu^+ \mu^-$	<i>B1</i>	$(3.34 \pm 0.27) \times 10^{-9}$	—	2681
$e^+ e^-$	<i>B1</i>	$< 9.4 \times 10^{-9}$ CL=90%	—	2683
$\tau^+ \tau^-$	<i>B1</i>	$< 6.8 \times 10^{-3}$ CL=95%	—	2011
$\mu^+ \mu^- \gamma$	<i>B1</i>	$< 2.0 \times 10^{-9}$	—	2681
$\mu^+ \mu^- \mu^+ \mu^-$	<i>B1</i>	$< 8.6 \times 10^{-10}$ CL=95%	—	2673
$SP, S \rightarrow \mu^+ \mu^-, P \rightarrow \mu^+ \mu^-$	<i>B1</i>	[b] $< 2.2 \times 10^{-9}$ CL=95%	—	—
$aa, a \rightarrow \mu^+ \mu^-$	<i>B1</i>	$< 5.8 \times 10^{-10}$ CL=95%	—	—
$\phi(1020)\mu^+ \mu^-$	<i>B1</i>	$(8.4 \pm 0.4) \times 10^{-7}$	—	2582
$f'_2(1525)\mu^+ \mu^-$	<i>B1</i>	$(1.62 \pm 0.22) \times 10^{-7}$	—	2464
$\bar{K}^*(892)^0 \mu^+ \mu^-$	<i>B1</i>	$(2.9 \pm 1.1) \times 10^{-8}$	—	2605
$\pi^+ \pi^- \mu^+ \mu^-$	<i>B1</i>	$(8.4 \pm 1.7) \times 10^{-8}$	—	2670
$\phi \nu \bar{\nu}$	<i>B1</i>	$< 5.4 \times 10^{-3}$ CL=90%	—	2587
$e^\pm \mu^\mp$	<i>LF</i>	[c] $< 5.4 \times 10^{-9}$ CL=90%	—	2682
$e^\pm \tau^\mp$	<i>LF</i>	$< 1.4 \times 10^{-3}$ CL=90%	—	2389
$\mu^\pm \tau^\mp$	<i>LF</i>	$< 4.2 \times 10^{-5}$ CL=95%	—	2388
$\phi \mu^\pm e^\mp$	<i>LF</i>	$< 1.6 \times 10^{-8}$ CL=90%	—	2586
$p \mu^-$	<i>L,B</i>	$< 1.21 \times 10^{-8}$ CL=90%	—	2600

B_s^*

$I(J^P) = 0(1^-)$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

Mass $m = 5415.4 \pm 1.4$ MeV ($S = 2.6$)

$m_{B_s^*} - m_{B_s} = 48.5 \pm 1.4$ MeV ($S = 2.6$)

B_s^* DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$B_s \gamma$	seen	48

$B_{s1}(5830)^0$

$I(J^P) = 0(1^+)$

I, J, P need confirmation.

Mass $m = 5828.73 \pm 0.20$ MeV

$m_{B_{s1}^0} - m_{B^{*+}} = 503.98 \pm 0.17$ MeV

Full width $\Gamma = 0.5 \pm 0.4$ MeV

$B_{s1}(5830)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$B^{*+} K^-$	seen	97

$B_{s2}^*(5840)^0$

$I(J^P) = 0(2^+)$

I, J, P need confirmation.

Mass $m = 5839.88 \pm 0.12$ MeV

$m_{B_{s2}^{*0}} - m_{B^+} = 560.48 \pm 0.12$ MeV

Full width $\Gamma = 1.49 \pm 0.27$ MeV

Branching fractions are given relative to the one **DEFINED AS 1**.

$B_{s2}^*(5840)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$B^+ K^-$	DEFINED AS 1	252
$B^{*+} K^-$	0.093 ± 0.018	141
$B^0 K_S^0$	0.43 ± 0.11	245
$B^{*0} K_S^0$	0.04 ± 0.04	—

NOTES

- [a] Not a pure measurement. See note at head of B_s^0 Decay Modes.
- [b] Here S and P are the hypothetical scalar and pseudoscalar particles with masses of $2.5 \text{ GeV}/c^2$ and $214.3 \text{ MeV}/c^2$, respectively.
- [c] The value is for the sum of the charge states or particle/antiparticle states indicated.