

$\Lambda_c(2880)^+$ $I(J^P) = 0(\frac{5}{2}^+)$ Status: ***

A narrow peak seen in $\Lambda_c^+\pi^+\pi^-$ and in pD^0 . It is not seen in pD^+ , and therefore it is a Λ_c^+ and not a Σ_c .

 $\Lambda_c(2880)^+ MASS$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2881.63±0.24 OUR FIT				
2881.62±0.24 OUR AVERAGE				
2881.75±0.29±0.07 ^{+0.14} _{-0.20}	1 AAIJ	17S LHCb	in $\Lambda_b^0 \rightarrow D^0 p\pi^-$	
2881.9 ± 0.1 ± 0.5	2.8k	AUBERT	07 BABR	in pD^0
2881.2 ± 0.2 ± 0.4	690	MIZUK	07 BELL	in $\Sigma_c(2455)^{0,++}\pi^\pm$

¹The third AAIJ 17S uncertainty comes from modeling the resonant shape of the $\Lambda_c(2880)^+$ and the background (non-resonant) amplitudes.

 $\Lambda_c(2880)^+ - \Lambda_c^+ MASS DIFFERENCE$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
595.17±0.28 OUR FIT				
596 ± 1 ± 2	350	ARTUSO	01 CLE2	in $\Lambda_c^+\pi^+\pi^-$

 $\Lambda_c(2880)^+ WIDTH$

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.6 ± 0.8 OUR AVERAGE					
5.43 ^{+0.77} _{-0.71} ^{+0.81} _{-0.29}		2 AAIJ	17S LHCb	in $\Lambda_b^0 \rightarrow D^0 p\pi^-$	
5.8 ± 1.5 ± 1.1	2.8k	AUBERT	07 BABR	in pD^0	
5.8 ± 0.7 ± 1.1	690	MIZUK	07 BELL	in $\Sigma_c(2455)^{0,++}\pi^\pm$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8	90	ARTUSO	01 CLEO	in $\Lambda_c^+\pi^+\pi^-$	

²AAIJ 17S reports $5.43^{+0.77}_{-0.71} \pm 0.29^{+0.75}_{-0.00}$ MeV value where the third uncertainty comes from modeling the resonant shape of the $\Lambda_c(2880)^+$ and the background (non-resonant) amplitudes. We have combined in quadrature the systematic uncertainties.

 $\Lambda_c(2880)^+ DECAY MODES$

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda_c^+\pi^+\pi^-$	seen
$\Gamma_2 \Sigma_c(2455)^{0,++}\pi^\pm$	seen
$\Gamma_3 \Sigma_c(2520)^{0,++}\pi^\pm$	seen
$\Gamma_4 \Lambda_c^+\eta$	
$\Gamma_5 pD^0$	seen

 $\Lambda_c(2880)^+ BRANCHING RATIOS$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.392±0.031 OUR AVERAGE				Error includes scale factor of 1.3.	
0.404±0.021±0.014		MIZUK	07 BELL	in $\Sigma_c(2455)^{0,++}\pi^\pm$	
0.31 ± 0.06 ± 0.03	96	ARTUSO	01 CLE2	$e^+ e^- \approx \Upsilon(4S)$	

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.091±0.025±0.010		MIZUK	07 BELL	in $\Sigma_c(2455)^{0,++}\pi^\pm$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.11	90	ARTUSO	01 CLE2	$e^+ e^- \approx \Upsilon(4S)$	

NODE=B151

NODE=B151

NODE=B151M

NODE=B151M

NODE=B151M;LINKAGE=A

NODE=B151D

NODE=B151D

NODE=B151W

NODE=B151W

NODE=B151W;LINKAGE=A

NODE=B151220;NODE=B151

DESIG=1;OUR EST

DESIG=2;OUR EST

DESIG=3;OUR EST

DESIG=5

DESIG=4;OUR EST

NODE=B151225

NODE=B151R1

NODE=B151R1

NODE=B151R2

NODE=B151R2

$\Gamma(\Sigma_c(2520)^0,++\pi^\pm)/\Gamma(\Sigma_c(2455)^0,++\pi^\pm)$				Γ_3/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.225 \pm 0.062 \pm 0.025$	³ MIZUK	07	BELL in $\Sigma_c(2455)^0,++\pi^\pm$	
³ This MIZUK 07 ratio is redundant with MIZUK 07 ratios given above.				
$\Gamma(\Lambda_c^+\eta)/\Gamma(\Sigma_c(2455)^0,++\pi^\pm)$				Γ_4/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.13	90	LI	24C	BELL e^+e^- at $\sim \gamma(nS)$
$\Gamma(pD^0)/\Gamma(\Sigma_c(2455)^0,++\pi^\pm)$				Γ_5/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.75±0.03±0.07	12k	LI	24C	BELL e^+e^- at $\sim \gamma(nS)$

$\Lambda_c(2880)^+$ REFERENCES

LI	24C	PR D110 032021	S.X. Li <i>et al.</i>	(BELLE Collab.)	REFID=62939
AAIJ	17S	JHEP 1705 030	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=57813
AUBERT	07	PRL 98 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51585
MIZUK	07	PRL 98 262001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=51816
ARTUSO	01	PRL 86 4479	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=48126