

$\Lambda_b(5920)^0$

$$J^P = \frac{3}{2}^-$$

Status: ***

Quantum numbers are based on quark model expectations.

NODE=B163

NODE=B163

NODE=B163M

NODE=B163M

SYCLP=A

SYCLP=A

SYCLP=A

SYCLP=A

NODE=B163M;LINKAGE=A

NODE=B163M;LINKAGE=B

NODE=B163M;LINKAGE=AL

NODE=B163M;LINKAGE=AT

NODE=B163M;LINKAGE=AA

NODE=B163M;LINKAGE=AI

NODE=B163W

NODE=B163W

NODE=B163215;NODE=B163

DESIG=1

NODE=B163220

NODE=B163R01

NODE=B163R01

NODE=B163

REFID=60505
REFID=60392
REFID=55450
REFID=54589 $\Lambda_b(5920)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5920.07±0.16 OUR AVERAGE			
5920.06±0.02±0.16	¹ AAIJ	20Q LHCB	pp at 7, 8, 13 TeV
5920.13±0.07±0.16	² SIRUNYAN	20K CMS	pp at 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5919.4 ±0.5 ±0.2	^{3,4} AALTONEN	13V CDF	$p\bar{p}$ at 1.96 TeV
5919.97±0.09±0.16	^{5,6} AAIJ	12AL LHCB	Repl. by AAIJ 20Q
¹ AAIJ 20Q measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.492 \pm 0.019 \pm 0.010$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.57 \pm 0.16$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values.			
² SIRUNYAN 20K measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.56 \pm 0.07 \pm 0.01$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.57 \pm 0.16$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values.			
³ Measured in $\Lambda_b(5920)^0 \rightarrow \Lambda_b^0 \pi^+ \pi^-$ decays with $17.3^{+5.3}_{-4.6}$ events, with a significance of 3.5 sigma.			
⁴ AALTONEN 13V measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) - 2m(\pi) = 20.68 \pm 0.35 \pm 0.30$ MeV. We have adjusted the measurement to our best values of $m(\Lambda_b^0) = 5619.57 \pm 0.16$ MeV and $m(\pi) = 139.57039 \pm 0.00018$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values.			
⁵ Observed in $\Lambda_b(5920)^0 \rightarrow \Lambda_b^0 \pi^+ \pi^-$ decays with 52.5 ± 8.1 candidates with a significance of 10.2 sigma.			
⁶ AAIJ 12AL measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.40 \pm 0.08 \pm 0.04$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.57 \pm 0.16$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values.			

 $\Lambda_b(5920)^0$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.19	90	AAIJ	20Q LHCB	pp at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.63	90	AAIJ	12AL LHCB	Repl. by AAIJ 20Q

 $\Lambda_b(5920)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Lambda_b^0 \pi^+ \pi^-$	seen

 $\Lambda_b(5920)^0$ BRANCHING RATIOS

$\Gamma(\Lambda_b^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	AAIJ	20Q	LHCB	pp at 7, 8, 13 TeV
seen	SIRUNYAN	20K	LHCB	pp at 13 TeV
seen	AALTONEN	13v	CDF	$p\bar{p}$ at 1.96 TeV
seen	AAIJ	12AL	LHCB	pp at 7 TeV

 $\Lambda_b(5920)^0$ REFERENCES

AAIJ	20Q	JHEP 2006 136	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	20K	PL B803 135345	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN	13V	PR D88 071101	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	12AL	PRL 109 172003	R. Aaij <i>et al.</i>	(LHCb Collab.)