

$f_2(1270)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f_2(1270)$ T-MATRIX POLE \sqrt{s}** Note that $\Gamma = -2 \operatorname{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1260–1283) – i (90–110) OUR ESTIMATE			
(1268 ± 8) – i (101 ± 6)	RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
(1263.3 ± 0.2 ± 1.5) – i (96.9 ± 0.2 ± 0.8)	ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
(1270 ± 8) – i (97 ± 8)	¹ ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$
(1278 ± 5) – i (102 ± 10)	¹ BERTIN	97C	OBLX 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

¹ Amplitude did not include dispersive corrections. **$f_2(1270)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1275.4 ± 0.8 OUR AVERAGE				
1275.8 ± 1.0 ± 0.4		1 BOGOLYUB...	13 SPEC	$7\pi^+(K^+, p)A \rightarrow n\gamma + X$
1262 $\pm \frac{1}{2}$ ± 8		2 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1275 ± 15		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1283 ± 5		ALDE	98 GAM4	$100\pi^-p \rightarrow \pi^0\pi^0n$
1272 ± 8	200k	PROKOSHKIN	94 GAM2	$38\pi^-p \rightarrow \pi^0\pi^0n$
1269.7 ± 5.2	5730	AUGUSTIN	89 DM2	$e^+e^- \rightarrow 5\pi$
1283 ± 8	400	³ ALDE	87 GAM4	$100\pi^-p \rightarrow 4\pi^0n$
1274 ± 5		³ AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1283 ± 6		⁴ LONGACRE	86 MPS	$22\pi^-p \rightarrow n2K_S^0$
1276 ± 7		COURAU	84 DLCO	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
1273.3 ± 2.3		⁵ CHABAUD	83 ASPK	17 π^-p polarized
1280 ± 4		⁶ CASON	82 STRC	$8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$
1281 ± 7	11600	GIDAL	81 MRK2	J/ψ decay
1282 ± 5		⁷ CORDEN	79 OMEG	$12\text{--}15\pi^-p \rightarrow n2\pi$
1269 ± 4	10k	APEL	75 NICE	$40\pi^-p \rightarrow n2\pi^0$
1272 ± 4	4600	ENGLER	74 DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$
1277 ± 4	5300	FLATTE	71 HBC	$7.0\pi^+p$
1273 ± 8		³ STUNTEBECK	70 HBC	$8\pi^-p, 5.4\pi^+d$
1265 ± 8		BOESEBECK	68 HBC	$8\pi^+p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1257 ± 6		⁸ KLEMPT	22 RVUE	$J/\psi(1S) \rightarrow \gamma\pi^0\pi^0, \gamma K_S^0 K_S^0$
1263 ± 12		CARVER	21 CLAS	$\gamma p \rightarrow \pi^0\pi^0p$
1259 ± 4 ± 4	1.7k	^{9,10} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1267 ± 4 ± 3	1.5k	^{9,10} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1277 ± 6	870	¹¹ SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1251 ± 10		TIKHOMIROV	03 SPEC	$40.0\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1260 ± 10		¹² ALDE	97 GAM2	$450pp \rightarrow pp\pi^0\pi^0$
1278 ± 6		¹² GRYGOREV	96 SPEC	$40\pi^-N \rightarrow K_S^0 K_S^0 X$
1262 ± 11		AGUILAR-...	91 EHS	$400pp$
1275 ± 10		AKER	91 CBAR	$0.0\bar{p}p \rightarrow 3\pi^0$
1220 ± 10		BREAKSTONE	90 SFM	$pp \rightarrow pp\pi^+\pi^-$
1288 ± 12		ABACHI	86B HRS	$e^+e^- \rightarrow \pi^+\pi^-X$
1284 ± 30	3k	BINON	83 GAM2	$38\pi^-p \rightarrow n2\eta$
1280 ± 20	3k	APEL	82 CNTR	$25\pi^-p \rightarrow n2\pi^0$
1284 ± 10	16000	DEUTSCH...	76 HBC	$16\pi^+p$
1258 ± 10	600	TAKAHASHI	72 HBC	$8\pi^-p \rightarrow n2\pi$
1275 ± 13		ARMENISE	70 HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$
1261 ± 5	1960	³ ARMENISE	68 DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$
1270 ± 10	360	³ ARMENISE	68 DBC	$5.1\pi^+n \rightarrow p\pi^0MM$
1268 ± 6		¹³ JOHNSON	68 HBC	$3.7\text{--}4.2\pi^-p$

NODE=M005PP

NODE=M005PP

NODE=M005PP

→ UNCHECKED ←

NODE=M005PP;LINKAGE=A

NODE=M005M

NODE=M005M

OCCUR=2

OCCUR=2

OCCUR=2

- ¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
² Breit-Wigner mass.
³ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
⁵ From an energy-independent partial-wave analysis.
⁶ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.
⁷ From an amplitude analysis of $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ scattering data.
⁸ Fit of the tensor partial waves from BES3 in the multipole basis.
⁹ Using CLEO-c data but not authored by the CLEO Collaboration.
¹⁰ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 185$ MeV.
¹¹ From analysis of L3 data at 91 and 183–209 GeV.
¹² Systematic uncertainties not estimated.
¹³ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

$f_2(1270)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$186.6^{+2.8}_{-2.2}$ OUR FIT				Error includes scale factor of 1.5.
$185.8^{+2.8}_{-2.1}$ OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
190.3 ± 1.9 ± 1.8		1 BOGOLYUB...	SPEC	$7\pi^+(K^+, p)A \rightarrow n\gamma + X$
175 ± 6 ± 10		2 ABLIKIM	06V	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
190 ± 20		ABLIKIM	05	$J/\psi \rightarrow \phi\pi^+\pi^-$
171 ± 10		ALDE	98	$100\pi^- p \rightarrow \pi^0\pi^0 n$
192 ± 5	200k	PROKOSHKIN	94	$38\pi^- p \rightarrow \pi^0\pi^0 n$
180 ± 24		AGUILAR-...	91	$400pp$
169 ± 9	5730	AUGUSTIN	89	$e^+ e^- \rightarrow 5\pi$
150 ± 30	400	ALDE	87	$100\pi^- p \rightarrow 4\pi^0 n$
186 ± 9		LONGACRE	86	$22\pi^- p \rightarrow n2K_S^0$
179.2 ± 6.9		CHABAUD	83	$17\pi^- p$ polarized
160 ± 11		DENNEY	83	$10\pi^+ N$
196 ± 10	3k	APEL	82	$25\pi^- p \rightarrow n2\pi^0$
152 ± 9		CASON	82	$8\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$
186 ± 27	11600	GIDAL	81	J/ψ decay
216 ± 13		CORDEN	79	$12-15\pi^- p \rightarrow n2\pi$
190 ± 10	10k	APEL	75	$40\pi^- p \rightarrow n2\pi^0$
192 ± 16	4600	ENGLER	74	$6\pi^+ n \rightarrow \pi^+\pi^- p$
183 ± 15	5300	FLATTE	71	$7\pi^+ p \rightarrow \Delta^{++}f_2$
196 ± 30		STUNTEBECK	70	$8\pi^- p, 5.4\pi^+ d$
216 ± 20	1960	ARMENISE	68	$5.1\pi^+ n \rightarrow p\pi^+ MM^-$
128 ± 27		BOESEBECK	68	$8\pi^+ p$
176 ± 21		JOHNSON	68	$3.7-4.2\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
168 ± 7		9 KLEMPT	22	$J/\psi(1S) \rightarrow \gamma\pi^0\pi^0, \gamma K_S^0 K_S^0$
183 ± 2		CARVER	21	$\gamma p \rightarrow \pi^0\pi^0 p$
195 ± 15	870	10 SCHEGELSKY	06A	$\gamma\gamma \rightarrow K_S^0 K_S^0$
121 ± 26		TIKHOMIROV	03	$40.0\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
187 ± 20		11 ALDE	97	$450pp \rightarrow pp\pi^0\pi^0$
184 ± 10		11 GRYGOREV	96	$40\pi^- N \rightarrow K_S^0 K_S^0 X$
200 ± 10		AKER	91	$0.0\bar{p}p \rightarrow 3\pi^0$
240 ± 40	3k	BINON	83	$38\pi^- p \rightarrow n2\eta$
187 ± 30	650	3 ANTIPOV	77	$25\pi^- p \rightarrow p3\pi$
225 ± 38	16000	DEUTSCH...	76	$16\pi^+ p$
166 ± 28	600	3 TAKAHASHI	72	$8\pi^- p \rightarrow n2\pi$
173 ± 53		3 ARMENISE	70	$9\pi^+ n \rightarrow p\pi^+\pi^-$

NODE=M005M;LINKAGE=B

NODE=M005M;LINKAGE=K

NODE=M005M;LINKAGE=T

NODE=M005M;LINKAGE=L

NODE=M005M;LINKAGE=O

NODE=M005M;LINKAGE=P

NODE=M005M;LINKAGE=S

NODE=M005M;LINKAGE=Q

NODE=M005M;LINKAGE=C

NODE=M005M;LINKAGE=D

NODE=M005M;LINKAGE=SC

NODE=M005M;LINKAGE=QQ

NODE=M005M;LINKAGE=J

NODE=M005W


NODE=M005W

OCCUR=2

OCCUR=2

- ¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
² Breit-Wigner width
³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
⁵ From an energy-independent partial-wave analysis.
⁶ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.
⁷ From an amplitude analysis of $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ scattering data.
⁸ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.
⁹ Fit of the tensor partial waves from BES3 in the multipole basis.
¹⁰ From analysis of L3 data at 91 and 183–209 GeV.
¹¹ Systematic uncertainties not estimated.

NODE=M005W;LINKAGE=C
NODE=M005W;LINKAGE=D
NODE=M005W;LINKAGE=T
NODE=M005W;LINKAGE=L
NODE=M005W;LINKAGE=R
NODE=M005W;LINKAGE=Q
NODE=M005W;LINKAGE=U
NODE=M005W;LINKAGE=J
NODE=M005W;LINKAGE=M
NODE=M005W;LINKAGE=SC
NODE=M005W;LINKAGE=QQ



$f_2(1270)$ width (MeV)

$f_2(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi\pi$	$(84.3 \pm 2.8) \%$	S=1.2
$\Gamma_2 \pi^+ \pi^- 2\pi^0$	$(7.7 \pm 1.2) \%$	S=1.2
$\Gamma_3 K\bar{K}$	$(4.6 \pm 0.4) \%$	S=2.7
$\Gamma_4 2\pi^+ 2\pi^-$	$(2.8 \pm 0.4) \%$	S=1.2
$\Gamma_5 \eta\eta$	$(4.0 \pm 0.8) \times 10^{-3}$	S=2.1
$\Gamma_6 4\pi^0$	$(3.0 \pm 1.0) \times 10^{-3}$	
$\Gamma_7 \gamma\gamma$	$(1.42 \pm 0.24) \times 10^{-5}$	S=1.4
$\Gamma_8 \eta\pi\pi$	$< 8 \times 10^{-3}$	CL=95%
$\Gamma_9 K^0 K^- \pi^+ + \text{c.c.}$	$< 3.4 \times 10^{-3}$	CL=95%
$\Gamma_{10} e^+ e^-$	$< 6 \times 10^{-10}$	CL=90%

NODE=M005215;NODE=M005

DESIG=1
DESIG=3
DESIG=4
DESIG=2
DESIG=7
DESIG=9
DESIG=8
DESIG=6
DESIG=5
DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 44 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 82.3$ for 37 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-91						
x_3	10 -39						
x_4	10 -38 1						
x_5	1 -6 0 0						
x_6	0 -7 0 0 0						
x_7	4 1 -15 0 0 0						
Γ	-72	66	-10	-7	-1	0	-6
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

	Mode	Rate (MeV)		Scale factor	
Γ_1	$\pi\pi$	157.2	$+5.0$ -1.0		DESIG=1
Γ_2	$\pi^+ \pi^- 2\pi^0$	14.3	$+2.3$ -6.0	1.3	DESIG=3
Γ_3	$K\bar{K}$	8.5	± 0.8	2.8	DESIG=4
Γ_4	$2\pi^+ 2\pi^-$	5.2	± 0.7	1.2	DESIG=2
Γ_5	$\eta\eta$	0.75	± 0.14	2.1	DESIG=7
Γ_6	$4\pi^0$	0.56	± 0.19		DESIG=9
Γ_7	$\gamma\gamma$	0.0026	± 0.0005	1.4	DESIG=8

$\Gamma_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$	Γ_1
VALUE (MeV) <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

157.2 $^{+5.0}_{-1.0}$ OUR FIT

157.0 $^{+6.0}_{-1.0}$ ¹ LONGACRE 86 MPS $22 \pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152 ± 8 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

2 From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

$\Gamma(K\bar{K})$	Γ_3
VALUE (MeV) <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

8.5 ± 0.8 OUR FIT Error includes scale factor of 2.8.

9.0 $^{+0.7}_{-0.3}$ ¹ LONGACRE 86 MPS $22 \pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.0 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

2 From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

$\Gamma(\eta\eta)$	Γ_5
VALUE (MeV) <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

0.75 ± 0.14 OUR FIT Error includes scale factor of 2.1.

1.0 ± 0.1 ¹ LONGACRE 86 MPS $22 \pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.4 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

2 From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005220

NODE=M005W1
NODE=M005W1

NODE=M005W1;LINKAGE=L
NODE=M005W1;LINKAGE=SC

NODE=M005W4
NODE=M005W4

NODE=M005W4;LINKAGE=L
NODE=M005W4;LINKAGE=SC

NODE=M005W7
NODE=M005W7

NODE=M005W7;LINKAGE=L
NODE=M005W7;LINKAGE=SC

$\Gamma(\gamma\gamma)$

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an S-wave contribution, values are systematically higher (typically around 3 keV).

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR FIT				Error includes scale factor of 1.4.
2.93±0.40		¹ DAI	14A RVUE	Compilation
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.14±0.20		2,3 PENNINGTON 08	RVUE	Compilation
3.82±0.30		3,4 PENNINGTON 08	RVUE	Compilation
2.55±0.15	870	5 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
2.84±0.35		BOGLIONE 99	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
2.93±0.23±0.32		6 YABUKI 95	VNS	
2.58±0.13 ^{+0.36} _{-0.27}		7 BEHREND 92	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.10±0.35±0.35		8 BLINOV 92	MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.27±0.47±0.11		ADACHI 90D	TOPZ	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.15±0.04±0.39		BOYER 90	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.19±0.16 ^{+0.29} _{-0.28}		MARSISKE 90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2.35±0.65		9 MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
3.19±0.09 ^{+0.22} _{-0.38}	2177	OEST 90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
3.2 ± 0.1 ± 0.4		10 AIHARA 86B	TPC	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.5 ± 0.1 ± 0.5		BEHREND 84B	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.85±0.25±0.5		11 BERGER 84	PLUT	$e^+ e^- \rightarrow e^+ e^- 2\pi$
2.70±0.05±0.20		COURAU 84	DLCO	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.52±0.13±0.38		12 SMITH 84C	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.7 ± 0.2 ± 0.6		EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
2.9 ^{+0.6} _{-0.4} ± 0.6		13 EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
3.2 ± 0.2 ± 0.6		BRANDELIK 81B	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.6 ± 0.3 ± 0.5		ROUSSARIE 81	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.3 ± 0.8		14 BERGER 80B	PLUT	$e^+ e^-$

1 Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.

2 Solution A (preferred solution based on χ^2 -analysis).

3 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

4 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

5 From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

6 With a narrow scalar state around 1220 MeV.

7 Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.

8 Using the unitarized model of LYTH 85.

9 Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370)$: $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.

10 Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.

11 Using the MENNESSIER 83 model.

12 Superseded by BOYER 90.

13 If helicity = 2 assumption is not made.

14 Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

 Γ_7

NODE=M005W8

NODE=M005W8

NODE=M005W8

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M005W8;LINKAGE=A

NODE=M005W8;LINKAGE=P1

NODE=M005W8;LINKAGE=P3

NODE=M005W8;LINKAGE=P2

NODE=M005W8;LINKAGE=SC

NODE=M005W8;LINKAGE=YA

NODE=M005W;LINKAGE=B

NODE=M005W;LINKAGE=A

NODE=M005PW;LINKAGE=C

NODE=M005PW;LINKAGE=B

NODE=M005PW;LINKAGE=X

NODE=M005PW;LINKAGE=V

NODE=M005PW;LINKAGE=H

NODE=M005PW;LINKAGE=A

NODE=M005W9

NODE=M005W9

 $\Gamma(e^+ e^-)$ Γ_{10}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.11	90	ACHASOV 00K	SND	$e^+ e^- \rightarrow \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.7	90	VOROBIEV 88	ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_7/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.121±0.020 OUR FIT	Error includes scale factor of 1.3.		
0.091±0.007±0.027	¹ ALBRECHT	90G ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.104±0.007±0.072	² ALBRECHT	90G ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
1 Using an incoherent background.			
2 Using a coherent background.			

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_5\Gamma_7/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
11.5^{+1.8+4.5}_{-2.0-3.7}	¹ UEHARA	10A BELL	$10.6 e^+e^- \rightarrow e^+e^-\eta\eta$
1 Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_0(Y)$.			

Helicity-0/Helicity-2 RATIO IN $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
3.7±0.3^{+15.9}_{-2.9}	UEHARA	08A BELL	$10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9.5±1.8	¹ DAI	14A RVUE	Compilation
13	2,3 PENNINGTON 08	RVUE	Compilation
26	3,4 PENNINGTON 08	RVUE	Compilation
1 Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.			
2 Solution A (preferred solution based on χ^2 -analysis).			
3 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.			
4 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).			

 $f_2(1270)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

0.843^{+0.028}_{-0.010} OUR FIT Error includes scale factor of 1.2.

0.837±0.020 OUR AVERAGE

0.849±0.025	CHABAUD	83	ASPK	17 $\pi^- p$ polarized
0.85 ± 0.05	250	BEAUPRE	71	HBC $8 \pi^+ p \rightarrow \Delta^{++} f_2$
0.8 ± 0.04	600	OH	70	HBC $1.26 \pi^- p \rightarrow \pi^+\pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.856±0.001±0.05	¹ ALBRECHT	20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi\pi)$

Should be twice $\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.091^{+0.015}_{-0.040} OUR FIT Error includes scale factor of 1.2.

0.15 ± 0.06 600 EISENBERG 74 HBC $4.9 \pi^+ p \rightarrow \Delta^{++} f_2$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 EMMS 75D DBC $4 \pi^+ n \rightarrow p f_2$

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.033±0.001±0.005	¹ ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M005223

NODE=M005G1

NODE=M005G1

OCCUR=2

NODE=M005G1;LINKAGE=A

NODE=M005G1;LINKAGE=K

NODE=M005G02

NODE=M005G02

NODE=M005G02;LINKAGE=UE

NODE=M005HR0

NODE=M005HR0

OCCUR=2

OCCUR=3

NODE=M005HR0;LINKAGE=A

NODE=M005HR0;LINKAGE=P1

NODE=M005HR0;LINKAGE=P3

NODE=M005HR0;LINKAGE=P2

NODE=M005225

NODE=M005R10

NODE=M005R10

NODE=M005R10;LINKAGE=A

NODE=M005R2

NODE=M005R2

NODE=M005R2

NODE=M005R00

NODE=M005R00

NODE=M005R00;LINKAGE=A

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.054^{+0.005}_{-0.006} OUR FIT Error includes scale factor of 2.7.

0.041^{+0.004}_{-0.005} OUR AVERAGE

0.045 \pm 0.01		¹ BARGIOTTI	03	OBLX $\bar{p}p$
0.037 ^{+0.008} _{-0.021}		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n2K_S^0$
0.045 \pm 0.009		CHABAUD	81	ASPK 17 $\pi^- p$ polarized
0.039 \pm 0.008		LOVERRE	80	HBC 4 $\pi^- p \rightarrow K\bar{K}N$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.052 \pm 0.025		ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
0.036 \pm 0.005		² COSTA	80	OMEG 1-2.2 $\pi^- p \rightarrow K^+ K^- n$
0.030 \pm 0.005		³ MARTIN	79	RVUE
0.027 \pm 0.009		⁴ POLYCHRO...	79	STRC 7 $\pi^- p \rightarrow n2K_S^0$
0.025 \pm 0.015		EMMS	75D	DBC 4 $\pi^+ n \rightarrow pf_2$
0.031 \pm 0.012	20	ADERHOLZ	69	HBC 8 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² Re-evaluated by CHABAUD 83.

³ Includes PAWLICKI 77 data.

⁴ Takes into account the $f_2(1270)$ - $f'_2(1525)$ interference.

NODE=M005R3

NODE=M005R3

NODE=M005R3

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.033 \pm 0.005 OUR FIT Error includes scale factor of 1.2.

0.033 \pm 0.004 OUR AVERAGE Error includes scale factor of 1.1.

0.024 \pm 0.006	160	EMMS	75D	DBC 4 $\pi^+ n \rightarrow pf_2$
0.051 \pm 0.025	70	EISENBERG	74	HBC 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.043 ^{+0.007} _{-0.011}	285	¹ LOUIE	74	HBC 3.9 $\pi^- p \rightarrow nf_2$
0.037 \pm 0.007	154	ANDERSON	73	DBC 6 $\pi^+ n \rightarrow pf_2$
0.047 \pm 0.013		OH	70	HBC 1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

¹ LOUIE 74 was quoted as 0.065 in PDG 74. Factor 2/3 to go from $\pi^+ \pi^- \rightarrow \pi\pi$ forgotten. Mike L.

NODE=M005R;LINKAGE=BG

NODE=M005R3;LINKAGE=D

NODE=M005R3;LINKAGE=F

NODE=M005R3;LINKAGE=M

NODE=M005R1

NODE=M005R1

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

4.0 \pm 0.8 OUR FIT Error includes scale factor of 2.1.

2.9 \pm 0.5 OUR AVERAGE

2.7 \pm 0.7		BINON	05	GAMS 33 $\pi^- p \rightarrow \eta\eta n$
2.8 \pm 0.7		ALDE	86D	GAM4 100 $\pi^- p \rightarrow 2\eta n$
5.2 \pm 1.7		BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
4.0 \pm 1.0 \pm 2.0		¹ ALBRECHT	20	RVUE 0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$, $\pi^0 \eta\eta$, $\pi^0 K^+ K^-$

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M005R7

NODE=M005R7

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.003 \pm 0.001

BARBERIS 00E 450 $p p \rightarrow p_f \eta\eta p_s$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<0.05	95	EDWARDS	82F	CBAL $e^+ e^- \rightarrow e^+ e^- 2\eta$
<0.016	95	EMMS	75D	DBC 4 $\pi^+ n \rightarrow pf_2$
<0.09	95	EISENBERG	74	HBC 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$

NODE=M005R6

NODE=M005R6

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0030 \pm 0.0010 OUR FIT

0.003 ± 0.001 400 ± 50 ALDE 87 GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

NODE=M005R11

NODE=M005R11

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_7/Γ				
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	NODE=M005R13	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.57 \pm 0.01^{+1.39}_{-0.14}$	UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	NODE=M005R13	
$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$	Γ_8/Γ_1				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	NODE=M005R5
<0.010	95	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$	NODE=M005R5
$\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$	Γ_9/Γ_1				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	NODE=M005R4
<0.004	95	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$	NODE=M005R4
$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$	Γ_{10}/Γ				
VALUE (units 10^{-10})	CL%	DOCUMENT ID	TECN	COMMENT	NODE=M005R12
<6	90	ACHASOV	00K SND	$e^+ e^- \rightarrow \pi^0 \pi^0$	NODE=M005R12

$f_2(1270)$ REFERENCES

KLEMPPT	22	PL B830 137171	E. Klempert et al.	(BONN)	REFID=61646
RODAS	22	EPJ C82 80	A. Rodas et al.	(JPAC Collab.)	REFID=61610
CARVER	21	PRL 126 082002	M. Carver et al.	(CLAS Collab.)	REFID=61097
ALBRECHT	20	EPJ C80 453	M. Albrecht et al.	(Crystal Barrel Collab.)	REFID=60439
DOBBS	15	PR D91 052006	S. Dobbs et al.	(NWES)	REFID=56805
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)	REFID=55923
BOGOLYUBOV	13	PAN 76 1324	M.Yu. Bogolyubsky et al.	(HYPERON-M Collab.)	REFID=55585
		Translated from YAF 76 1389.			
UEHARA	13	PTEP 2013 123C01	S. Uehara et al.	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara et al.	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara et al.	(BELLE Collab.)	REFID=52761
PDG	08	PL B667 1	C. Amsler et al.	(PDG Collab.)	REFID=52166
PENNINGTON	08	EPJ C56 1	M.R. Pennington et al.	(BELLE Collab.)	REFID=52303
UEHARA	08A	PR D78 052004	S. Uehara et al.	(BELLE Collab.)	REFID=52309
MORI	07	PR D75 051101	T. Mori et al.	(BELLE Collab.)	REFID=51652
ABLIKIM	06V	PL B642 441	M. Ablikim et al.	(BES Collab.)	REFID=51507
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky et al.	(BES Collab.)	REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim et al.	(BES Collab.)	REFID=50450
BINON	05	PAN 68 960	F. Binon et al.	(BES Collab.)	REFID=50780
		Translated from YAF 68 998.			
ABLIKIM	04E	PL B603 138	M. Ablikim et al.	(BES Collab.)	REFID=50174
BARGIOTTI	03	EPJ C26 371	M. Bargiotti et al.	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov et al.		REFID=49423
		Translated from YAF 66 860.			
ACHASOV	00K	PL B492 8	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00E	PL B479 59	D. Barberis et al.	(WA 102 Collab.)	REFID=47961
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
ALDE	98	EPJ A3 361	D. Alde et al.	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde et al.	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ALDE	97	PL B397 350	D.M. Alde et al.	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin et al.	(OBELIX Collab.)	REFID=45701
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)	REFID=45566
		Translated from YAF 59 2187.			
YABUKI	95	JPSJ 64 435	F. Yabuki et al.	(VENUS Collab.)	REFID=46384
PROKOSHKIN	94	PD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)	REFID=44094
		Translated from DANS 336 613.			
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
BLINOV	92	ZPHY C53 33	A.E. Blinov et al.	(NOVO)	REFID=41858
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez et al.	(LEBC-EHS Collab.)	REFID=41637
AKER	91	PL B260 249	E. Aker et al.	(Crystal Barrel Collab.)	REFID=41587
ADACHI	90D	PL B234 185	I. Adachi et al.	(TOPAZ Collab.)	REFID=41345
ALBRECHT	90G	ZPHY C48 183	H. Albrecht et al.	(ARGUS Collab.)	REFID=41374
BOYER	90	PR D42 1350	J. Boyer et al.	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone et al.	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske et al.	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest et al.	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev et al.	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
ALDE	87	PL B198 286	D.M. Alde et al.	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin et al.	(LALO, CLER, FRAS+)	REFID=40268
ABACHI	86B	PRL 57 1990	S. Abachi et al.	(PURD, ANL, IND, MICH+)	REFID=20394
AIHARA	86B	PR D57 404	H. Aihara et al.	(TPC-2 γ Collab.)	REFID=20764
ALDE	86D	NP B269 485	D.M. Alde et al.	(BELG, LAPP, SERP, CERN+)	REFID=20765
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)	REFID=20767
LONGACRE	86	PL B177 223	R.S. Longacre et al.	(BNL, BRAN, CUNY+)	REFID=20768
LYTH	85	JP G11 459	D.H. Lyth		REFID=42169
BEHREND	84B	ZPHY C23 223	H.J. Behrend et al.	(CELLO Collab.)	REFID=20757
BERGER	84	ZPHY C26 199	C. Berger et al.	(PLUTO Collab.)	REFID=20760
COURAU	84	PL 147B 227	A. Courau et al.	(CIT, SLAC)	REFID=20758
SMITH	84C	PR D30 851	J.R. Smith et al.	(SLAC, LBL, HARV)	REFID=20759
BINON	83	NC 78A 313	F.G. Binon et al.	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon et al.	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934.			

CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)	REFID=20393
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20745
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=20741
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)	REFID=20388
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20736
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)	REFID=20382
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>		REFID=40124
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)	REFID=20367
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)	REFID=20721
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=20715
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)	REFID=20719
PDG	74	PL 50B 1	V. Chaloupka <i>et al.</i>		REFID=40125
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)	REFID=20710
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)	REFID=20103
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)	REFID=20698
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)	REFID=20700
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP	REFID=20335
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)	REFID=20681
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)	REFID=20065
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)	REFID=20668
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)	REFID=20663
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)	REFID=20657