

NODE=M030

$h_1(1170)$

$I^G(J^{PC}) = 0^-(1^{+-})$

$h_1(1170)$ MASS

NODE=M030M

NODE=M030M
OCCUR=2

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1166± 5±3	¹ ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1168± 4	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1190±60	² DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
¹ Average and spread of values using 2 variants of the model of BOWLER 75.				
² Uses the model of BOWLER 75.				

NODE=M030M;LINKAGE=B
NODE=M030M;LINKAGE=C

$h_1(1170)$ WIDTH

NODE=M030W

NODE=M030W
OCCUR=2

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
375± 6±34	³ ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
345± 6	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
320±50	⁴ DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
³ Average and spread of values using 2 variants of the model of BOWLER 75.				
⁴ Uses the model of BOWLER 75.				

NODE=M030W;LINKAGE=B
NODE=M030W;LINKAGE=C

$h_1(1170)$ DECAY MODES

NODE=M030215;NODE=M030

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \rho \pi$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

$h_1(1170)$ BRANCHING RATIOS

NODE=M030220

NODE=M030R1
NODE=M030R1

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
seen	ATKINSON	84	OMEG	20-70 $\gamma p \rightarrow \pi^+ \pi^- \pi^0 p$
seen	DANKOWY...	81	SPEC	8 $\pi p \rightarrow 3\pi n$

$h_1(1170)$ REFERENCES

NODE=M030

REFID=43171
REFID=20574
REFID=20572
REFID=20571

ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)