

$J/\psi(1S)$ $I^G(J^{PC}) = 0^-(1^-^-)$ **$J/\psi(1S)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.900 ± 0.006 OUR AVERAGE				
3096.900 $\pm 0.002 \pm 0.006$		¹ ANASHIN 15	KEDR	$e^+ e^- \rightarrow$ hadrons
3096.89 ± 0.09	502	² ARTAMONOV 00	OLYA	$e^+ e^- \rightarrow$ hadrons
3096.91 $\pm 0.03 \pm 0.01$		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
3096.95 $\pm 0.1 \pm 0.3$	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc.				
3096.66 $\pm 0.19 \pm 0.02$	6.1k	⁴ AAIJ 15BI	LHCb	$p p \rightarrow J/\psi X$
3096.917 $\pm 0.010 \pm 0.007$		AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
3097.5 ± 0.3		GRIBUSHIN 96	FMPS	$515 \pi^- Be \rightarrow 2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE 82	GOLI	$185 \pi^- Be \rightarrow \gamma \mu^+ \mu^- A$
3096.93 ± 0.09	502	⁵ ZHOLENTZ 80	REDE	$e^+ e^-$
3097.0 ± 1		⁶ BRANDELIK 79C	DASP	$e^+ e^-$

1 Supersedes AULCHENKO 03.

2 Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

3 Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.4 From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.

5 Superseded by ARTAMONOV 00.

6 From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$ and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

NODE=M070M

NODE=M070M

 $J/\psi(1S)$ WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.6 ± 1.7 OUR AVERAGE				
92.45 $\pm 1.40 \pm 1.48$		Error includes scale factor of 1.1.		
96.1 ± 3.2	13k	¹ ANASHIN 20	KEDR	$e^+ e^-$
84.4 ± 8.9		² ADAMS 06A	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
91 $\pm 11 \pm 6$		BAI 95B	BES	$e^+ e^-$
85.5 ± 6.1 $- 5.8$		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
92.94 ± 1.83		^{5,6} ANASHIN 18A	KEDR	$e^+ e^-$
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	$3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$

1 Based on the same dataset as ANASHIN 18A and correlated to the values reported there.

2 Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(e^+ e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.

3 The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

4 Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

5 Using $\Gamma(e^+ e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ from PDG 16.

6 Superseded by ANASHIN 20 that is based on the same dataset.

7 Assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$ and using $\Gamma(e^+ e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.NODE=M070M;LINKAGE=A
NODE=M070M;LINKAGE=AR

NODE=M070M;LINKAGE=NW

NODE=M070M;LINKAGE=B

NODE=M070M;LINKAGE=RZ

NODE=M070M;LINKAGE=F

NODE=M070W

NODE=M070W

 $J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.46 \pm 0.07) %	
Γ_3 $g g g$	(64.1 \pm 1.0) %	
Γ_4 $\gamma g g$	(8.8 \pm 1.1) %	
Γ_5 $e^+ e^-$	(5.971 \pm 0.032) %	
Γ_6 $e^+ e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+ \mu^-$	(5.961 \pm 0.033) %	
Γ_8 $e^+ e^- e^+ e^-$	(5.5 \pm 0.5) $\times 10^{-5}$	
Γ_9 $e^+ e^- \mu^+ \mu^-$	(3.53 \pm 0.26) $\times 10^{-5}$	
Γ_{10} $\mu^+ \mu^- \mu^+ \mu^-$	(1.11 \pm 0.11) $\times 10^{-6}$	

DESIG=3

DESIG=4

DESIG=249

DESIG=250

DESIG=1

DESIG=5

DESIG=2

DESIG=472

DESIG=473

DESIG=471

NODE=M070215;NODE=M070

Decays involving hadronic resonances

Γ_{11}	$\rho\pi$	(1.88 \pm 0.12) %	S=2.6	NODE=M070;CLUMP=A
Γ_{12}	$\rho^0\pi^0$	(6.2 \pm 0.6) $\times 10^{-3}$		DESIG=20
Γ_{13}	$a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0$	(2.8 \pm 0.6) $\times 10^{-3}$		DESIG=21
Γ_{14}	$a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow 2(\pi^+\pi^-)\pi^0$	(3.7 \pm 0.7) $\times 10^{-3}$		DESIG=442
Γ_{15}	$a_2(1320)\rho$	(1.09 \pm 0.22) %		DESIG=443
Γ_{16}	$\eta\pi^+\pi^-$	(3.8 \pm 0.7) $\times 10^{-4}$		DESIG=239
Γ_{17}	$\eta\rho$	(1.93 \pm 0.23) $\times 10^{-4}$		DESIG=22
Γ_{18}	$\eta\pi^+\pi^-\pi^0$	(1.17 \pm 0.20) %		DESIG=420
Γ_{19}	$\eta\pi^+\pi^-3\pi^0$	(4.9 \pm 1.0) $\times 10^{-3}$		DESIG=422
Γ_{20}	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	(1.2 \pm 0.4) $\times 10^{-4}$		DESIG=287
Γ_{21}	$\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0$	< 2.52 $\times 10^{-4}$ CL=90%		DESIG=253
Γ_{22}	ηK^+K^-	(8.6 \pm 3.0) $\times 10^{-4}$		DESIG=455
Γ_{23}	$\eta K^\pm K_S^0\pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$		DESIG=230
Γ_{24}	$\eta K^*(892)^0 \bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$		DESIG=252
Γ_{25}	$\rho\eta'(958)$	(8.1 \pm 0.8) $\times 10^{-5}$	S=1.6	DESIG=23
Γ_{26}	$\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0$	(2.8 \pm 0.8) %		DESIG=415
Γ_{27}	$\rho^+\rho^-\pi^+\pi^-\pi^0$	(6 \pm 4) $\times 10^{-3}$		DESIG=416
Γ_{28}	$\rho^+K^+K^-\pi^- + \text{c.c.} \rightarrow K^+K^-\pi^+\pi^-\pi^0$	(3.5 \pm 0.8) $\times 10^{-3}$		DESIG=444
Γ_{29}	$\rho^\mp K^\pm K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$		DESIG=342
Γ_{30}	$h_1(1415)\eta' \rightarrow \gamma\eta\eta'$			DESIG=435
Γ_{31}	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$			DESIG=437
Γ_{32}	$\rho(1450)\pi$	seen		DESIG=310;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{33}	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	(2.2 \pm 1.1) $\times 10^{-4}$		DESIG=328
Γ_{34}	$\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp$	(3.3 \pm 0.6) $\times 10^{-4}$		DESIG=329
Γ_{35}	$\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0$	(2.7 \pm 0.6) $\times 10^{-4}$		DESIG=312
Γ_{36}	$\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958)$	(3.3 \pm 0.7) $\times 10^{-6}$		DESIG=345
Γ_{37}	$\rho(1700)\pi$	seen		DESIG=325;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{38}	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	(1.6 \pm 1.1) $\times 10^{-4}$		DESIG=313
Γ_{39}	$\rho(2150)\pi$	seen		DESIG=326;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{40}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	(10 \pm 40) $\times 10^{-6}$		DESIG=314
Γ_{41}	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$			DESIG=316
Γ_{42}	$\omega\pi^0$	(4.5 \pm 0.5) $\times 10^{-4}$	S=1.4	DESIG=32
Γ_{43}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	(1.6 \pm 0.7) $\times 10^{-5}$		DESIG=327
Γ_{44}	$\omega\pi^+\pi^-$	(8.5 \pm 1.0) $\times 10^{-3}$	S=1.3	DESIG=24
Γ_{45}	$\omega\pi^0\pi^0$	(3.4 \pm 0.8) $\times 10^{-3}$		DESIG=140
Γ_{46}	$\omega 3\pi^0$	(1.9 \pm 0.6) $\times 10^{-3}$		DESIG=421
Γ_{47}	$\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$		DESIG=28
Γ_{48}	$\omega\eta$	(1.74 \pm 0.20) $\times 10^{-3}$	S=1.6	DESIG=30
Γ_{49}	$\omega\pi^+\pi^-\pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$		DESIG=211
Γ_{50}	$\omega\pi^0\eta$	(3.4 \pm 1.7) $\times 10^{-4}$		DESIG=360
Γ_{51}	$\omega\pi^+\pi^-\pi^-\pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$		DESIG=26
Γ_{52}	$\omega\pi^+\pi^-2\pi^0$	(3.3 \pm 0.5) %		DESIG=412
Γ_{53}	$\omega\eta'\pi^+\pi^-$	(1.12 \pm 0.13) $\times 10^{-3}$		DESIG=385
Γ_{54}	$\omega\eta'(958)$	(1.89 \pm 0.18) $\times 10^{-4}$		DESIG=31
Γ_{55}	$\omega f_0(980)$	(1.4 \pm 0.5) $\times 10^{-4}$		DESIG=150
Γ_{56}	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	(4.8 \pm 1.1) $\times 10^{-4}$		DESIG=130
Γ_{57}	$\omega f_1(1420)$	(6.8 \pm 2.4) $\times 10^{-4}$		DESIG=105
Γ_{58}	$\omega f'_2(1525)$	< 2.2 $\times 10^{-4}$ CL=90%		DESIG=29
Γ_{59}	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9 $\times 10^{-6}$ CL=95%		DESIG=263
Γ_{60}	$\omega K^+K^-\eta$	(3.33 \pm 0.12) $\times 10^{-4}$		DESIG=478
Γ_{61}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	< 6.2 $\times 10^{-5}$		DESIG=386
Γ_{62}	ωK^+K^-	(1.52 \pm 0.31) $\times 10^{-3}$		DESIG=441
Γ_{63}	$\omega K^\pm K_S^0\pi^\mp$	[b] (3.4 \pm 0.5) $\times 10^{-3}$		DESIG=101
Γ_{64}	$\omega K\bar{K}$	(1.9 \pm 0.4) $\times 10^{-3}$		DESIG=27

Γ_{65}	$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	DESIG=102	
Γ_{66}	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$	DESIG=355	
Γ_{67}	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$	DESIG=357	
Γ_{68}	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$	DESIG=353	
Γ_{69}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$	DESIG=354	
Γ_{70}	$\eta' h_1(1415) \rightarrow \gamma \eta' \eta'$	$(4.7 \pm 1.1) \times 10^{-7}$	DESIG=430	
Γ_{71}	$\bar{K} K^*(892) + \text{c.c.}$	seen	DESIG=331; OUR EST; → UNCHECKED ←	
Γ_{72}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(4.8 \pm 0.5) \times 10^{-3}$	DESIG=332	
Γ_{73}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm 0.8) \times 10^{-3}$	S=2.9	DESIG=121
Γ_{74}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.13) \times 10^{-3}$	DESIG=231	
Γ_{75}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	DESIG=232	
Γ_{76}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	DESIG=122	
Γ_{77}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	DESIG=233	
Γ_{78}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$	DESIG=214	
Γ_{79}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	DESIG=343	
Γ_{80}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$	DESIG=299	
Γ_{81}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$	DESIG=300	
Γ_{82}	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$	DESIG=445	
Γ_{83}	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm 0.6) \times 10^{-6}$	DESIG=376	
Γ_{84}	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$	DESIG=344	
Γ_{85}	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm 1.0) \times 10^{-3}$	DESIG=257	
Γ_{86}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$	DESIG=46	
Γ_{87}	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm 0.22) \times 10^{-3}$	DESIG=256	
Γ_{88}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	DESIG=132	
Γ_{89}	$K^*(1410) \bar{K} + \text{c.c.}$	seen	DESIG=317; OUR EST; → UNCHECKED ←	
Γ_{90}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$	DESIG=330	
Γ_{91}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(8 \pm 5) \times 10^{-5}$	DESIG=318	
Γ_{92}	$K_2^*(1430) \bar{K} + \text{c.c.}$	seen	DESIG=319; OUR EST; → UNCHECKED ←	
Γ_{93}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$	DESIG=321	
Γ_{94}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.8 \pm 1.0) \times 10^{-4}$	DESIG=320	
Γ_{95}	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%	DESIG=45
Γ_{96}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.25) \times 10^{-4}$	DESIG=381	
Γ_{97}	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$	DESIG=446	
Γ_{98}	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	DESIG=301	
Γ_{99}	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$	DESIG=48	
Γ_{100}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	DESIG=303	
Γ_{101}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$	DESIG=304	
Γ_{102}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%	DESIG=47
Γ_{103}	$\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	DESIG=235	
Γ_{104}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(1.10 \pm 0.60) \times 10^{-5}$	DESIG=382	

Γ_{105}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	(6.2 \pm 2.9) $\times 10^{-6}$	DESIG=383
Γ_{106}	$K_1(1270)^\pm K^\mp$	< 3.0 $\times 10^{-3}$ CL=90%	DESIG=131
Γ_{107}	$K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0$	(8.5 \pm 2.5) $\times 10^{-7}$	DESIG=377
Γ_{108}	$a_2(1320)^\pm \pi^\mp$	[b] < 4.3 $\times 10^{-3}$ CL=90%	DESIG=42
Γ_{109}	$\phi \pi^0$	3 $\times 10^{-6}$ or 1 $\times 10^{-7}$	DESIG=33; OUR EVAL; → UNCHECKED ←
Γ_{110}	$\phi \pi^+ \pi^-$	(9.4 \pm 1.5) $\times 10^{-4}$ S=1.7	DESIG=34
Γ_{111}	$\phi \pi^0 \pi^0$	(4.9 \pm 1.0) $\times 10^{-4}$	DESIG=76
Γ_{112}	$\phi 2(\pi^+ \pi^-)$	(1.60 \pm 0.32) $\times 10^{-3}$	DESIG=35
Γ_{113}	$\phi \eta$	(7.4 \pm 0.6) $\times 10^{-4}$ S=1.2	DESIG=37
Γ_{114}	$\phi \eta'(958)$	(4.6 \pm 0.5) $\times 10^{-4}$ S=2.2	DESIG=38
Γ_{115}	$\phi \eta \eta'$	(2.32 \pm 0.17) $\times 10^{-4}$	DESIG=387
Γ_{116}	$\phi f_0(980)$	(3.2 \pm 0.9) $\times 10^{-4}$ S=1.9	DESIG=41
Γ_{117}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	(2.58 \pm 0.34) $\times 10^{-4}$	DESIG=236
Γ_{118}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	(1.7 \pm 0.5) $\times 10^{-4}$	DESIG=237
Γ_{119}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	(4.5 \pm 1.0) $\times 10^{-6}$	DESIG=278
Γ_{120}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 p^0 \pi^0$	(1.7 \pm 0.6) $\times 10^{-6}$	DESIG=279
Γ_{121}	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	(3.2 \pm 1.0) $\times 10^{-4}$	DESIG=229
Γ_{122}	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$		DESIG=258
Γ_{123}	$\phi(1680)^0 \pi^0 \rightarrow \phi \eta \pi^0$	(6.7 \pm 1.1) $\times 10^{-6}$	DESIG=480
Γ_{124}	$X(2000)^0 \pi^0 \rightarrow \phi \eta \pi^0$	(1.70 \pm 0.50) $\times 10^{-6}$	DESIG=481
Γ_{125}	$h_1(1900)^0 \pi^0 \rightarrow \phi \eta \pi^0$	(8.4 \pm 1.4) $\times 10^{-6}$	DESIG=482
Γ_{126}	$\phi f_2(1270)$	(3.2 \pm 0.6) $\times 10^{-4}$	DESIG=39
Γ_{127}	$\phi f_1(1285)$	(2.6 \pm 0.5) $\times 10^{-4}$	DESIG=106
Γ_{128}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	(9.4 \pm 2.8) $\times 10^{-7}$	DESIG=280
Γ_{129}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0$	(2.1 \pm 2.2) $\times 10^{-7}$	DESIG=281
Γ_{130}	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	(2.0 \pm 1.0) $\times 10^{-5}$	DESIG=128
Γ_{131}	$\phi f'_2(1525)$	(8 \pm 4) $\times 10^{-4}$ S=2.7	DESIG=40
Γ_{132}	$\phi X(1835) \rightarrow \phi p \bar{p}$	< 2.1 $\times 10^{-7}$ CL=90%	DESIG=291
Γ_{133}	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	< 2.8 $\times 10^{-4}$ CL=90%	DESIG=288
Γ_{134}	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	< 6.13 $\times 10^{-5}$ CL=90%	DESIG=289
Γ_{135}	$\phi K \bar{K}$	(1.77 \pm 0.16) $\times 10^{-3}$ S=1.3	DESIG=36
Γ_{136}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	(3.6 \pm 0.6) $\times 10^{-4}$	DESIG=129
Γ_{137}	$\phi K^+ K^-$	(8.2 \pm 1.1) $\times 10^{-4}$	DESIG=295
Γ_{138}	$\phi K_S^0 K_S^0$	(5.8 \pm 1.5) $\times 10^{-4}$	DESIG=305
Γ_{139}	$\phi K^\pm K_S^0 \pi^\mp$	[b] (7.2 \pm 0.8) $\times 10^{-4}$	DESIG=103
Γ_{140}	$\phi K^*(892) \bar{K} + \text{c.c.}$	(2.18 \pm 0.23) $\times 10^{-3}$	DESIG=104
Γ_{141}	$b_1(1235)^\pm \pi^\mp$	[b] (3.0 \pm 0.5) $\times 10^{-3}$	DESIG=49
Γ_{142}	$b_1(1235)^0 \pi^0$	(2.3 \pm 0.6) $\times 10^{-3}$	DESIG=160
Γ_{143}	$f'_2(1525) K^+ K^-$	(1.04 \pm 0.35) $\times 10^{-3}$	DESIG=308
Γ_{144}	$\Delta(1232)^+ \bar{p}$	< 1 $\times 10^{-4}$ CL=90%	DESIG=112
Γ_{145}	$\Delta(1232)^{++} \bar{p} \pi^-$	(1.6 \pm 0.5) $\times 10^{-3}$	DESIG=70
Γ_{146}	$\Delta(1232)^{++} \Delta(1232)^{--}$	(1.10 \pm 0.29) $\times 10^{-3}$	DESIG=66
Γ_{147}	$\Sigma(1385)^0 p K^-$	(5.1 \pm 3.2) $\times 10^{-4}$	DESIG=74
Γ_{148}	$\Sigma(1385)^0 \Lambda + \text{c.c.}$	< 8.2 $\times 10^{-6}$ CL=90%	DESIG=111
Γ_{149}	$\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.}$	[b] (3.0 \pm 0.7) $\times 10^{-4}$	DESIG=68
Γ_{150}	$\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.}$	(3.3 \pm 0.8) $\times 10^{-4}$	DESIG=450
Γ_{151}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.}$	[b] (1.08 \pm 0.06) $\times 10^{-3}$	DESIG=67
Γ_{152}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.}$	(1.25 \pm 0.07) $\times 10^{-3}$	DESIG=451
Γ_{153}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	(1.07 \pm 0.08) $\times 10^{-3}$	DESIG=309
Γ_{154}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	< 4.1 $\times 10^{-6}$ CL=90%	DESIG=260
Γ_{155}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	< 1.80 $\times 10^{-3}$ CL=90%	DESIG=364
Γ_{156}	$\Xi^0 \bar{\Xi}^0$	(1.17 \pm 0.04) $\times 10^{-3}$	DESIG=248
Γ_{157}	$\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.}$	(3.18 \pm 0.08) $\times 10^{-4}$	DESIG=107
Γ_{158}	$\Xi(1530)^0 \bar{\Xi}^0$	(3.2 \pm 1.4) $\times 10^{-4}$	DESIG=108
Γ_{159}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	[c] < 1.1 $\times 10^{-5}$ CL=90%	DESIG=205
Γ_{160}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[c] < 2.1 $\times 10^{-5}$ CL=90%	DESIG=206
Γ_{161}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[c] < 1.6 $\times 10^{-5}$ CL=90%	DESIG=207
Γ_{162}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[c] < 5.6 $\times 10^{-5}$ CL=90%	DESIG=208
Γ_{163}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[c] < 1.1 $\times 10^{-5}$ CL=90%	DESIG=209

Decays into stable hadrons

			NODE=M070;CLUMP=B
Γ_{164}	$2(\pi^+\pi^-)\pi^0$	(4.2 \pm 0.4) %	S=2.1 DESIG=9
Γ_{165}	$3(\pi^+\pi^-)\pi^0$	(2.9 \pm 0.6) %	DESIG=11
Γ_{166}	$\pi^+\pi^-3\pi^0$	(1.9 \pm 0.9) %	DESIG=358
Γ_{167}	$\rho^\pm\pi^\mp\pi^0\pi^0$	(1.41 \pm 0.22) %	DESIG=362
Γ_{168}	$\rho^+\rho^-\pi^0$	(6.0 \pm 1.1) $\times 10^{-3}$	DESIG=363
Γ_{169}	$\pi^+\pi^-4\pi^0$	(6.5 \pm 1.3) $\times 10^{-3}$	DESIG=419
Γ_{170}	$\pi^+\pi^-\pi^0$	(2.00 \pm 0.07) %	S=2.0 DESIG=7
Γ_{171}	$2(\pi^+\pi^-\pi^0)$	(1.61 \pm 0.20) %	DESIG=210
Γ_{172}	$\pi^+\pi^-\pi^0K^+K^-$	(1.52 \pm 0.27) %	S=1.4 DESIG=18
Γ_{173}	$\pi^+\pi^-$	(1.47 \pm 0.14) $\times 10^{-4}$	DESIG=6
Γ_{174}	$2(\pi^+\pi^-)$	(3.20 \pm 0.25) $\times 10^{-3}$	S=1.2 DESIG=8
Γ_{175}	$3(\pi^+\pi^-)$	(4.3 \pm 0.4) $\times 10^{-3}$	DESIG=10
Γ_{176}	$2(\pi^+\pi^-)3\pi^0$	(6.2 \pm 0.9) %	DESIG=411
Γ_{177}	$4(\pi^+\pi^-)\pi^0$	(9.0 \pm 3.0) $\times 10^{-3}$	DESIG=12
Γ_{178}	$2(\pi^+\pi^-)\eta$	(2.29 \pm 0.28) $\times 10^{-3}$	DESIG=201
Γ_{179}	$3(\pi^+\pi^-)\eta$	(7.2 \pm 1.5) $\times 10^{-4}$	DESIG=202
Γ_{180}	$2(\pi^+\pi^-\pi^0)\eta$	(1.6 \pm 0.5) $\times 10^{-3}$	DESIG=418
Γ_{181}	$\pi^+\pi^-\pi^0\pi^0\eta$	(2.4 \pm 0.5) $\times 10^{-3}$	DESIG=359
Γ_{182}	$\rho^\pm\pi^\mp\pi^0\eta$	(1.9 \pm 0.8) $\times 10^{-3}$	DESIG=361
Γ_{183}	K^+K^-	(3.06 \pm 0.05) $\times 10^{-4}$	DESIG=13
Γ_{184}	$K_S^0K_L^0$	(1.95 \pm 0.11) $\times 10^{-4}$	S=2.4 DESIG=75
Γ_{185}	$K_S^0K_S^0$	< 1.4 $\times 10^{-8}$	CL=95% DESIG=14
Γ_{186}	$K\bar{K}\pi$	(6.1 \pm 1.0) $\times 10^{-3}$	DESIG=15
Γ_{187}	$K^+K^-\pi^0$	(2.88 \pm 0.12) $\times 10^{-3}$	DESIG=334
Γ_{188}	$K_S^0K^\pm\pi^\mp$	(5.3 \pm 0.5) $\times 10^{-3}$	DESIG=335
Γ_{189}	$K_S^0K_L^0\pi^0$	(2.06 \pm 0.26) $\times 10^{-3}$	DESIG=336
Γ_{190}	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	(1.21 \pm 0.18) $\times 10^{-3}$	DESIG=339
Γ_{191}	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	(4.3 \pm 1.3) $\times 10^{-4}$	DESIG=338
Γ_{192}	$K^+K^-\pi^+\pi^-$	(7.0 \pm 1.0) $\times 10^{-3}$	DESIG=16
Γ_{193}	$K^+K^-\pi^0\pi^0$	(2.13 \pm 0.22) $\times 10^{-3}$	DESIG=234
Γ_{194}	$K^+K^-\pi^0\pi^0\pi^0$	(1.61 \pm 0.29) $\times 10^{-3}$	DESIG=452
Γ_{195}	$K_S^0K^\pm\pi^\mp\pi^0\pi^0$	(5.3 \pm 0.7) $\times 10^{-3}$	DESIG=453
Γ_{196}	$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	(6.3 \pm 0.4) $\times 10^{-3}$	DESIG=454
Γ_{197}	$K_S^0K^\pm\rho(770)^\pm\pi^0$	(2.9 \pm 0.8) $\times 10^{-3}$	DESIG=463
Γ_{198}	$K_S^0K_L^0\pi^+\pi^-$	(3.8 \pm 0.6) $\times 10^{-3}$	DESIG=296
Γ_{199}	$K_S^0K_L^0\pi^0\pi^0$	(1.9 \pm 0.4) $\times 10^{-3}$	DESIG=337
Γ_{200}	$K_S^0K_L^0\eta$	(1.45 \pm 0.33) $\times 10^{-3}$	DESIG=340
Γ_{201}	$K_S^0K_S^0\pi^+\pi^-$	(1.68 \pm 0.19) $\times 10^{-3}$	DESIG=297
Γ_{202}	$K^\mp K_S^0\pi^\pm\pi^0$	(5.7 \pm 0.5) $\times 10^{-3}$	DESIG=341
Γ_{203}	$K_S^0K^\pm\pi^\mp\rho(770)^0$	(3.1 \pm 0.5) $\times 10^{-3}$	DESIG=456
Γ_{204}	$K^+K^-2(\pi^+\pi^-)$	(3.1 \pm 1.3) $\times 10^{-3}$	DESIG=17
Γ_{205}	$K^+K^-\pi^+\pi^-\eta$	(4.7 \pm 0.7) $\times 10^{-3}$	DESIG=238
Γ_{206}	$2(K^+K^-)$	(7.2 \pm 0.8) $\times 10^{-4}$	DESIG=19
Γ_{207}	$K^+K^-K_S^0K_S^0$	(4.2 \pm 0.7) $\times 10^{-4}$	DESIG=298
Γ_{208}	$K_S^0K^*(892)^0\pi^+\pi^-$	(1.7 \pm 0.6) $\times 10^{-3}$	DESIG=457
Γ_{209}	$K_S^0K^*(892)^0\pi^0\pi^0$	(1.01 \pm 0.18) $\times 10^{-3}$	DESIG=462
Γ_{210}	$K^\mp K^*(892)^\pm\pi^+\pi^-$	(3.4 \pm 1.2) $\times 10^{-3}$	DESIG=458
Γ_{211}	$K^*(892)^\pm K^*(892)^0\pi^\mp$	(4.8 \pm 1.0) $\times 10^{-3}$	DESIG=459
Γ_{212}	$K^\mp K^*(892)^\pm\pi^0\pi^0$	(1.57 \pm 0.32) $\times 10^{-3}$	DESIG=461
Γ_{213}	$K^*(892)^+K^*(892)^-\pi^0$	(1.12 \pm 0.23) %	DESIG=460
Γ_{214}	$p\bar{p}$	(2.120 \pm 0.029) $\times 10^{-3}$	DESIG=50
Γ_{215}	$p\bar{p}\pi^0$	(1.19 \pm 0.08) $\times 10^{-3}$	S=1.1 DESIG=52
Γ_{216}	$p\bar{p}\pi^+\pi^-$	(6.0 \pm 0.5) $\times 10^{-3}$	S=1.3 DESIG=54
Γ_{217}	$p\bar{p}\pi^+\pi^-\pi^0$	[d] (2.3 \pm 0.9) $\times 10^{-3}$	S=1.9 DESIG=55

Γ_{218}	$p\bar{p}\eta$	$(2.00 \pm 0.12) \times 10^{-3}$	DESIG=56
Γ_{219}	$p\bar{p}\rho$	$< 3.1 \times 10^{-4}$	CL=90% DESIG=57
Γ_{220}	$p\bar{p}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3 DESIG=58
Γ_{221}	$p\bar{p}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0 DESIG=59
Γ_{222}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$	DESIG=276
Γ_{223}	$p\bar{p}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	DESIG=127
Γ_{224}	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	DESIG=53
Γ_{225}	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	DESIG=64
Γ_{226}	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$	DESIG=65
Γ_{227}	$nN(1440)$	seen	DESIG=215;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{228}	$nN(1520)$	seen	DESIG=216;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{229}	$nN(1535)$	seen	DESIG=217;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{230}	$\Lambda\bar{\Lambda}$	$(1.88 \pm 0.08) \times 10^{-3}$	S=2.6 DESIG=60
Γ_{231}	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	DESIG=109
Γ_{232}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	DESIG=261
Γ_{233}	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$	DESIG=228
Γ_{234}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	[b] $(1.26 \pm 0.05) \times 10^{-3}$	S=1.2 DESIG=71
Γ_{235}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.21 \pm 0.07) \times 10^{-3}$	S=1.8 DESIG=449
Γ_{236}	$pK^-\bar{\Lambda} + \text{c.c.}$	$(8.6 \pm 1.1) \times 10^{-4}$	DESIG=72
Γ_{237}	$pK^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	DESIG=73
Γ_{238}	$pK_S^0\bar{\Sigma}^- + \text{c.c.}$	$(2.73 \pm 0.05) \times 10^{-4}$	DESIG=474
Γ_{239}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	DESIG=225
Γ_{240}	$\Lambda\bar{\Sigma}^+ + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	DESIG=61
Γ_{241}	$\Sigma^+\bar{\Sigma}^-$	$(1.07 \pm 0.04) \times 10^{-3}$	DESIG=247
Γ_{242}	$\Sigma^0\bar{\Sigma}^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4 DESIG=63
Γ_{243}	$\Sigma^+\bar{\Sigma}^-\eta$	$(6.3 \pm 0.4) \times 10^{-5}$	DESIG=448
Γ_{244}	$\Xi^-\bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4 DESIG=62

Radiative decays

Γ_{245}	$\gamma\eta_c(1S)$	$(1.41 \pm 0.14) \%$	S=1.3 NODE=M070;CLUMP=C
Γ_{246}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	seen	DESIG=85
Γ_{247}	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$	seen	DESIG=246;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{248}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	DESIG=391;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{249}	4γ	$< 9 \times 10^{-6}$	DESIG=81
Γ_{250}	5γ	$< 1.5 \times 10^{-5}$	DESIG=244
Γ_{251}	$\gamma\pi^0$	$(3.39 \pm 0.08) \times 10^{-5}$	DESIG=245
Γ_{252}	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	DESIG=82
Γ_{253}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	DESIG=283
Γ_{254}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	DESIG=95
Γ_{255}	$\gamma f_2(1270) f_2(1270) (\text{non resonant})$	$(8.2 \pm 1.9) \times 10^{-4}$	DESIG=203
Γ_{256}	$\gamma\pi^+\pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	DESIG=204
Γ_{257}	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	DESIG=99
Γ_{258}	$\gamma(K\bar{K}\pi) [J^{PC} = 0-+]$	$(7 \pm 4) \times 10^{-4}$	S=2.1 DESIG=378
Γ_{259}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	DESIG=176
Γ_{260}	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	DESIG=143
Γ_{261}	$\gamma\eta$	$(1.090 \pm 0.013) \times 10^{-3}$	DESIG=145
Γ_{262}	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	DESIG=83
Γ_{263}	$\gamma f_0(500) \rightarrow \gamma\pi\pi$		DESIG=292
Γ_{264}	$\gamma f_0(500) \rightarrow \gamma K\bar{K}$		DESIG=398
Γ_{265}	$\gamma f_0(500) \rightarrow \gamma\eta\eta$		DESIG=399
Γ_{266}	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	$< 2.5 \times 10^{-6}$	DESIG=400 CL=95%
Γ_{267}	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	$< 6.6 \times 10^{-6}$	DESIG=293
Γ_{268}	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	DESIG=294
Γ_{269}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	DESIG=96
Γ_{270}	$\gamma\eta'(958)$	$(5.28 \pm 0.06) \times 10^{-3}$	DESIG=142
Γ_{271}	$\gamma f_0(980) \rightarrow \gamma\pi\pi$		DESIG=84
Γ_{272}	$\gamma f_0(980) \rightarrow \gamma K\bar{K}$		DESIG=393
Γ_{273}	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$	DESIG=394
Γ_{274}	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	DESIG=226 CL=90%

Γ_{275}	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=227
Γ_{276}	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$		DESIG=97
Γ_{277}	$\gamma\phi\phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1	DESIG=98
Γ_{278}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6	DESIG=89
Γ_{279}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8	DESIG=171
Γ_{280}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=170
Γ_{281}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3	DESIG=124
Γ_{282}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%	DESIG=212
Γ_{283}	$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0$	$(1.50 \pm 0.16) \times 10^{-5}$		DESIG=476
Γ_{284}	$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^0\pi^0\pi^0$	$(7.1 \pm 1.1) \times 10^{-6}$		DESIG=477
Γ_{285}	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	$< 2.63 \times 10^{-6}$	CL=90%	DESIG=348
Γ_{286}	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	$< 1.86 \times 10^{-6}$	CL=90%	DESIG=349
Γ_{287}	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		DESIG=125
Γ_{288}	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	$(1.98 \pm 0.33) \times 10^{-3}$		DESIG=224
Γ_{289}	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	$< 4.80 \times 10^{-6}$	CL=90%	DESIG=347
Γ_{290}	$\gamma\eta(2225)$	$(3.14 \pm 0.50) \times 10^{-4}$		DESIG=126
Γ_{291}	$\gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=86
Γ_{292}	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm 0.60) \times 10^{-5}$		DESIG=373
Γ_{293}	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		DESIG=88
Γ_{294}	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$			DESIG=395
Γ_{295}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		DESIG=284
Γ_{296}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$		DESIG=368
Γ_{297}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$			DESIG=396
Γ_{298}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$			DESIG=397
Γ_{299}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$		DESIG=175
Γ_{300}	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$(1.09 \pm 0.24) \times 10^{-4}$		DESIG=172
Γ_{301}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$(1.7 \pm 0.6) \times 10^{-5}$		DESIG=265
Γ_{302}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm 0.24) \times 10^{-5}$		DESIG=369
Γ_{303}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$			DESIG=401
Γ_{304}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		DESIG=141
Γ_{305}	$\gamma f'_2(1525)$	$(5.7 \pm 0.8) \times 10^{-4}$	S=1.5	DESIG=87
Γ_{306}	$\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm 0.7) \times 10^{-5}$		DESIG=374
Γ_{307}	$\gamma f'_2(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$		DESIG=268
Γ_{308}	$\gamma f_2(1565) \rightarrow \gamma\eta\eta'$			DESIG=432
Γ_{309}	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$		DESIG=222
Γ_{310}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=135
Γ_{311}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm 1.0) \times 10^{-4}$	S=1.5	DESIG=91
Γ_{312}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$		DESIG=221
Γ_{313}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm 1.2) \times 10^{-4}$		DESIG=266
Γ_{314}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta'$			DESIG=402
Γ_{315}	$\gamma f_0(1710) \rightarrow \gamma\omega\phi$	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=262
Γ_{316}	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm 0.20) \times 10^{-5}$		DESIG=370
Γ_{317}	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm 3.5) \times 10^{-5}$		DESIG=269
Γ_{318}	$\gamma\eta_1(1855) \rightarrow \gamma\eta\eta'$	$(2.7 \pm 0.4) \times 10^{-6}$		DESIG=447
Γ_{319}	$\gamma f_0(1770) \rightarrow \gamma\eta\eta'$			DESIG=431
Γ_{320}	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	$(2.0 \pm 1.4) \times 10^{-4}$		DESIG=223
Γ_{321}	$\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$		DESIG=144
Γ_{322}	$\gamma f_2(2010) \rightarrow \gamma\eta\eta'$			DESIG=440

Γ_{323}	$\gamma f_0(2020) \rightarrow \gamma\pi\pi$	DESIG=403
Γ_{324}	$\gamma f_0(2020) \rightarrow \gamma K\bar{K}$	DESIG=404
Γ_{325}	$\gamma f_0(2020) \rightarrow \gamma\eta\eta$	DESIG=405
Γ_{326}	$\gamma f_0(2020) \rightarrow \gamma\eta'\eta'$	$(2.63 \pm 0.32) \times 10^{-4}$ DESIG=426
Γ_{327}	$\gamma f_0(2020) \rightarrow \gamma\eta\eta'$	DESIG=438
Γ_{328}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$ DESIG=100
Γ_{329}	$\gamma f_4(2050) \rightarrow \gamma\eta\eta'$	DESIG=433
Γ_{330}	$\gamma f_0(2100) \rightarrow \gamma\eta\eta$	$(1.13 \pm 0.60) \times 10^{-4}$ DESIG=267
Γ_{331}	$\gamma f_0(2100) \rightarrow \gamma K\bar{K}$	DESIG=406
Γ_{332}	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	$(6.2 \pm 1.0) \times 10^{-4}$ DESIG=286
Γ_{333}	$\gamma f_0(2200)$	seen DESIG=123; OUR EST; → UNCHECKED ←
Γ_{334}	$\gamma f_0(2200) \rightarrow \gamma K\bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$ DESIG=285
Γ_{335}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm 0.19) \times 10^{-4}$ DESIG=371
Γ_{336}	$\gamma f_0(2200) \rightarrow \gamma\pi\pi$	DESIG=407
Γ_{337}	$\gamma f_0(2200) \rightarrow \gamma\eta\eta$	DESIG=408
Γ_{338}	$\gamma f_J(2220)$	seen DESIG=92; OUR EST; → UNCHECKED ←
Γ_{339}	$\gamma f_J(2220) \rightarrow \gamma\pi\pi$	$< 3.9 \times 10^{-5}$ CL=90% DESIG=136
Γ_{340}	$\gamma f_J(2220) \rightarrow \gamma K\bar{K}$	$< 4.1 \times 10^{-5}$ CL=90% DESIG=137
Γ_{341}	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$ DESIG=138
Γ_{342}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$ DESIG=372
Γ_{343}	$\gamma f_0(2330) \rightarrow \gamma\pi\pi$	DESIG=409
Γ_{344}	$\gamma f_0(2330) \rightarrow \gamma\eta\eta$	DESIG=410
Γ_{345}	$\gamma f_0(2330) \rightarrow \gamma\eta'\eta'$	$(6.1 \pm 4.0) \times 10^{-6}$ DESIG=427
Γ_{346}	$\gamma f_0(2330) \rightarrow \gamma\eta\eta'$	DESIG=439
Γ_{347}	$\gamma f_2(2340) \rightarrow \gamma\eta\eta$	$(5.6 \pm 2.4) \times 10^{-5}$ DESIG=270
Γ_{348}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm 4.0) \times 10^{-5}$ DESIG=375
Γ_{349}	$\gamma f_2(2340) \rightarrow \gamma\eta'\eta'$	$(8.7 \pm 0.9) \times 10^{-6}$ DESIG=428
Γ_{350}	$\gamma f_0(2470) \rightarrow \gamma\eta'\eta'$	$(8.2 \pm 4.0) \times 10^{-7}$ DESIG=429
Γ_{351}	$\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta'$	$(2.7 \pm 0.6) \times 10^{-4}$ S=1.6 DESIG=213
Γ_{352}	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	$(7.7 \pm 1.5) \times 10^{-5}$ DESIG=254
Γ_{353}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm 2.0) \times 10^{-5}$ DESIG=282
Γ_{354}	$\gamma X(1835) \rightarrow \gamma\gamma\phi(1020)$	DESIG=346
Γ_{355}	$\gamma X(1835) \rightarrow \gamma\gamma\gamma$	$< 3.56 \times 10^{-6}$ CL=90% DESIG=350
Γ_{356}	$\gamma X(1835) \rightarrow \gamma 3(\pi^+\pi^-)$	$(2.4 \pm 0.7) \times 10^{-5}$ DESIG=264
Γ_{357}	$\gamma\eta(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$ DESIG=388
Γ_{358}	$\gamma\eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$ DESIG=389
Γ_{359}	$\gamma\eta(2370) \rightarrow \gamma\eta\eta\eta'$	$< 9.2 \times 10^{-6}$ CL=90% DESIG=390
Γ_{360}	$\gamma D^0 + c.c.$	$< 9.1 \times 10^{-8}$ CL=90% DESIG=479
Γ_{361}	$\gamma p\bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$ DESIG=90
Γ_{362}	$\gamma p\bar{p}\pi^+\pi^-$	$< 7.9 \times 10^{-4}$ CL=90% DESIG=93
Γ_{363}	$\gamma\Lambda\bar{\Lambda}$	$< 1.3 \times 10^{-4}$ CL=90% DESIG=200
Γ_{364}	$\gamma A^0 \rightarrow \gamma \text{invisible}$	[e] $< 1.7 \times 10^{-6}$ CL=90% DESIG=251
Γ_{365}	$\gamma A^0 \rightarrow \gamma\gamma\gamma$	$< 4.9 \times 10^{-7}$ CL=95% DESIG=469
Γ_{366}	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	[f] $< 7.8 \times 10^{-7}$ CL=90% DESIG=259

Dalitz decays

Γ_{367}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$	NODE=M070;CLUMP=G
Γ_{368}	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$	DESIG=271
Γ_{369}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$	DESIG=272
Γ_{370}	$\eta(1405) e^+ e^- \rightarrow f_0(980) \pi^0 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 e^+ e^-$	$(2.04 \pm 0.22) \times 10^{-7}$	DESIG=273
Γ_{371}	$X(1835) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(3.58 \pm 0.25) \times 10^{-6}$	DESIG=475
Γ_{372}	$X(2120) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(8.2 \pm 1.3) \times 10^{-7}$	DESIG=423
Γ_{373}	$\eta(2370) e^+ e^-$, $\eta \rightarrow \pi^+ \pi^- \eta'$	$(1.08 \pm 0.17) \times 10^{-6}$	DESIG=425
Γ_{374}	$\eta U \rightarrow \eta e^+ e^-$	$[g] < 9.11 \times 10^{-7}$ CL=90%	DESIG=424
Γ_{375}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	$[g] < 2.0 \times 10^{-7}$ CL=90%	DESIG=352
Γ_{376}	$\phi e^+ e^-$	$< 1.2 \times 10^{-7}$ CL=90%	DESIG=366
			DESIG=384

Weak decays

Γ_{377}	$D^- e^+ \nu_e + \text{c.c.}$	$< 7.1 \times 10^{-8}$ CL=90%	NODE=M070;CLUMP=E
Γ_{378}	$D^- \mu^+ \nu_\mu + \text{c.c.}$	$< 5.6 \times 10^{-7}$ CL=90%	DESIG=218
Γ_{379}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 8.5 \times 10^{-8}$ CL=90%	DESIG=468
Γ_{380}	$\bar{D}_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$ CL=90%	DESIG=219
Γ_{381}	$\bar{D}_s^* - e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$ CL=90%	DESIG=220
Γ_{382}	$D^- \pi^+ + \text{c.c.}$	$< 7.0 \times 10^{-8}$ CL=90%	DESIG=290
Γ_{383}	$D^- \rho^+ + \text{c.c.}$	$< 6.0 \times 10^{-7}$ CL=90%	DESIG=241
Γ_{384}	$\bar{D}^0 \pi^0 + \text{c.c.}$	$< 4.7 \times 10^{-7}$ CL=90%	DESIG=464
Γ_{385}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$ CL=90%	DESIG=275
Γ_{386}	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	$< 2.5 \times 10^{-6}$ CL=90%	DESIG=465
Γ_{387}	$\bar{D}^0 \eta + \text{c.c.}$	$< 6.8 \times 10^{-7}$ CL=90%	DESIG=242
Γ_{388}	$\bar{D}^0 \rho^0 + \text{c.c.}$	$< 5.2 \times 10^{-7}$ CL=90%	DESIG=466
Γ_{389}	$\bar{D}_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$ CL=90%	DESIG=467
Γ_{390}	$\bar{D}_s^- \rho^+ + \text{c.c.}$	$< 1.3 \times 10^{-5}$ CL=90%	DESIG=243
			DESIG=274

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{391}	$\gamma\gamma$	$C < 2.7 \times 10^{-7}$ CL=90%	NODE=M070;CLUMP=D
Γ_{392}	$\gamma\phi$	$C < 1.4 \times 10^{-6}$ CL=90%	DESIG=80
Γ_{393}	$e^\pm \mu^\mp$	$LF < 1.6 \times 10^{-7}$ CL=90%	DESIG=277
Γ_{394}	$e^\pm \tau^\mp$	$LF < 7.5 \times 10^{-8}$ CL=90%	DESIG=177
Γ_{395}	$\mu^\pm \tau^\mp$	$LF < 2.0 \times 10^{-6}$ CL=90%	DESIG=178
Γ_{396}	$\Lambda_c^+ e^- + \text{c.c.}$	$< 6.9 \times 10^{-8}$ CL=90%	DESIG=179
			DESIG=379

Other decays

Γ_{397}	invisible	$< 7 \times 10^{-4}$ CL=90%	DESIG=240
Γ_{398}	$\mu^+ \mu^- X^0 \rightarrow \mu^+ \mu^- + \text{invisible}$		DESIG=470

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] $\Theta(1540)$ is a hypothetical pentaquark state of 1.54 GeV/c 2 mass and a width of less than 25 MeV/c 2 .

[d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.

[e] For a narrow state A with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

[g] For a dark photon U with mass between 100 and 2100 MeV.

LINKAGE=EGM

LINKAGE=SG

LINKAGE=THT

LINKAGE=MF

LINKAGE=NSA

LINKAGE=NA0

LINKAGE=DPH

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

J/ ψ (1S) PARTIAL WIDTHS **Γ (hadrons)**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_1
81.37 ± 1.36 ± 1.30	¹ ANASHIN	20	KEDR $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
74.1 ± 8.1	BAI	95B	BES $e^+ e^-$	
59 ± 24	BALDINI-...	75	FRAG $e^+ e^-$	
59 ± 14	BOYARSKI	75	MRK1 $e^+ e^-$	
50 ± 25	ESPOSITO	75B	FRAM $e^+ e^-$	

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there.

 $\Gamma(e^+ e^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5
5.53 ± 0.10 OUR AVERAGE					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.550 ± 0.056 ± 0.089	1,2 ANASHIN	18A	KEDR $e^+ e^-$		
5.36 ^{+0.29} _{-0.28}	³ HSUEH	92	RVUE See γ mini-review		

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.58 ± 0.05 ± 0.08	⁴ ABLIKIM	16Q	BES3 $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$		
5.71 ± 0.16	^{13k} ADAMS	06A	CLEO $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$		
5.57 ± 0.19	^{7.8k} AUBERT	04	BABR $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$		
5.14 ± 0.39	BAI	95B	BES $e^+ e^-$		
4.72 ± 0.35	ALEXANDER	89	RVUE See γ mini-review		
4.4 ± 0.6	³ BRANDELIK	79C	DASP $e^+ e^-$		
4.6 ± 0.8	⁶ BALDINI-...	75	FRAG $e^+ e^-$		
4.8 ± 0.6	BOYARSKI	75	MRK1 $e^+ e^-$		
4.6 ± 1.0	ESPOSITO	75B	FRAM $e^+ e^-$		

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the J/ ψ (1S) peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$.

 $\Gamma(\mu^+ \mu^-)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_7
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.13 ± 0.52	BAI	95B	BES $e^+ e^-$	
4.8 ± 0.6	BOYARSKI	75	MRK1 $e^+ e^-$	
5 ± 1	ESPOSITO	75B	FRAM $e^+ e^-$	

 $\Gamma(\gamma\gamma)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{391}
<5.4	90	BRANDELIK	79C	DASP $e^+ e^-$	

J/ ψ (1S) $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel(I) in the $e^+ e^-$ annihilation.

 $\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_5 / \Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.884 ± 0.048 ± 0.078	^{1,2} ANASHIN	18A	KEDR $e^+ e^-$	
4 ± 0.8	³ BALDINI-...	75	FRAG $e^+ e^-$	
3.9 ± 0.8	³ ESPOSITO	75B	FRAM $e^+ e^-$	

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the J/ ψ (1S) peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

NODE=M070220

NODE=M070W3

NODE=M070W3

NODE=M070W3;LINKAGE=A

NODE=M070W1

NODE=M070W1

OCCUR=4

NODE=M070W1;LINKAGE=D

NODE=M070W1;LINKAGE=E

NODE=M070W1;LINKAGE=F

NODE=M070W1;LINKAGE=A

NODE=M070W1;LINKAGE=AA

NODE=M070W1;LINKAGE=B

NODE=M070W2

NODE=M070W2

NODE=M070W70

NODE=M070W70

NODE=M070225

NODE=M070225

NODE=M070G3

NODE=M070G3

NODE=M070G3;LINKAGE=A

NODE=M070G3;LINKAGE=B

NODE=M070G3;LINKAGE=S

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_5\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
333.1 ± 6.6 ± 4.0	1,2	ANASHIN	18A	KEDR e^+e^-	
332.3 ± 6.4 ± 4.8		ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow e^+e^-$	
350 ± 20		BRANDELIK	79C	DASP e^+e^-	
320 ± 70	3	BALDINI...	75	FRAG e^+e^-	
340 ± 90	3	ESPOSITO	75B	FRAM e^+e^-	
360 ± 100	3	FORD	75	SPEC e^+e^-	

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
333 ± 4 OUR AVERAGE					
333.4 ± 2.5 ± 4.4		ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$	
331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow \mu^+\mu^-$	
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$	
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
510 ± 90		DASP	75	DASP e^+e^-	
380 ± 50		1 ESPOSITO	75B	FRAM e^+e^-	

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{16}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.3 ± 0.4 OUR AVERAGE					
2.34 ± 0.43 ± 0.16	49	LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$	
2.22 ± 0.96 ± 0.02	9	1 AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$	
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{18}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
64.8 ± 11.1 ± 0.4					
200	1	LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$	
¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{19}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
26.9 ± 5.7 ± 0.1					
101	1	LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-3\pi^0\gamma\gamma)$	
¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\eta K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{22}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.76 ± 1.64 ± 0.03					
1	LEES	23	BABR	$e^+e^- \rightarrow \gamma_{ISR}$ hadrons	
¹ LEES 23 reports $[\Gamma(J/\psi(1S) \rightarrow \eta K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 1.55 \pm 0.51 \pm 0.16$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{23}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.3 ± 1.4 ± 0.4					
44	LEES	17D	BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$	

NODE=M070G1
NODE=M070G1

OCCUR=2

NODE=M070G1;LINKAGE=A

NODE=M070G1;LINKAGE=B

NODE=M070G1;LINKAGE=S

NODE=M070G2
NODE=M070G2

NODE=M070G2;LINKAGE=S

NODE=M070G25
NODE=M070G25

NODE=M070G25;LINKAGE=AU

NODE=M070Q20
NODE=M070Q20

NODE=M070Q20;LINKAGE=A

NODE=M070Q19
NODE=M070Q19

NODE=M070Q19;LINKAGE=A

NODE=M070Q54
NODE=M070Q54

NODE=M070Q54;LINKAGE=A

NODE=M070G38
NODE=M070G38

$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{26}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
155±26±36	14k	LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$	

NODE=M070Q13
NODE=M070Q13

$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{27}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
32±13±15	14k	LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$	

NODE=M070Q14
NODE=M070Q14

$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{29}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.4±1.0±1.9	130	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$	

NODE=M070G31
NODE=M070G31

$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{44}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
53.6±5.0±0.4	788	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$	

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24
NODE=M070G24

$\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{45}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
27.8±3.5±0.2	398	¹ LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$	

NODE=M070P54
NODE=M070P54

¹LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24;LINKAGE=AU

$\Gamma(\omega 3\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{46}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.5±3.1±0.1	89	¹ LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR} (\pi^+ \pi^- 4\pi^0)$	

NODE=M070Q18
NODE=M070Q18

¹LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P54;LINKAGE=A

$\Gamma(\omega \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{48}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
16.9±7.6±0.2	1	LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR} (\pi^+ \pi^- 4\pi^0)$	

NODE=M070Q21
NODE=M070Q21

¹Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best values $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070Q21;LINKAGE=A

$\Gamma(\omega \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{49}\Gamma_5/\Gamma$
VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.2±0.3±0.2	170	AUBERT	06D	BABR $10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \pi^0 \gamma$	

NODE=M070G8
NODE=M070G8

$\Gamma(\omega \pi^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{50}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.90±0.96±0.01	27	¹ LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$	

NODE=M070P58
NODE=M070P58

¹LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P58;LINKAGE=A

$\Gamma(\omega \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{52}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
185±30±1	14k	¹ LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$	

NODE=M070Q11
NODE=M070Q11

¹LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^- 2\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 165 \pm 9 \pm 25$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q11;LINKAGE=A

$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{64}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.70±1.98±0.03	24	1 AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega K^+K^-\gamma$	NODE=M070G29 NODE=M070G29
1 AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 3.3 \pm 1.3 \pm 1.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{73}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
29.0±1.7±1.3		AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^*(892)^-\gamma$	NODE=M070G18 NODE=M070G18
$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{74}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.96±0.85±0.70	155	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\gamma$	NODE=M070G20 NODE=M070G20
$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{75}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
16.76±1.70±1.00	89	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\gamma$	NODE=M070G21 NODE=M070G21
$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{76}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
26.6±2.5±1.5		AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^0\bar{K}^*(892)^0\gamma$	NODE=M070G19 NODE=M070G19
$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{77}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
17.70±1.70±1.00	94	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\gamma$	NODE=M070G22 NODE=M070G22
$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{78}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
42.6±4.8±7.2	99	1 LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\pi^0\gamma$	NODE=M070G39 NODE=M070G39
1 Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0\pi^0) = 1/6$.					
$\Gamma(K^*(892)^\pm K^\mp\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{79}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
22.8±2.8±6.8	80	1 LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\pi^0\gamma$	NODE=M070G32 NODE=M070G32
1 Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0\pi^\pm) = 1/4$.					
$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{80}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.0±2.8 OUR AVERAGE					NODE=M070GY4 NODE=M070GY4
9.2±1.2±3.2	64	1 LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\pi^0\gamma$	
14.8±4.8±1.2	53	2 LEES	14H BABR	$e^+e^- \rightarrow \pi^+\pi^- K_S^0 K_S^0\gamma$	
1 Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm\pi^\mp) = 1/2$.					
2 Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0\pi) = 1/4$.					
$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{81}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.7±1.2±0.3	53	LEES	14H BABR	$e^+e^- \rightarrow \pi^+\pi^- K_S^0 K_S^0\gamma$	NODE=M070GY5 NODE=M070GY5
$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{84}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.60±0.75±2.25	34	1 LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp\pi^0\gamma$	NODE=M070G33 NODE=M070G33
1 Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+\pi^-) = 2/3$.					
$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{86}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.28±0.34±0.07	47±12	1 LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^- K^+K^-\gamma$	NODE=M070G01 NODE=M070G01
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.28±0.40±0.11	25 ± 8	1,2 AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^- K^+K^-\gamma$	
1 Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3$ $B(K^{*0} \rightarrow K\pi)$.					
2 Superseded by LEES 12F.					

$\Gamma(K^*(892)^{\pm} K^*(892)^{\mp}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{87}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.80±0.48±0.32	1 ± 5	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{98}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
20.1±9.8±0.5	35	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by $1/4$ to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 10.0 \pm 4.8 \pm 0.8$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{99}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
25.8±1.4±0.6	710	1,2,3 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1 317 2,4 AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by $2/3$ to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{100}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
18.6±16.1±0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{101}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.32±2.00±0.08	8 ± 8	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

¹ Dividing by $1/4$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{103}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.8±0.4±0.3	110 ± 14	1 AUBERT 07AK BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	

¹ Dividing by $2/3$ to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

NODE=M070GY8
NODE=M070GY8

NODE=M070GY8;LINKAGE=A

NODE=M070GY6
NODE=M070GY6

NODE=M070GY6;LINKAGE=A

NODE=M070GY6;LINKAGE=B

NODE=M070G02
NODE=M070G02

NODE=M070G02;LINKAGE=A

NODE=M070G02;LINKAGE=AE
NODE=M070G02;LINKAGE=B
NODE=M070G02;LINKAGE=UB

NODE=M070GY9
NODE=M070GY9

NODE=M070GY9;LINKAGE=A

NODE=M070GY9;LINKAGE=B

NODE=M070GZ0
NODE=M070GZ0

NODE=M070GZ0;LINKAGE=A

NODE=M070G03
NODE=M070G03

NODE=M070G03;LINKAGE=AE

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{110}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.41±0.34 OUR AVERAGE				
4.39±0.48±0.05	181	1 LEES	12F BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
4.44±0.48±0.05	254 ± 23	2 SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.7 ± 0.1	103	3 AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14

NODE=M070G14

NODE=M070G14;LINKAGE=B

NODE=M070G14;LINKAGE=SH

NODE=M070G14;LINKAGE=AU

 $\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{111}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.73±0.56±0.03	45	1 LEES	12F BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.09±0.86±0.03	23	2 AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G15

NODE=M070G15

NODE=M070G15;LINKAGE=A

NODE=M070G15;LINKAGE=AU

 $\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{112}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.94±0.19±0.01	35	1 AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \phi 2(\pi^+\pi^-)\gamma$
¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G10

NODE=M070G10

NODE=M070G10;LINKAGE=AU

 $\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{113}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.6±1.4 OUR AVERAGE				

4.1±1.6±0.4	1 LEES	23 BABR	$e^+e^- \rightarrow \gamma_{ISR}$ hadrons
6.1±2.7±0.4	6 2 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$

¹ LEES 23 quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi^0) = 0.64 \pm 0.26 \pm 0.06$ eV.

² AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

NODE=M070G28

NODE=M070G28

OCCUR=2

NODE=M070G28;LINKAGE=A

NODE=M070G28;LINKAGE=AU

 $\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{117}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.42±0.18 OUR AVERAGE				

1.38±0.24±0.02	57 ± 9	1 LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
1.48±0.27±0.09	60±11	2 SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.00±0.23±0.01	20 ± 5	3 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	----------	-----------	--

NODE=M070G05

NODE=M070G05

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.

³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{118}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.96±0.26±0.01 16 ± 4 ¹ LEES 12F BABR $10.6 \text{ e}^+\text{e}^- \rightarrow \pi^0\pi^0K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.94 \pm 0.39 \pm 0.01$ 7.0 ± 2.8 ² AUBERT 07AK BABR $10.6 \text{ e}^+\text{e}^- \rightarrow \pi^0\pi^0K^+K^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.48 \pm 0.12 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.47 \pm 0.19 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{126}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.79±0.32±0.02 61 1,2,3 LEES 12F BABR $10.6 \text{ e}^+\text{e}^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.08 \pm 0.73 \pm 0.05$ 44 2,4 AUBERT 07AK BABR $10.6 \text{ e}^+\text{e}^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 1.51 \pm 0.25 \pm 0.10 \text{ eV}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3 \pm 2.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+\pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28 \text{ eV}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3 \pm 2.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{131}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

8.1±3.2±0.2 11 1,2 LEES 14H BABR $e^+\text{e}^- \rightarrow K_S^0K_S^0K^+K^-\gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ and using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3 \text{ eV}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{137}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.53±0.61±0.05 163 ¹ LEES 12F BABR $10.6 \text{ e}^+\text{e}^- \rightarrow K^+K^-K^+K^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.26 \pm 0.26 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G05;LINKAGE=A

NODE=M070G05;LINKAGE=SH

NODE=M070G05;LINKAGE=UB

NODE=M070G06
NODE=M070G06

NODE=M070G06;LINKAGE=A

NODE=M070G06;LINKAGE=UB

NODE=M070G07
NODE=M070G07

NODE=M070G07;LINKAGE=A

NODE=M070G07;LINKAGE=AE
NODE=M070G07;LINKAGE=B

NODE=M070G07;LINKAGE=UB

NODE=M070GZ2
NODE=M070GZ2

NODE=M070GZ2;LINKAGE=A

NODE=M070GZ2;LINKAGE=B

NODE=M070G09
NODE=M070G09

NODE=M070G09;LINKAGE=A

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{138}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.21±0.83±0.03	29	1 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$	
1 LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(f'_2(1525) K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{143}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$	
1 Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$. 2 LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525) K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{164}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$	
$\Gamma(\pi^+ \pi^- 3\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{166}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
100 ± 50 OUR AVERAGE	Error includes scale factor of 4.3.				
55 ± 16 ± 1	14k	1 LEES	21 BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$	
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$	
1 LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)/\Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2$ eV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)/\Gamma_{\text{total}} = 0.3469 \pm 0.0034$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(\pi^+ \pi^- 4\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{169}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
35.8±4.4±5.4	340	LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR} (\pi^+ \pi^- 4\pi^0)$	
$\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{167}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
78.0±9.0±8.0	1.2k	LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$	
$\Gamma(\rho^+ \rho^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{168}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
33.0±5.0±3.3	529	LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$	
$\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{170}\Gamma_5/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.1248±0.0019±0.0026	LEES 21B BABR $10.5 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$				
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.122 ± 0.005 ± 0.008	AUBERT,B	04N BABR	10.6 BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$	
$\Gamma(2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{171}\Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.9±0.5±1.0	761	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^- \pi^0) \gamma$	
$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{172}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
107.0±4.3±6.4	768	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$	
$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{174}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
20.4±0.9±0.4	LEES 12E BABR $10.6 e^+ e^- \rightarrow 2\pi^+ 2\pi^- \gamma$				
• • • We do not use the following data for averages, fits, limits, etc. • • •					
19.5 ± 1.4 ± 1.3	270	1 AUBERT	05D BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \gamma$	

¹ Superseded by LEES 12E.

NODE=M070GZ1
NODE=M070GZ1

NODE=M070GZ1;LINKAGE=A

NODE=M070GZ4
NODE=M070GZ4

NODE=M070GZ4;LINKAGE=A

NODE=M070GZ4;LINKAGE=B

NODE=M070G23
NODE=M070G23

NODE=M070P53
NODE=M070P53

NODE=M070P53;LINKAGE=A

NODE=M070Q17
NODE=M070Q17

NODE=M070P55
NODE=M070P55

NODE=M070P56
NODE=M070P56

NODE=M070G5
NODE=M070G5

NODE=M070G7
NODE=M070G7

NODE=M070G27
NODE=M070G27

NODE=M070G11
NODE=M070G11

NODE=M070G11;LINKAGE=AU

$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{175}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D	BABR $10.6 e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$

 $\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{176}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
345±10±50	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

 $\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{178}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{180}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

 $\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{181}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3 14k ¹LEES 21 BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
 12.8± 1.8±2.0 203 LEES 18E BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 6 \pm 4 \pm 1 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{182}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

 $\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{183}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

1.78±0.11±0.05 462 ¹LEES 15J BABR $e^+e^- \rightarrow K^+K^-\gamma$
 1.94±0.11±0.05 462 ²LEES 15J BABR $e^+e^- \rightarrow K^+K^-\gamma$
 1.42±0.23±0.08 51 ³LEES 13Q BABR $e^+e^- \rightarrow K^+K^-\gamma$

1 $\sin\phi > 0$.
 2 $\sin\phi < 0$.

3 Interference with non-resonant K^+K^- production not taken into account.

 $\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{189}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES	17A	BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{190}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.9±0.4	106	LEES	17A	BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$

 $\Gamma(K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{191}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.7±0.1	37	LEES	17A	BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_{192}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94±0.81±1.10	3.1k	LEES	12F	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ± 1.3 ± 2.1 1.5k ¹AUBERT 07AK BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
 33.6 ± 2.7 ± 2.7 233 ²AUBERT 05D BABR $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

1 Superseded by LEES 12F.

2 Superseded by AUBERT 07AK.

NODE=M070G6
 NODE=M070G6

NODE=M070Q12
 NODE=M070Q12

NODE=M070G26
 NODE=M070G26

NODE=M070Q15
 NODE=M070Q15

NODE=M070P57
 NODE=M070P57

NODE=M070G08
 NODE=M070G08

OCCUR=2

NODE=M070G08;LINKAGE=A
 NODE=M070G08;LINKAGE=B
 NODE=M070G08;LINKAGE=BA

NODE=M070G41
 NODE=M070G41

NODE=M070G42
 NODE=M070G42

NODE=M070G43
 NODE=M070G43

NODE=M070G12
 NODE=M070G12

NODE=M070G12;LINKAGE=B
 NODE=M070G12;LINKAGE=AU

$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{193} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.75 ± 0.81 ± 0.90	388	LEES	12F	BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ± 1.1 ± 1.3	203	¹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
------------------	-----	---------------------	-----------	---

¹ Superseded by LEES 12F.

$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{194} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.9 ± 1.3 ± 0.9		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons	

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{195} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
29.3 ± 2.6 ± 2.9		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons	

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{196} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
34.6 ± 1.4 ± 1.8		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons	

$\Gamma(K_S^0 K^\pm \rho(770)^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{197} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
16.0 ± 4.1 ± 1.6		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons	

$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{198} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
20.8 ± 2.3 ± 2.1	248	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$	

$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{199} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.3 ± 2.3 ± 0.5	47	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$	

$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{200} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.0 ± 1.8 ± 0.4	45	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$	

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{201} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.3 ± 0.9 ± 0.5	133	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$	

$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{202} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
31.7 ± 1.9 ± 1.8	393	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$	

$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{203} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
17.3 ± 2.1 ± 1.7		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons	

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{204} \Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.75 ± 0.23 ± 0.17	205	AUBERT	06D	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$	

$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{205} \Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
25.9 ± 3.9 ± 0.1	73	¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$	

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G04
NODE=M070G04

NODE=M070G04;LINKAGE=A

NODE=M070Q51
NODE=M070Q51

NODE=M070Q52
NODE=M070Q52

NODE=M070Q62
NODE=M070Q62

NODE=M070GY1
NODE=M070GY1

NODE=M070G40
NODE=M070G40


NODE=M070G35
NODE=M070G35

NODE=M070G34
NODE=M070G34

NODE=M070Q55
NODE=M070Q55

NODE=M070G9
NODE=M070G9

NODE=M070G30
NODE=M070G30



$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7±0.9±0.7	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma_{230}\Gamma_5/\Gamma$

NODE=M070G16
NODE=M070G16

$\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\Gamma_{241}\Gamma_5/\Gamma$

NODE=M070Q50
NODE=M070Q50

$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.2±1.5±0.6	GONG	23 BELL	$e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

$\Gamma_{242}\Gamma_5/\Gamma$

NODE=M070G17
NODE=M070G17

6.4±1.2±0.6

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.4±1.2±0.6	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

OCCUR=2

J/ψ(1S) BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.877±0.005 OUR AVERAGE			

Γ_1/Γ

NODE=M070R3
NODE=M070R3

0.878±0.005

VALUE	DOCUMENT ID	TECN	COMMENT
0.878±0.005	BAI	95B BES	$e^+ e^-$

0.86 ± 0.02

VALUE	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.02	BOYARSKI	75 MRK1	$e^+ e^-$

VALUE	DOCUMENT ID	TECN	COMMENT
0.1346±0.0007	1 LIAO	23 RVUE	$e^+ e^-$

Γ_1/Γ

NODE=M070R3
NODE=M070R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.135 ± 0.003

VALUE	DOCUMENT ID	TECN	COMMENT
0.135 ± 0.003	2,3 SETH	04 RVUE	$e^+ e^-$

0.17 ± 0.02

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.02	2 BOYARSKI	75 MRK1	$e^+ e^-$

Γ_2/Γ

NODE=M070R4
NODE=M070R4

¹ Using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (5.967 \pm 0.023)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

NODE=M070R4;LINKAGE=A

² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

NODE=M070R4;LINKAGE=C

³ Using $B(J/\psi \rightarrow \ell^+ \ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

NODE=M070R4;LINKAGE=SE

$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
64.1±1.0	6 M	¹ BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+ \ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\eta_C)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.79±1.05	200 k	¹ BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_4/Γ_3
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
13.7±0.1±0.7	6 M	BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.971±0.032 OUR AVERAGE					
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B	BES	$e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.8±1.3±0.4		¹ ARMSTRONG	96	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$

¹ For $E_\gamma > 100$ MeV.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.961±0.033 OUR AVERAGE					
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B	BES	$e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$					Γ_5/Γ_7
VALUE	DOCUMENT ID	TECN	COMMENT		
1.0016±0.0031 OUR AVERAGE					
1.0022±0.0044±0.0048	¹ AULCHENKO	14	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$	
1.0017±0.0017±0.0033	² ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.002 ± 0.021 ± 0.013	³ ANASHIN	10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$	
0.997 ± 0.012 ± 0.006	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.011 ± 0.013 ± 0.016	BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.00 ± 0.07	BAI	95B	BES	$e^+ e^-$	
1.00 ± 0.05	BOYARSKI	75	MRK1	$e^+ e^-$	
0.91 ± 0.15	ESPOSITO	75B	FRAM	$e^+ e^-$	
0.93 ± 0.10	FORD	75	SPEC	$e^+ e^-$	

¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$.

NODE=M070S65
NODE=M070S65

NODE=M070S65;LINKAGE=BE

NODE=M070S66
NODE=M070S66

NODE=M070S66;LINKAGE=BE

NODE=M070S67
NODE=M070S67

NODE=M070R1
NODE=M070R1

NODE=M070S33
NODE=M070S33

NODE=M070S33;LINKAGE=A

NODE=M070R2
NODE=M070R2

NODE=M070R5
NODE=M070R5

NODE=M070R5;LINKAGE=A
NODE=M070R5;LINKAGE=AB

NODE=M070R5;LINKAGE=AN

$\Gamma(e^+e^-e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.48±0.31±0.45	700	1 ABLIKIM	24L BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$1 B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

 Γ_8/Γ

NODE=M070Q71
NODE=M070Q71

 $\Gamma(e^+e^-\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53±0.22±0.13	354	1 ABLIKIM	24L BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$1 B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

 Γ_9/Γ

NODE=M070Q72
NODE=M070Q72

 $\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.3±1.1±0.1		452	1 AAIJ	24AE LHCb	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$
<16		90	2 ABLIKIM	24L BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
10.1 ^{+3.3} _{-2.8} ±0.1		12	3 HAYRAPETYAN	24A CMS	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$

¹ AAIJ 24AE reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (1.89 \pm 0.17 \pm 0.09) \times 10^{-5}$ which we multiply by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Measured with $\psi(2S) \rightarrow \pi^+\pi^- J/\psi, J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$. $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

³ HAYRAPETYAN 24A reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (16.9^{+5.5}_{-4.6} \pm 0.6) \times 10^{-6}$ which we multiply by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{10}/Γ

NODE=M070Q74
NODE=M070Q74

 $\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma(\mu^+\mu^-)$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

18.7±1.8 OUR AVERAGE

18.9±1.7±0.9	452	1 AAIJ	24AE LHCb	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$
16.9 ^{+5.5} _{-4.6} ±0.6	12	HAYRAPETYAN	24A CMS	$J/\psi \rightarrow \mu^+\mu^-\mu^+\mu^-$

¹ Includes prompt production and inclusive decays of b -hadrons.

 Γ_{10}/Γ_7

NODE=M070Q70
NODE=M070Q70

HADRONIC DECAYS $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.88 ±0.12 OUR AVERAGE

2.072±0.017±0.062	19.8k	1 ANASHIN	23 KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \rho^0\pi^0$
2.18 ± 0.19		2,3 AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \rho^0\pi^0\gamma$
2.184±0.005±0.201	220k	3,4 BAI	04H BES	$e^+e^- \rightarrow J/\psi \rightarrow \rho^0\pi^0\pi^0$
2.091±0.021±0.116		3,5 BAI	04H BES	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
1.21 ± 0.20		BAI	96D BES	$e^+e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN	88 MRK3	$e^+e^- \rightarrow \rho\pi$
1.3 ± 0.3	150	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \rho\pi$
1.6 ± 0.4	183	ALEXANDER	78 PLUT	$e^+e^- \rightarrow \rho\pi$
1.33 ± 0.21		BRANDELIK	78B DASP	$e^+e^- \rightarrow \rho\pi$
1.0 ± 0.2	543	BARTEL	76 CNTR	$e^+e^- \rightarrow \rho\pi$
1.3 ± 0.3	153	JEAN-MARIE	76 MRK1	$e^+e^- \rightarrow \rho\pi$

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0, J/\psi \rightarrow \rho^+\pi^-, J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.

² From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).

³ Not independent of their $B(\pi^+\pi^-\pi^0)$.

⁴ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

⁵ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

 Γ_{11}/Γ

NODE=M070R20
NODE=M070R20

OCCUR=2

NODE=M070R20;LINKAGE=A


NODE=M070R20;LINKAGE=AU

NODE=M070R20;LINKAGE=BU

NODE=M070R20;LINKAGE=BA

NODE=M070R20;LINKAGE=BI

NODE=M070305

 $\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.142±0.011±0.026	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

1.331±0.033 20k 2 LEES 17C BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

1 From a Dalitz plot analysis in an isobar model.

2 From a Dalitz plot analysis in a Veneziano model.

 Γ_{11}/Γ_{170} NODE=M070P18
NODE=M070P18 $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.328±0.005±0.027	COFFMAN 88	MRK3	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ± 0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL 76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE 76	MRK1	e^+e^-

 Γ_{12}/Γ_{11} NODE=M070R21
NODE=M070R21 $\Gamma(a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ NODE=M070Q39
NODE=M070Q39

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.84±0.08±0.60	1317	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ NODE=M070Q40
NODE=M070Q40

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.67±0.09±0.73	1628	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$ Γ_{15}/Γ NODE=M070R43
NODE=M070R43

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.2 OUR AVERAGE				
11.7±0.7±2.5	7584	AUGUSTIN 89	DM2	$J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
8.4±4.5	36	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ NODE=M070P81
NODE=M070P81

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.68	471	1 ABLIKIM 19Q BES3	$e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$	

1 From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$. $\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{17}/Γ NODE=M070R22
NODE=M070R22

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN 88	MRK3	$e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{20}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$
$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{21}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$
$\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$					Γ_{23}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$21.8 \pm 2.2 \pm 3.4$	232 ± 23	ABLIKIM	08E	BES2	$e^+ e^- \rightarrow J/\psi$
$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{24}/Γ
<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM	10C	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$
$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$					Γ_{25}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.1 ± 0.8 OUR AVERAGE		Error includes scale factor of 1.6.			
7.90 $\pm 0.19 \pm 0.49$	3476	¹ ABLIKIM	17AK	BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
8.3 $\pm 3.0 \pm 1.2$	19	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 $\pm 1.4 \pm 1.6$		COFFMAN	88	MRK3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
1 From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.					
$\Gamma(\rho^+ K^+ K^- \pi^- + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{28}/Γ
<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$3.53 \pm 0.16 \pm 0.81$	485	ANASHIN	22	KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$\Gamma(h_1(1415)\eta' \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$					Γ_{30}/Γ
<u>VALUE</u> (units 10^{-5})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.08 $\pm 0.01^{+0.01}_{-0.02}$		¹ ABLIKIM	22AS	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
1 From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta'$ P -wave.					
$\Gamma(h_1(1595)\eta' \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$					Γ_{31}/Γ
<u>VALUE</u> (units 10^{-5})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.16 $\pm 0.02^{+0.03}_{-0.01}$		¹ ABLIKIM	22AS	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
1 From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta'$ P -wave.					
$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{33}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.2 \pm 0.2 \pm 1.1$	19.8k	¹ ANASHIN	23	KEDR	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
1 By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.					
$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{33}/Γ_{170}
<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
10.9 $\pm 1.7 \pm 2.7$	20k	¹ LEES	17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
0.80 ± 0.27	20k	² LEES	17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
1 From a Dalitz plot analysis in an isobar model.					
2 From a Dalitz plot analysis in a Veneziano model.					
$\Gamma(\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$					Γ_{34}/Γ_{188}
<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$6.3 \pm 0.8 \pm 0.6$	4k	¹ LEES	17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
1 From a Dalitz plot analysis in an isobar model.					

$\Gamma(\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{35}/Γ_{187}
9.3±2.0±0.6	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$	

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+ \pi^- \eta'(958)) / \Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ
3.28±0.55±0.44	119	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$	

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

 $\Gamma(\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{38}/Γ_{170}
8±2±5	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

22±6	20k	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
------	-----	--------	----------	--

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

 $\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ_{170}
4± 1±20	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

600±250	20k	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
---------	-----	--------	----------	--

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

 $\Gamma(\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{41}/Γ_{170}
• • • We do not use the following data for averages, fits, limits, etc. • • •					

4.0±0.8	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
---------	-----	--------	----------	--

¹ From a Dalitz plot analysis in a Veneziano model.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{42}/Γ
0.45 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.	

0.538±0.012±0.065 2090 1 ABLIKIM 06F BES2 $J/\psi \rightarrow \omega\pi^0$

0.360±0.028±0.054 222 JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$

0.482±0.019±0.064 COFFMAN 88 MRK3 $e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.



NODE=M070P27

NODE=M070P27

NODE=M070P27;LINKAGE=A

NODE=M070P36

NODE=M070P36

NODE=M070P36;LINKAGE=A

NODE=M070P21

NODE=M070P21

OCCUR=2

NODE=M070P21;LINKAGE=A

NODE=M070P21;LINKAGE=B

NODE=M070P22

NODE=M070P22

OCCUR=2

NODE=M070P22;LINKAGE=A

NODE=M070P22;LINKAGE=B

NODE=M070P24

NODE=M070P24

NODE=M070P24;LINKAGE=A

NODE=M070R32

NODE=M070R32

NODE=M070R32;LINKAGE=BL

$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8±3±2	20k	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model and significance 4.9 σ .

 Γ_{43}/Γ_{170}

NODE=M070P23
NODE=M070P23

 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±1.0 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.


10.6±1.2±0.1	3531	1 ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8±1.6	215	BURMESTER	77D PLUT	e^+e^-
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{44}/Γ

NODE=M070R24
NODE=M070R24

WEIGHTED AVERAGE
8.5±1.0 (Error scaled by 1.3)

 $\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.4±0.3±0.7	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+\pi^-3\pi^0$

 Γ_{45}/Γ

NODE=M070S26
NODE=M070S26

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±0.6 OUR AVERAGE				

4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	e^+e^-
4.0±1.6	70	BURMESTER	77D PLUT	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------	----	----------	---------	---

 Γ_{47}/Γ

NODE=M070R28
NODE=M070R28

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ±0.20 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.

2.352±0.273	5k	1 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta$
1.44 ± 0.40 ± 0.14	13	2 AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta$

 Γ_{48}/Γ


NODE=M070R30
NODE=M070R30

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R30;LINKAGE=BL

NODE=M070R30;LINKAGE=EE



$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$				Γ_{51}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85±34	140	VANNUCCI	77	$e^+ e^- \rightarrow 3(\pi^+ \pi^-) \pi^0$

NODE=M070R26
NODE=M070R26

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{53}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.12±0.02±0.13	14k	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

NODE=M070P83
NODE=M070P83

¹ Using the decays $\omega \rightarrow \pi^+ \pi^- \pi^0$ and $\eta' \rightarrow \eta \pi^+ \pi^-$.

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$				Γ_{54}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89±0.18 OUR AVERAGE				
2.08±0.30±0.14	137	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
2.26±0.43	218	² ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
$1.8^{+1.0}_{-0.8} \pm 0.3$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.66±0.17±0.19		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi\eta'$

NODE=M070R31
NODE=M070R31

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

² Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

NODE=M070R31;LINKAGE=A
NODE=M070R31;LINKAGE=BL

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$				Γ_{55}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41±0.27±0.47		¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M070S27
NODE=M070S27

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

NODE=M070S27;LINKAGE=K

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$				Γ_{56}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.8±1.1±0.3		^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S25
NODE=M070S25

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.

NODE=M070S25;LINKAGE=F
NODE=M070S25;LINKAGE=G

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$				Γ_{57}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8^{+1.9}_{-1.6} \pm 1.7	111^{+31}_{-26}	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S5
NODE=M070S5

$\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$					Γ_{58}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.2 \times 10^{-4}$	90	1 VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<2.8 \times 10^{-4}$	90	1 FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

NODE=M070R29
NODE=M070R29

$\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$					Γ_{59}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.9 \times 10^{-6}$	95	ABLIKIM	13P	BES3 $J/\psi \rightarrow \gamma \pi^0 p\bar{p}$	

NODE=M070R29;LINKAGE=C

$\Gamma(\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{61}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
$<6.2 \times 10^{-5}$	1 ABLIKIM	19AC	BES3 $J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$		

NODE=M070P84
NODE=M070P84

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$					Γ_{62}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.52 \pm 0.30 \pm 0.01$	276	1 ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P84;LINKAGE=A

$\Gamma(\omega K^+ K^- \eta)/\Gamma_{\text{total}}$					Γ_{60}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.33 \pm 0.02 \pm 0.12$		ABLIKIM	24BQ	BES3 $e^+ e^- \rightarrow J/\psi(1S)$	I

NODE=M070Q38
NODE=M070Q38

$\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$					Γ_{63}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
34 ± 5 OUR AVERAGE					
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$	
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M070S1
NODE=M070S1

$\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$					Γ_{64}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
19 ± 4 OUR AVERAGE					
19.8 ± 2.1 ± 3.9		1 FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	
16 ± 10	22	FELDMAN	77	MRK1 $e^+ e^-$	

NODE=M070R27
NODE=M070R27

¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

NODE=M070R27;LINKAGE=B

$\Gamma(\omega K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{65}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
61 ± 9 OUR AVERAGE					
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E	BES2 $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$	
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E	BES2 $J/\psi \rightarrow \omega K^+ K^- \pi^0$	
53 ± 14 ± 14	530 ± 140	BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M070S2
NODE=M070S2

OCCUR=2

$\Gamma(\eta' K^{\pm} K^{\mp})/\Gamma_{\text{total}}$					Γ_{66}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.48 ± 0.13 OUR AVERAGE					
1.50 ± 0.02 ± 0.19		¹ ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
1.47 ± 0.03 ± 0.17		² ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
1 From $\eta' K^+ K^- \pi^0$.					
2 From $\eta' K_S^0 K^\pm \pi^\mp$.					

NODE=M070P48
NODE=M070P48

OCCUR=2

NODE=M070P48;LINKAGE=A
NODE=M070P48;LINKAGE=B

$\Gamma(\eta' K^* \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{67}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.66 ± 0.03 ± 0.21		¹ ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
1 From $\eta' K_S^0 K^\pm \pi^\mp$.					

NODE=M070P49;LINKAGE=A

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{68}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.16 \pm 0.12 \pm 0.29$	1.1k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$	
1 From $\eta' K_S^0 K^\pm \pi^\mp$.					

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$					Γ_{69}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.51 \pm 0.09 \pm 0.21$	1.0k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$	
1 From $\eta' K^+ K^- \pi^0$.					

$\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$					Γ_{70}/Γ
<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$4.69 \pm 0.80 \pm 0.74$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$		

1 From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.


$\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$					Γ_{72}/Γ_{188}
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$90.5 \pm 0.9 \pm 3.8$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	

1 From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{73}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

6.0 ± 0.8 OUR AVERAGE Error includes scale factor of 2.9. See the ideogram below.

$8.07 \pm 0.04 \pm 0.38$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
$4.57 \pm 0.17 \pm 0.70$	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$5.26 \pm 0.13 \pm 0.53$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$



$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{74}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.69 \pm 0.01 \pm 0.13$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	
-0.20					

NODE=M070P52
NODE=M070P52

NODE=M070P52;LINKAGE=A

NODE=M070P44
NODE=M070P44

NODE=M070P44;LINKAGE=A

NODE=M070Q28
NODE=M070Q28

NODE=M070Q28;LINKAGE=A

NODE=M070P30
NODE=M070P30

NODE=M070P30;LINKAGE=A

NODE=M070S15
NODE=M070S15

NODE=M070P79
NODE=M070P79

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$
 Γ_{74}/Γ_{187}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
92.4±1.5±3.4	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma_{\text{total}}$
 Γ_{76}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
3.96±0.15±0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33±0.12±0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
-----------	----	----------	---------	--

 $\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$
 Γ_{78}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.73±0.14±0.82		1 ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	2 ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
------	-----------	----------	---

1 Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value 2/3 for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

2 A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$
 Γ_{82}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.81±0.10±0.54	1559	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$
 Γ_{83}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
6.28±0.16±0.59		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(K^*(892)^\pm K^*(700)^\mp) / \Gamma_{\text{total}}$
 Γ_{85}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.18±0.94	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$
 Γ_{86}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<5	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
----	----	----------	---------	---

 $\Gamma(K^*(892)^\pm K^*(892)^\mp) / \Gamma_{\text{total}}$
 Γ_{87}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.19±0.11	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

 $\Gamma(K_1(1400)^\pm K^\mp) / \Gamma_{\text{total}}$
 Γ_{88}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±0.8±1.2		1 BAI	99C BES	$e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

 $\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0) / \Gamma(K^+ K^- \pi^0)$
 Γ_{90}/Γ_{187}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±1.1±0.7	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$
 Γ_{91}/Γ_{188}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.5±0.9	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P26

NODE=M070P26

NODE=M070P26;LINKAGE=A

NODE=M070S16

NODE=M070S16

NODE=M070S52

NODE=M070S52

NODE=M070S52;LINKAGE=A

NODE=M070S52;LINKAGE=AB

NODE=M070P71

NODE=M070P71

NODE=M070S74

NODE=M070S74

NODE=M070R46

NODE=M070R46

NODE=M070S73

NODE=M070S73

NODE=M070S35;LINKAGE=M3

NODE=M070P28

NODE=M070P28

NODE=M070P28;LINKAGE=A

NODE=M070P32

NODE=M070P32

NODE=M070P32;LINKAGE=A

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.3±0.9	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

 Γ_{93}/Γ_{187}

NODE=M070P29
NODE=M070P29

 $\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
7.1±1.3±1.2	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K_2^*(1430)K + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<40 × 10⁻⁴	90	VANNUCCI	77	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<66 \times 10^{-4}$	90	BRAUNSCH...	76	DASP $e^+ e^- \rightarrow K^\pm \bar{K}_2^{*0}$
----------------------	----	-------------	----	---

 $\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
2.69±0.04^{+0.25}_{-0.19}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
2.65±0.80±0.44	1094	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ± 2.6	40	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
---------------	----	----------	----	--

 $\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<29 × 10⁻⁴	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 $\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

1.1±0.1^{+0.6}_{-0.1}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
--	------	---------	----------	------------------------------------

 $\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
6.2±0.7^{+2.8}_{-1.4}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.0 × 10⁻³	90	1 BAI	99C BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

 $\Gamma(K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁷)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

8.54^{+1.07}_{-1.20}^{+2.35}_{-2.13}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	-----------	---

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<43 × 10⁻⁴	90	BRAUNSCH...	76	DASP $e^+ e^-$

 Γ_{93}/Γ_{187}

NODE=M070P29
NODE=M070P29

NODE=M070P29;LINKAGE=A

NODE=M070P33
NODE=M070P33

NODE=M070P33;LINKAGE=A

NODE=M070R45
NODE=M070R45

NODE=M070P76
NODE=M070P76

NODE=M070Q43
NODE=M070Q43

NODE=M070R48
NODE=M070R48

NODE=M070R47
NODE=M070R47

NODE=M070P77
NODE=M070P77

NODE=M070P78
NODE=M070P78

NODE=M070S34
NODE=M070S34

NODE=M070S34;LINKAGE=M2

NODE=M070P72
NODE=M070P72

NODE=M070R42
NODE=M070R42

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{109}/Γ

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$
$0.124 \pm 0.033 \pm 0.030$	35 ± 9		² ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.4	90		³ ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
<6.8	90		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94 ± 0.15 OUR AVERAGE				Error includes scale factor of 1.7.
$1.09 \pm 0.02 \pm 0.13$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
$0.78 \pm 0.03 \pm 0.12$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ± 0.9	23	FELDMAN	77 MRK1	$e^+ e^-$

 $\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
16.0 $\pm 1.0 \pm 3.0$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{113}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.74 ± 0.06 OUR AVERAGE				Error includes scale factor of 1.2.
$0.71 \pm 0.10 \pm 0.05$	99 ± 14	¹ ZHU	23 BELL	$e^+ e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$

¹ From a fit to the combined $\phi\eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

 $\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.5 OUR AVERAGE					Error includes scale factor of 2.2. See the ideogram below.
$5.10 \pm 0.03 \pm 0.32$		31k	ABLIKIM	16M BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$5.46 \pm 0.31 \pm 0.56$			ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$4.1 \pm 0.3 \pm 0.8$		167	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$3.08 \pm 0.34 \pm 0.36$			COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 13	90		VANNUCCI	77 MRK1	$e^+ e^-$

NODE=M070R33

NODE=M070R33

NODE=M070R33

OCCUR=2

NODE=M070R33;LINKAGE=A

NODE=M070R33;LINKAGE=C

NODE=M070R33;LINKAGE=B

NODE=M070R34

NODE=M070R34

NODE=M070R35

NODE=M070R35


NODE=M070R37

NODE=M070R37

NODE=M070R37;LINKAGE=A

NODE=M070R38

NODE=M070R38



$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±0.06±0.16	2.2k	1 ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2±0.9 OUR AVERAGE				Error includes scale factor of 1.9.
4.6±0.4±0.8		1 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

2.6±0.6 50 1 GIDAL 81 MRK2 $J/\psi \rightarrow K^+K^-K^+K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50±0.80±0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.50±0.24	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\phi f_0(980)\eta \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23±0.75±0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$

$\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.24±0.20 ^{+0.52} _{-0.22}	1 ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi\eta\pi^0$
2.74±0.13 ^{+0.15} _{-0.16}	2 ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi\eta\pi^0$
4.37±1.35	3,4 ABLIKIM	18D BES3	$J/\psi \rightarrow \phi\eta\pi^0$
5.0 ± 2.7 ± 2.5	5 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi\eta\pi^0$

¹ $J/\psi \rightarrow \phi a_0(980)$ electromagnetic decay.

² $J/\psi \rightarrow \phi f_0(980)$, $\phi a_0(980)$ mixing.

³ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

⁴ Superseded by ABLIKIM 24CB.

⁵ Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and K^*K loops.

NODE=M070P85
NODE=M070P85

NODE=M070P85;LINKAGE=A

NODE=M070R41
NODE=M070R41

NODE=M070R41;LINKAGE=A

NODE=M070S97
NODE=M070S97

NODE=M070S98
NODE=M070S98

NODE=M070R08
NODE=M070R08

NODE=M070S75
NODE=M070S75

OCCUR=2

NODE=M070S75;LINKAGE=B
NODE=M070S75;LINKAGE=C
NODE=M070S75;LINKAGE=A

NODE=M070S75;LINKAGE=D
NODE=M070S75;LINKAGE=AB

$\Gamma(\phi(1680)^0 \pi^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{123}/Γ	NODE=M070Q80 NODE=M070Q80
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>COMMENT</u>		
$6.66 \pm 0.26 \pm 1.1$	ABLIKIM	24CB	BES3	$J/\psi \rightarrow \phi\eta\pi^0$		
$\Gamma(X(2000)^0 \pi^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{124}/Γ	NODE=M070Q81 NODE=M070Q81
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>COMMENT</u>		
$1.70 \pm 0.19 \pm 0.48$	ABLIKIM	24CB	BES3	$J/\psi \rightarrow \phi\eta\pi^0$		
$\Gamma(h_1(1900)^0 \pi^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{125}/Γ	NODE=M070Q82 NODE=M070Q82
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>COMMENT</u>		
$8.44 \pm 0.35 \pm 1.4$	ABLIKIM	24CB	BES3	$J/\psi \rightarrow \phi\eta\pi^0$		
$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$					Γ_{126}/Γ	NODE=M070R39 NODE=M070R39
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.45	90	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$		
< 0.37	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$		
$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$					Γ_{127}/Γ	NODE=M070S6 NODE=M070S6
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
2.6 ± 0.5 OUR AVERAGE						
3.4 ± 1.8 ± 1.5	1.1k	1 ABLIKIM	15H	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+ \pi^-$		
3.2 ± 0.6 ± 0.4		JOUSSET	90	DM2 $J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$		
2.1 ± 0.5 ± 0.4	25	2 JOUSSET	90	DM2 $J/\psi \rightarrow \phi\eta\pi^+ \pi^-$		OCCUR=2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.6 ± 0.2 ± 0.1	16	BECKER	87	MRK3 $J/\psi \rightarrow \phi K \bar{K} \pi$		
1 ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						NODE=M070S6;LINKAGE=A
2 We attribute to the $f_1(1285)$ the signal observed in the $\pi^+ \pi^- \eta$ invariant mass distribution at 1297 MeV.						NODE=M070S6;LINKAGE=Q
$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{128}/Γ	NODE=M070S99 NODE=M070S99
<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$9.36 \pm 2.31 \pm 1.54$	78	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$		
$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi 3\pi^0)/\Gamma_{\text{total}}$					Γ_{129}/Γ	NODE=M070S00 NODE=M070S00
<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.08 \pm 1.63 \pm 1.47$	9	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$		
$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{130}/Γ	NODE=M070S23 NODE=M070S23
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.01 \pm 0.58 \pm 0.82$		172	1 ABLIKIM	15H	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 17	90	2 FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$		
1 With 3.6 σ significance.						NODE=M070S23;LINKAGE=B
2 Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.						NODE=M070S23;LINKAGE=A
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$					Γ_{131}/Γ	NODE=M070R40 NODE=M070R40
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
8 ± 4 OUR AVERAGE Error includes scale factor of 2.7.						
12.3 ± 0.6 ± 2.0		1,2 FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$		
4.8 ± 1.8	46	1 GIDAL	81	MRK2 $J/\psi \rightarrow K^+ K^- K^+ K^-$		
1 Re-evaluated using $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$.						NODE=M070R40;LINKAGE=B
2 Including interference with $f_0(1710)$.						NODE=M070R40;LINKAGE=C

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{132}/Γ
$<2.1 \times 10^{-7}$	90	1 ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p} K_S^0 K_L^0$ $p\bar{p} K^+ K^-$	

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

 $\Gamma(\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{133}/Γ
$<2.8 \times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	

 $\Gamma(\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{134}/Γ
$<6.13 \times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	

 $\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{135}/Γ
--------------------------	------	-------------	------	---------	-----------------------


17.7 ± 1.6 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

16.6 ± 1.9 ± 1.2	163 ± 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
48 ± 20 ± 6	9.0 ± 3.7	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K\bar{K}$.

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

 $\Gamma(\phi f_0(1710) \rightarrow \phi K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{136}/Γ
$3.6 \pm 0.2 \pm 0.6$	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$	

¹ Including interference with $f'_2(1525)$.

² Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

 $\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{139}/Γ
7.2 ± 0.8 OUR AVERAGE					

7.4 ± 0.6 ± 1.4	227 ± 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070P00

NODE=M070P00

OCCUR=2

NODE=M070P00;LINKAGE=A

NODE=M070B10

NODE=M070B10

NODE=M070B11

NODE=M070B11

NODE=M070R36

NODE=M070R36

NODE=M070R36;LINKAGE=AA

NODE=M070R36;LINKAGE=CC

NODE=M070R36;LINKAGE=A

NODE=M070S24

NODE=M070S24

NODE=M070S24;LINKAGE=D

NODE=M070S24;LINKAGE=E

NODE=M070S3

NODE=M070S3

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{140}/Γ	NODE=M070S4 NODE=M070S4
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
21.8±2.3 OUR AVERAGE						
20.8±2.7±3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$		
29.6±3.7±4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^\mp K^\pm \pi^0$		OCCUR=2
20.7±2.4±3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$		
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$		
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$					Γ_{141}/Γ	NODE=M070R49 NODE=M070R49
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
30±5 OUR AVERAGE						
31±6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$		
29±7	87	BURMESTER	77D PLUT	$e^+ e^-$		
$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{142}/Γ	NODE=M070S28 NODE=M070S28
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
23±3±5	229	AUGUSTIN	89 DM2	$e^+ e^-$		
$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$					Γ_{144}/Γ	NODE=M070S14 NODE=M070S14
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.1 × 10⁻³	90	HENRARD	87 DM2	$e^+ e^-$		
$\Gamma(\Delta(1232)^{++} \bar{p}\pi^-)/\Gamma_{\text{total}}$					Γ_{145}/Γ	NODE=M070R70 NODE=M070R70
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.58±0.23±0.40	332	EATON	84 MRK2	$e^+ e^-$		
$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$					Γ_{146}/Γ	NODE=M070R66 NODE=M070R66
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.10±0.09±0.28	233	EATON	84 MRK2	$e^+ e^-$		
$\Gamma(\bar{\Sigma}(1385)^0 p K^-)/\Gamma_{\text{total}}$					Γ_{147}/Γ	NODE=M070R74 NODE=M070R74
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.51±0.26±0.18	89	EATON	84 MRK2	$e^+ e^-$		
$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{148}/Γ	NODE=M070S13 NODE=M070S13
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.82 × 10⁻⁵	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.2 × 10 ⁻³	90	HENRARD	87 DM2	$e^+ e^-$		
$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{149}/Γ	NODE=M070R68 NODE=M070R68
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.30±0.07 OUR AVERAGE						
0.30±0.03±0.08	74 ± 8	HENRARD	87 DM2	$e^+ e^-$		
0.29±0.11±0.10	26	EATON	84 MRK2	$e^+ e^-$		
$\Gamma(\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{150}/Γ	NODE=M070Q47 NODE=M070Q47
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.33±0.08 OUR AVERAGE						
0.34±0.04±0.08	77	HENRARD	87 DM2	$e^+ e^-$		
0.31±0.11±0.11	28	EATON	84 MRK2	$e^+ e^-$		
$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{151}/Γ	NODE=M070R67 NODE=M070R67
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.08 ± 0.06 OUR AVERAGE						
1.096±0.012±0.071	43k	ABLIKIM	16L BES3	$e^+ e^-$		
1.23 ± 0.07 ± 0.30	0.8k	ABLIKIM	12P BES2	$e^+ e^-$		
1.00 ± 0.04 ± 0.21	0.6k	HENRARD	87 DM2	$e^+ e^-$		
0.86 ± 0.18 ± 0.22	56	EATON	84 MRK2	$e^+ e^-$		


$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{152}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070Q48 NODE=M070Q48
1.25 ± 0.07 OUR AVERAGE						
1.258 ± 0.014 ± 0.078	53k	ABLIKIM	16L	BES3 $e^+ e^-$		
1.50 ± 0.08 ± 0.38	1k	ABLIKIM	12P	BES2 $e^+ e^-$		
1.19 ± 0.04 ± 0.25	0.7k	HENRARD	87	DM2 $e^+ e^-$		
1.03 ± 0.24 ± 0.25	68	EATON	84	MRK2 $e^+ e^-$		
$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$					Γ_{153}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P17 NODE=M070P17
1.071 ± 0.009 ± 0.082	103k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$		OCCUR=2
$\Gamma(\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{154}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S77 NODE=M070S77
<4.1 × 10⁻⁶	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda \bar{\Lambda} \gamma$		
$\Gamma(\bar{\Lambda}(1520) \Lambda + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{155}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P60 NODE=M070P60
<1.80 × 10⁻³	90	LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$		
$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$					Γ_{156}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S64 NODE=M070S64
1.17 ± 0.04 OUR AVERAGE						
1.165 ± 0.004 ± 0.043	135k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$		
1.20 ± 0.12 ± 0.21	206	ABLIKIM	080	BES2 $e^+ e^- \rightarrow J/\psi$		
$\Gamma(\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{157}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S9 NODE=M070S9
0.318 ± 0.008 OUR AVERAGE						
0.317 ± 0.002 ± 0.008	70k	ABLIKIM	20	BES3 $e^+ e^- \rightarrow J/\psi$		
0.59 ± 0.09 ± 0.12	75	HENRARD	87	DM2 $e^+ e^-$		
$\Gamma(\Xi(1530)^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$					Γ_{158}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S10 NODE=M070S10
0.32 ± 0.12 ± 0.07	24 ± 9	HENRARD	87	DM2 $e^+ e^-$		
$\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{159}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S47 NODE=M070S47
<1.1 × 10⁻⁵	90	BAI	04G	BES2 $e^+ e^-$		
$\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{160}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S48 NODE=M070S48
<2.1 × 10⁻⁵	90	BAI	04G	BES2 $e^+ e^-$		
$\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{161}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S49 NODE=M070S49
<1.6 × 10⁻⁵	90	BAI	04G	BES2 $e^+ e^-$		
$\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{162}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S50 NODE=M070S50
<5.6 × 10⁻⁵	90	BAI	04G	BES2 $e^+ e^-$		
$\Gamma(\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{163}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S51 NODE=M070S51
<1.1 × 10⁻⁵	90	BAI	04G	BES2 $e^+ e^-$		

STABLE HADRONS **$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{164}/Γ
4.2 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.	
5.44 ± 0.07 ± 0.33	23K	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$	
3.5 ± 0.8 ± 0.1	14k	1 LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$	
4.73 ± 0.44	228k	2 ABLIKIM	19Q	BES3 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$	
3.25 ± 0.49	46055	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$	
3.17 ± 0.42	147	FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$	
3.64 ± 0.52	1500	BURMESTER	77D	PLUT e^+e^-	
4 ± 1	675	JEAN-MARIE	76	MRK1 e^+e^-	

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3} \text{ keV}$ which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04 \text{ keV}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) = (18.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$, assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

 **$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{165}/Γ
0.029±0.006 OUR AVERAGE					
0.028 ± 0.009	11	FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$	
0.029 ± 0.007	181	JEAN-MARIE	76	MRK1 e^+e^-	

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{170}/Γ
20.0 ± 0.7 OUR AVERAGE				Error includes scale factor of 2.0. See the ideogram below.	
18.78 ± 0.13 ± 0.51	19.8k	1 ANASHIN	23	KEDR $e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
21.37 ± 0.04 ± 0.64	1.8M	2 ABLIKIM	12H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
23.0 ± 2.0 ± 0.4	256	3 AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$	
21.84 ± 0.05 ± 2.01	220k	4,5 BAI	04H	BES e^+e^-	
20.91 ± 0.21 ± 1.16		5,6 BAI	04H	BES e^+e^-	
15 ± 2	168	FRANKLIN	83	MRK2 e^+e^-	

NODE=M070307

NODE=M070R9

NODE=M070R9

NODE=M070R9;LINKAGE=B

NODE=M070R9;LINKAGE=A

NODE=M070R11


NODE=M070R11

NODE=M070R7

NODE=M070R7

OCCUR=2

- ¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.
- ² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.
- ³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.014$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.
- ⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.
- ⁶ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+\pi^-\pi^0 K^+ K^-)/\Gamma_{\text{total}}$		Γ_{172}/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52±0.27 OUR AVERAGE		Error includes scale factor of 1.4.		
1.74±0.08±0.24	2616	ANASHIN	22	$J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$
1.2 ± 0.3	309	VANNUCCI	77	e^+e^-

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_{173}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE		Error includes scale factor of 1.2.		
1.47±0.13±0.13	140	1 METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ± 0.5	5	BRANDELIK	78B	DASP
1.6 ± 1.6	1	VANNUCCI	77	MRK1

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$		Γ_{174}/Γ		
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.20±0.25 OUR AVERAGE		Error includes scale factor of 1.2.		
2.88±0.14±0.24	2654	ANASHIN	22	$J/\psi \rightarrow 2(\pi^+\pi^-)$
3.53±0.12±0.29	1107	1 ABLIKIM	05H	$BES2$
				$e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\pi^+\pi^-$, $J/\psi \rightarrow 2(\pi^+\pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1
				e^+e^-

¹ Computed using $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

NODE=M070R7;LINKAGE=B

NODE=M070R7;LINKAGE=AB

NODE=M070R7;LINKAGE=AU

NODE=M070R;LINKAGE=BA

NODE=M070R;LINKAGE=BU

NODE=M070R;LINKAGE=BI

NODE=M070R18

NODE=M070R18

NODE=M070R6

NODE=M070R6

NODE=M070R6;LINKAGE=ME

NODE=M070R8

NODE=M070R8

NODE=M070R8;LINKAGE=AB

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

40±20 32 JEAN-MARIE 76 MRK1 e^+e^-

 Γ_{175}/Γ

NODE=M070R10
NODE=M070R10

 $\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

90±30 13 JEAN-MARIE 76 MRK1 e^+e^-

 Γ_{177}/Γ

NODE=M070R12
NODE=M070R12

 $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.29±0.28 OUR AVERAGE

3.1 ± 1.5 ± 0.1 14k ¹ LEES 21 BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
 2.26 ± 0.08 ± 0.27 4.8k ABLIKIM 05C BES2 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta$
¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best values $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 Γ_{178}/Γ

NODE=M070S42
NODE=M070S42

 $\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

7.24±0.96±1.11 616 ABLIKIM 05C BES2 $e^+e^- \rightarrow 3(\pi^+\pi^-)\eta$

 Γ_{179}/Γ

NODE=M070S43
NODE=M070S43

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

3.06 ± 0.05 OUR AVERAGE

3.072 ± 0.023 ± 0.050 1.8 k ABLIKIM 24AB BES3 $\psi(2S) \rightarrow \pi^+\pi^- K^+K^-$
 2.86 ± 0.09 ± 0.19 1k ² METREVELI 12 $\psi(2S) \rightarrow \pi^+\pi^- K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.39 ± 0.24 ± 0.22 107 ³ BALTRUSAIT..85D MRK3 e^+e^-
 2.2 ± 0.9 6 ³ BRANDELIK 79C DASP e^+e^-

 Γ_{183}/Γ

NODE=M070R13
NODE=M070R13

¹ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.916 \pm 0.033)\%$.

² Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

³ Interference with non-resonant K^+K^- production not taken into account.

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{184}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

1.95±0.11 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.

1.93 ± 0.01 ± 0.05 110k ABLIKIM 17AH BES3 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^- X$
 2.62 ± 0.15 ± 0.14 0.3k ¹ METREVELI 12 $\psi(2S) \rightarrow \pi^+\pi^- K_S^0 K_L^0$
 1.82 ± 0.04 ± 0.13 2.1k ² BAI 04A BES2 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.18 ± 0.12 ± 0.18 JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$
 1.01 ± 0.16 ± 0.09 74 BALTRUSAIT..85D MRK3 e^+e^-

NODE=M070R13;LINKAGE=D

NODE=M070R13;LINKAGE=ME

NODE=M070R13;LINKAGE=BA

NODE=M070R75


NODE=M070R75

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.

NODE=M070R75;LINKAGE=ME

NODE=M070R;LINKAGE=HZ

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-8}$	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$
$<1 \times 10^{-6}$	95	¹ BAI	04D BES	$e^+ e^-$
$<5.2 \times 10^{-6}$	90	¹ BALTRUSAIT..85C	MRK3	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{185}/Γ NODE=M070R14
NODE=M070R14 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ±10 OUR AVERAGE				
55.2±12.0	25	FRANKLIN	83	$e^+ e^- \rightarrow K^+ K^- \pi^0$
78.0±21.0	126	VANNUCCI	77	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

 Γ_{186}/Γ

NODE=M070R14;LINKAGE=C

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.01±0.12	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 Γ_{187}/Γ NODE=M070R15
NODE=M070R15 $\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23k	LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

 $\Gamma_{187}/\Gamma_{170}$ NODE=M070P34
NODE=M070P34 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24k	LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

 $\Gamma_{188}/\Gamma_{170}$ NODE=M070P35
NODE=M070P35 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.04±0.26±0.92	2671	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 Γ_{192}/Γ NODE=M070R16
NODE=M070R16 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
31±13	30	VANNUCCI	77	MRK1 $e^+ e^-$

 Γ_{204}/Γ NODE=M070R17
NODE=M070R17

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ **Γ_{206}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.4 ^{+0.5} _{-0.4} ± 0.2	11.0 ^{+4.3} _{-3.5}	1 HUANG 03	BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$
0.7 ± 0.3		VANNUCCI 77	MRK1	$e^+ e^-$

¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

NODE=M070R19
NODE=M070R19

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ **Γ_{214}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.120± 0.029 OUR AVERAGE				
2.112 ± 0.004 ± 0.031	314k	ABLIKIM	12C	BES3 $e^+ e^-$
2.17 $\pm 0.16 \pm 0.04$	317	¹ WU	06	BELL $B^+ \rightarrow p\bar{p} K^+$
2.26 $\pm 0.01 \pm 0.14$	63316	BAI	04E	BES2 $e^+ e^- \rightarrow J/\psi$
1.97 ± 0.22	99	BALDINI	98	FENI $e^+ e^-$
1.91 $\pm 0.04 \pm 0.30$		PALLIN	87	DM2 $e^+ e^-$
2.16 $\pm 0.07 \pm 0.15$	1420	EATON	84	MRK2 $e^+ e^-$
2.5 ± 0.4	133	BRANDELIK	79C	DASP $e^+ e^-$
2.0 ± 0.5		BESCH	78	BONA $e^+ e^-$
2.2 ± 0.2	331	² PERUZZI	78	MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3 48 ANTONELLI 93 SPEC $e^+ e^-$

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ **Γ_{215}/Γ**


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19± 0.08 OUR AVERAGE Error includes scale factor of 1.1.				
1.33 $\pm 0.02 \pm 0.11$	11k	ABLIKIM	09B	BES2 $e^+ e^-$
1.13 $\pm 0.09 \pm 0.09$	685	EATON	84	MRK2 $e^+ e^-$
1.4 ± 0.4		BRANDELIK	79C	DASP $e^+ e^-$
1.00 ± 0.15	109	PERUZZI	78	MRK1 $e^+ e^-$

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{216}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
6.46 $\pm 0.17 \pm 0.43$	1435	EATON	84	MRK2 $e^+ e^-$
3.8 ± 1.6	48	BESCH	81	BONA $e^+ e^-$
5.5 ± 0.6	533	PERUZZI	78	MRK1 $e^+ e^-$

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)

NODE=M070R54
NODE=M070R54



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9.
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

 Γ_{217}/Γ

NODE=M070R55

NODE=M070R55

NODE=M070R55

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{218}/Γ

NODE=M070R56

NODE=M070R56

2.00±0.12 OUR AVERAGE

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.91 ± 0.02 ± 0.17	13k	¹ ABLIKIM	09	BES2 e^+e^-
$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79c	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels. $\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$ Γ_{219}/Γ

NODE=M070R57

NODE=M070R57

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.31 × 10⁻³	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$


 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{220}/Γ

NODE=M070R58

NODE=M070R58

0.98±0.10 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
0.98±0.10 (Error scaled by 1.3) $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ (units 10^{-3}) $\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$ Γ_{221}/Γ

NODE=M070R59

NODE=M070R59

0.129±0.014 OUR AVERAGE Error includes scale factor of 2.0.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.129 ± 0.014 OUR AVERAGE				Error includes scale factor of 2.0.
$0.126 \pm 0.002 \pm 0.007$	16k	¹ ABLIKIM	19N	BES3 e^+e^-
$0.200 \pm 0.023 \pm 0.028$	265 ± 31	² ABLIKIM	09	BES2 e^+e^-
$0.68 \pm 0.23 \pm 0.17$	19	EATON	84	MRK2 e^+e^-
1.8 ± 0.6	19	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.² From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels. $\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{222}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.8 ± 1.2 ± 1.3	ABLIKIM	14N	BES3 $e^+e^- \rightarrow J/\psi$


NODE=M070R59;LINKAGE=A

NODE=M070R59;LINKAGE=AB

NODE=M070S94

NODE=M070S94

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$				Γ_{223}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.519 ± 0.033 OUR AVERAGE					NODE=M070S22 NODE=M070S22
$0.523 \pm 0.006 \pm 0.033$	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p} K_S^0 K_L^0$	
$0.45 \pm 0.13 \pm 0.07$		FALVARD	88	DM2 $p\bar{p} K^+ K^-$ $J/\psi \rightarrow \text{hadrons}$	
$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$				Γ_{224}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.12 ± 0.09 OUR AVERAGE					NODE=M070R53 NODE=M070R53
$2.36 \pm 0.02 \pm 0.21$	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow p\pi^- \bar{n}$	
$2.47 \pm 0.02 \pm 0.24$	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{p}\pi^+ n$	OCCUR=2
$2.02 \pm 0.07 \pm 0.16$	1288	EATON	84	MRK2 $e^+ e^- \rightarrow p\pi^-$	OCCUR=2
$1.93 \pm 0.07 \pm 0.16$	1191	EATON	84	MRK2 $e^+ e^- \rightarrow \bar{p}\pi^+$	OCCUR=2
1.7 ± 0.7	32	BESCH	81	BONA $e^+ e^- \rightarrow p\pi^-$	OCCUR=2
1.6 ± 1.2	5	BESCH	81	BONA $e^+ e^- \rightarrow \bar{p}\pi^+$	OCCUR=2
2.16 ± 0.29	194	PERUZZI	78	MRK1 $e^+ e^- \rightarrow p\pi^-$	OCCUR=2
2.04 ± 0.27	204	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \bar{p}\pi^+$	OCCUR=2
$\Gamma(n\bar{n})/\Gamma_{\text{total}}$				Γ_{225}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.09 ± 0.16 OUR AVERAGE					NODE=M070R64 NODE=M070R64
$2.07 \pm 0.01 \pm 0.17$	36k	ABLIKIM	12C	BES3 $e^+ e^-$	
2.31 ± 0.49	79	BALDINI	98	FENI $e^+ e^-$	
1.8 ± 0.9		BESCH	78	BONA $e^+ e^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
1.90 ± 0.55	40	ANTONELLI	93	SPEC $e^+ e^-$	
$\Gamma(n\bar{n}\pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{226}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.8 ± 3.6	5	BESCH	81	BONA $e^+ e^-$	NODE=M070R65 NODE=M070R65
$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$				Γ_{230}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.88 ± 0.08 OUR AVERAGE		Error includes scale factor of 2.6. See the ideogram below.			
$1.64 \pm 0.12 \pm 0.09$		ABLIKIM	23S	BES3 $e^+ e^- \rightarrow \gamma\Lambda\bar{\Lambda}$	
$1.943 \pm 0.003 \pm 0.033$	441k	ABLIKIM	17L	BES3 $e^+ e^-$	
$2.03 \pm 0.03 \pm 0.15$	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$	
$1.96^{+0.47}_{-0.44} \pm 0.04$	46	¹ WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$	
$1.08 \pm 0.06 \pm 0.24$	631	BAI	98G	BES $e^+ e^-$	
$1.38 \pm 0.05 \pm 0.20$	1847	PALLIN	87	DM2 $e^+ e^-$	
$1.58 \pm 0.08 \pm 0.19$	365	EATON	84	MRK2 $e^+ e^-$	
2.6 ± 1.6	5	BESCH	81	BONA $e^+ e^-$	
1.1 ± 0.2	196	PERUZZI	78	MRK1 $e^+ e^-$	
$1^1 WU$ 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78\pm0.27\pm0.30		323	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6.4		90	² ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$
23 \pm 7 \pm 8		11	BAI	98G	BES e^+e^-
22 \pm 5 \pm 5		19	HENRARD	87	DM2 e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

Γ_{231}/Γ

NODE=M070S11
NODE=M070S11

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30\pm0.13\pm0.99	2.4k	ABLIKIM	12P	BES2 J/ψ

Γ_{232}/Γ

NODE=M070S11;LINKAGE=AL

NODE=M070S11;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

Γ_{233}/Γ

NODE=M070S78
NODE=M070S78

16.2 \pm 1.7 OUR AVERAGE

15.7 \pm 0.80 \pm 1.54	454	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2 \pm 6.0 \pm 4.4	44	² ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Sigma}^-\pi^++\text{c.c.})/\Gamma_{\text{total}}$

Γ_{234}/Γ

NODE=M070R07
NODE=M070R07

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.26 \pm0.05 OUR AVERAGE				Error includes scale factor of 1.2.

1.244 \pm 0.002 \pm 0.045	2.6M	ABLIKIM	23BUBES3	e^+e^-
1.52 \pm 0.08 \pm 0.16	589	¹ ABLIKIM	07H	BES2 e^+e^-
1.11 \pm 0.06 \pm 0.20	342 \pm 18	HENRARD	87	DM2 e^+e^-
1.38 \pm 0.21 \pm 0.35	118	EATON	84	MRK2 e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(\Lambda\bar{\Sigma}^+\pi^-+\text{c.c.})/\Gamma_{\text{total}}$

Γ_{235}/Γ

NODE=M070R71
NODE=M070R71

OCCUR=3

OCCUR=2

OCCUR=2

NODE=M070R71;LINKAGE=AB

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 \pm0.07 OUR AVERAGE				Error includes scale factor of 1.8.

1.221 \pm 0.002 \pm 0.038	2.7M	ABLIKIM	23BU BES3	e^+e^-
0.90 \pm 0.06 \pm 0.16	225	HENRARD	87	DM2 e^+e^-
1.53 \pm 0.17 \pm 0.38	135	EATON	84	MRK2 e^+e^-

NODE=M070Q46
NODE=M070Q46

$\Gamma(pK^-\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}$	Γ_{236}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.11 OUR AVERAGE				
$0.84^{+0.17}_{-0.15} \pm 0.02$	45	1 LU	19 BELL	$B^+ \rightarrow p\Lambda K^+ K^+$

$0.89 \pm 0.07 \pm 0.14$ 307 EATON 84 MRK2 $e^+ e^-$

1 LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$	Γ_{237}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.06±0.05	90	EATON	84	MRK2 $e^+ e^-$

$\Gamma(pK_S^0\bar{\Sigma}^-+c.c.)/\Gamma_{\text{total}}$	Γ_{238}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.725±0.009±0.050	120k	1 ABLIKIM	24H BES3	$e^+ e^- \rightarrow J/\psi$

1 The branching fractions for the charge-conjugate channels are measured separately as $(1.361 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $pK_S^0\bar{\Sigma}^+$ and $(1.352 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $pK_S^0\bar{\Sigma}^-$.

$\Gamma(\bar{\Lambda}nK_S^0+c.c.)/\Gamma_{\text{total}}$	Γ_{239}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	1 ABLIKIM	08C BES2	$e^+ e^- \rightarrow J/\psi$

1 Using $B(\bar{\Lambda} \rightarrow p\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\Lambda\bar{\Sigma}+c.c.)/\Gamma_{\text{total}}$	Γ_{240}/Γ				
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					
2.74 ± 0.24 ± 0.22		234 ± 21	1 ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92 ± 0.22 ± 0.24		308 ± 24	2 ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\Lambda\bar{\Sigma}+c.c.)/\Gamma_{\text{total}}$	Γ_{240}/Γ				
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					
<18			2 HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15		90	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Lambda X$

1 ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

2 ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$	Γ_{241}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.04 OUR AVERAGE				
1.061 ± 0.004 ± 0.036	87k	ABLIKIM	21AT BES3	$J/\psi \rightarrow p\pi^0\bar{p}\pi^0$
1.50 ± 0.10 ± 0.22	399	ABLIKIM	08O BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$	Γ_{242}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.172±0.032 OUR AVERAGE				Error includes scale factor of 1.4.
1.164 ± 0.004 ± 0.023	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.33 ± 0.04 ± 0.11	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884	PALLIN	87 DM2	$e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$	Γ_{242}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.172±0.032 OUR AVERAGE				Error includes scale factor of 1.4.
2.4 ± 2.6	3	BESCH	81 BONA	$e^+ e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

$\Gamma(\Sigma^+\bar{\Sigma}^-\eta)/\Gamma_{\text{total}}$	Γ_{243}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.34±0.21±0.37	1821	ABLIKIM	22AY BES3	$J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-\eta$

NODE=M070R72
NODE=M070R72

NODE=M070R72;LINKAGE=A

NODE=M070Q73
NODE=M070Q73

NODE=M070Q73;LINKAGE=A

NODE=M070S56
NODE=M070S56

NODE=M070S56;LINKAGE=AB


NODE=M070R61;LINKAGE=AC

NODE=M070S09
NODE=M070S09

NODE=M070R63
NODE=M070R63

NODE=M070Q45
NODE=M070Q45

$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$	Γ_{244}/Γ				
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.97 ± 0.08 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.	
1.040 ± 0.006 ± 0.074	43k	ABLIKIM	16L	BES3 $J/\psi \rightarrow \Xi^-\bar{\Xi}^+$	
0.90 ± 0.03 ± 0.18	961	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Xi^-\bar{\Xi}^+$	
0.70 ± 0.06 ± 0.12	132	HENRARD	87	DM2 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$	
1.14 ± 0.08 ± 0.20	194	EATON	84	MRK2 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$	
1.4 ± 0.5	51	PERUZZI	78	MRK1 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$	



RADIATIVE DECAYS

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$	Γ_{245}/Γ				
VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.41±0.14 OUR FIT					Error includes scale factor of 1.3.
1.7 ± 0.4 OUR AVERAGE					Error includes scale factor of 1.5.
2.00 ± 0.31 ± 0.02	1	MITCHELL	09	CLEO	$e^+e^- \rightarrow \gamma X$
1.27 ± 0.36		GAISER	86	CBAL	$J/\psi \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.40 ± 0.33	2	ANASHIN	14	KEDR	$J/\psi \rightarrow \gamma\eta_c$
1 MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Statistical uncertainty only.					

NODE=M070310

NODE=M070R85
NODE=M070R85

NODE=M070R85;LINKAGE=MI

NODE=M070R85;LINKAGE=A

NODE=M070R81
NODE=M070R81

$\Gamma(3\gamma)/\Gamma_{\text{total}}$	Γ_{248}/Γ				
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6±2.2 OUR AVERAGE					
11.3 ± 1.8 ± 2.0	113 ± 18	ABLIKIM	13I	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ± 3 ± 2	24.2 ± 7.2	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<55	90	PARTTRIDGE	80	CBAL	e^+e^-

NODE=M070S06
NODE=M070S06

$\Gamma(4\gamma)/\Gamma_{\text{total}}$	Γ_{249}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<9 × 10 ⁻⁶	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<15 × 10 ⁻⁶	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

NODE=M070S07
NODE=M070S07

$\Gamma(5\gamma)/\Gamma_{\text{total}}$	Γ_{250}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<15 × 10 ⁻⁶	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$				Γ_{251}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.39 ± 0.08 OUR AVERAGE				
$3.34 \pm 0.02 \pm 0.09$	176k	ABLIKIM	23BD BES3	$J/\psi \rightarrow \pi^0\gamma$
$3.59 \pm 0.20 \pm 0.03$	1.6k	¹ ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$3.63 \pm 0.36 \pm 0.13$		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0\gamma$
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 1.1 \pm 0.7$	BLOOM	83 CBAL	e^+e^-
7.3 ± 4.7	10	BRANDELIK	79C DASP e^+e^-

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$		Γ_{252}/Γ	
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.05	¹ ABLIKIM	15AE BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$


1 The uncertainty is systematic as statistical is negligible.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$		Γ_{253}/Γ	
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.5 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.
$4.32 \pm 0.14 \pm 0.73$	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
$2.08 \pm 0.13 \pm 0.35$	² BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
$3.05 \pm 0.08 \pm 0.45$	² BALTRUSAIT...86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
$4.85 \pm 0.45 \pm 1.20$	³ BURKE	82 MRK2	e^+e^-

1 4π mass less than 3.0 GeV.

2 4π mass less than 2.0 GeV.

3 4π mass less than 2.5 GeV.



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$		Γ_{254}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.7 \pm 1.6$	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070R82
NODE=M070R82

NODE=M070R82;LINKAGE=A

NODE=M070B00
NODE=M070B00

NODE=M070B00;LINKAGE=A

NODE=M070R95
NODE=M070R95

OCCUR=2

NODE=M070R95;LINKAGE=A
NODE=M070R95;LINKAGE=B
NODE=M070R95;LINKAGE=M

NODE=M070S45
NODE=M070S45

$\Gamma(\gamma f_2(1270) f_2(1270) \text{(non resonant)})/\Gamma_{\text{total}}$				Γ_{255}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.2 \pm 0.8 \pm 1.7$	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$	NODE=M070S46 NODE=M070S46
1 Subtracting contribution from intermediate $\eta_c(1S)$ decays.				
$\Gamma(\gamma\pi^+\pi^- 2\pi^0)/\Gamma_{\text{total}}$				Γ_{256}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.3 \pm 0.2 \pm 3.1$	¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$	NODE=M070R99 NODE=M070R99
1 4π mass less than 2.0 GeV.				
$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{257}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.1 ± 0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	NODE=M070P73 NODE=M070P73
$\Gamma(\gamma(K\bar{K}\pi)[J^{PC}=0^- +])/ \Gamma_{\text{total}}$				Γ_{258}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.			
$0.58 \pm 0.03 \pm 0.20$	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$	
$2.1 \pm 0.1 \pm 0.7$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$	
1 For a broad structure around 1800 MeV. 2 For a broad structure around 2040 MeV.				
$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{259}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.1 \pm 0.1 \pm 0.6$	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$				Γ_{260}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
1 Summed over all charges.				
$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$				Γ_{261}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.090 ± 0.013 OUR AVERAGE				
$1.096 \pm 0.001 \pm 0.019$	2.2M	ABLIKIM	23BD BES3	$J/\psi \rightarrow \eta\gamma$
$1.067 \pm 0.005 \pm 0.023$	87.9k	ABLIKIM	21AMBES3	$e^+ e^- \rightarrow J/\psi$
$1.12 \pm 0.05 \pm 0.01$	18.6k	¹ ABLIKIM	180 BESS3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
$1.101 \pm 0.029 \pm 0.022$		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta\gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.88 $\pm 0.08 \pm 0.11$		BLOOM	83 CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77 CNTR	$e^+ e^-$
1 ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.				
$\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$				Γ_{262}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$21.4 \pm 1.8 \pm 2.5$	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$
$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$				Γ_{263}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.5 ± 2.0	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$		
$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$				Γ_{264}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5 ± 5	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$		

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4±3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 Γ_{265}/Γ

NODE=M070P98
NODE=M070P98

 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.5 \times 10^{-6}$	95	ABLIKIM	16P	$BES3 \quad J/\psi \rightarrow 5\gamma$

 Γ_{266}/Γ

NODE=M070P02
NODE=M070P02

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.6 \times 10^{-6}$	95	ABLIKIM	16P	$BES3 \quad J/\psi \rightarrow 5\gamma$

 Γ_{267}/Γ

NODE=M070P03
NODE=M070P03

 $\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

6.1 ±1.0 OUR AVERAGE

$5.85 \pm 0.3 \pm 1.05$ ¹ EDWARDS 83B CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$
 $7.8 \pm 1.2 \pm 2.4$ ¹ EDWARDS 83B CBAL $J/\psi \rightarrow \eta 2\pi^0$

¹ Broad enhancement at 1700 MeV.

 Γ_{268}/Γ

NODE=M070R96
NODE=M070R96

 $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.2 \pm 2.2 \pm 0.9$	BAI	99	$BES \quad J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

 Γ_{269}/Γ

NODE=M070S37
NODE=M070S37

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

5.28±0.06 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

5.40±0.01±0.11	638k	ABLIKIM	23BD	BES3 $J/\psi \rightarrow \gamma\eta'$
5.27±0.03±0.05	36k	ABLIKIM	19T	BES $J/\psi \rightarrow \gamma\eta'$
5.43±0.23±0.09	5.0k	¹ ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
4.76±0.22±0.06		² ABLIKIM	11	BES3 $J/\psi \rightarrow \eta'\gamma$
5.24±0.12±0.11		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
5.55±0.44	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •


4.50±0.14±0.53		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
4.30±0.31±0.71		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
4.04±0.16±0.85	622	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
4.39±0.09±0.66	2420	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
4.1 ± 0.3 ± 0.6		BLOOM	83	CBAL $e^+e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ± 1.1	6	BRANDELIK	79C	DASP $e^+e^- \rightarrow 3\gamma$
2.4 ± 0.7	57	BARTEL	76	CNTR $e^+e^- \rightarrow 2\gamma\rho$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R84;LINKAGE=A

NODE=M070R84;LINKAGE=AB

 $\Gamma(\gamma f_0(980) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 Γ_{271}/Γ

NODE=M070P90
NODE=M070P90

 $\Gamma(\gamma f_0(980) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8 ± 0.3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 Γ_{272}/Γ

NODE=M070P91
NODE=M070P91

 $\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

4.5 ± 0.8 OUR AVERAGE

$4.7 \pm 0.3 \pm 0.9$ 1 BALTRUSAIT..86B MRK3 $J/\psi \rightarrow 4\pi\gamma$
 $3.75 \pm 1.05 \pm 1.20$ 2 BURKE 82 MRK2 $J/\psi \rightarrow 4\pi\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09 90 3 BISELLO 89B $J/\psi \rightarrow 4\pi\gamma$

1.4π mass less than 2.0 GeV.

2.4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

3.4π mass in the range 2.0–25 GeV.

 Γ_{273}/Γ

NODE=M070R94
NODE=M070R94

 $\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<5.4 \times 10^{-4}$ 90 ABLIKIM 08A BES2 $e^+ e^- \rightarrow J/\psi$

 Γ_{274}/Γ

NODE=M070R05
NODE=M070R05

 $\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<8.8 \times 10^{-5}$ 90 ABLIKIM 08A BES2 $e^+ e^- \rightarrow J/\psi$

 Γ_{275}/Γ

NODE=M070R06
NODE=M070R06

 $\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 1.61 ± 0.33 OUR AVERAGE

$6.0 \pm 4.8 \pm 1.8$ ABLIKIM 08A BES2 $J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
 $1.41 \pm 0.2 \pm 0.42$ 120 ± 17 BISELLO 87 SPEC $e^+ e^-, \text{hadrons}\gamma$
 $1.76 \pm 0.09 \pm 0.45$ BALTRUSAIT..85C MRK3 $e^+ e^- \rightarrow \text{hadrons}\gamma$

 Γ_{276}/Γ

NODE=M070R97
NODE=M070R97


$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{277}/Γ
4.0±1.2 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.					
7.5±0.6±1.2	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
3.4±0.8±0.6	33 ± 7	1 BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
3.1±0.7±0.4		1 BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	

1 $\phi\phi$ mass less than 2.9 GeV, η_C excluded.

NODE=M070R98
NODE=M070R98

NODE=M070R98;LINKAGE=C

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_{278}/Γ
2.8 ±0.6 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.				
1.66±0.1 ±0.58	1,2 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$	
3.8 ±0.3 ±0.6	3 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	
4.0 ±0.7 ±1.0	3 EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$	
4.3 ±1.7	3,4 SCHARRE	80 MRK2	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.78±0.21±0.33	3,5,6 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	
0.83±0.13±0.18	3,7,8 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	OCCUR=2
0.66±0.17±0.24	3,6,9 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
0.16±0.15				OCCUR=2
1.03±0.21±0.26	3,8,10 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
-0.18±0.19				

1 Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

2 Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

3 Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

4 Corrected for spin-zero hypothesis for $\eta(1405)$.

5 From fit to the $a_0(980)\pi^-$ partial wave.

6 $a_0(980)\pi^-$ mode.

7 From fit to the $K^*(892)K^-$ partial wave.

8 $K^* K^-$ mode.

9 From $a_0(980)\pi^-$ final state.

10 From $K^*(890)K^-$ final state.

NODE=M070R89
NODE=M070R89

OCCUR=2

OCCUR=2

NODE=M070R89;LINKAGE=BD

NODE=M070R89;LINKAGE=BE

NODE=M070R89;LINKAGE=B

NODE=M070R89;LINKAGE=C

NODE=M070R89;LINKAGE=H


NODE=M070R89;LINKAGE=K9

NODE=M070R89;LINKAGE=J

NODE=M070R89;LINKAGE=K8

NODE=M070R89;LINKAGE=D

NODE=M070R89;LINKAGE=E

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{279}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78±0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07±0.17±0.11	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64±0.12±0.07	¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{280}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ±0.5 OUR AVERAGE				
2.6 ±0.7 ±0.4		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38±0.33±0.64		¹ BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ±0.6 ±1.1	261	² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
---------------	-----	-----------------------	--------	---

¹ Via $a_0(980)\pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{281}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ±0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ±0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36±0.38	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

1 Estimated by us from various fits.

2 Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{282}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<82	95		BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.03±0.92±0.91	1.3k	¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
10.36±1.51±1.54	1.9k	² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{283}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.11±0.11	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma\eta(1405)$

NODE=M070S30
NODE=M070S30

NODE=M070S30;LINKAGE=C

NODE=M070S29
NODE=M070S29

NODE=M070S29;LINKAGE=RR
NODE=M070S29;LINKAGE=R

NODE=M070S19
NODE=M070S19

NODE=M070S19;LINKAGE=A
NODE=M070S19;LINKAGE=B

NODE=M070R77
NODE=M070R77

OCCUR=2

NODE=M070R77;LINKAGE=B

NODE=M070R77;LINKAGE=A

NODE=M070Q76
NODE=M070Q76


$\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^0\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_{284}/Γ	NODE=M070Q77 NODE=M070Q77
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$7.10 \pm 0.82 \pm 0.72$	198	ABLIKIM
		TECN
		J/ ψ $\rightarrow \gamma\eta(1405)$
$\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{285}/Γ	NODE=M070P38 NODE=M070P38
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>
$<2.63 \times 10^{-6}$	90	ABLIKIM
		TECN
		$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{286}/Γ	NODE=M070P39 NODE=M070P39
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>
$<1.86 \times 10^{-6}$	90	ABLIKIM
		TECN
		$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$	Γ_{287}/Γ	NODE=M070S20 NODE=M070S20
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
0.13 ± 0.09	1,2 BISELLO	89B DM2
		TECN
		J/ ψ $\rightarrow 4\pi\gamma$
1 Estimated by us from various fits.		
2 Includes unknown branching fraction to $\rho^0\rho^0$.		
$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$	Γ_{288}/Γ	NODE=M070R04 NODE=M070R04
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM
		TECN
		J/ ψ $\rightarrow \gamma\omega\omega$
$\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{289}/Γ	NODE=M070P40 NODE=M070P40
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>
$<4.80 \times 10^{-6}$	90	ABLIKIM
		TECN
		$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$	Γ_{290}/Γ	NODE=M070S21 NODE=M070S21
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$3.14^{+0.50}_{-0.19}$ OUR AVERAGE		
		TECN
		COMMENT
2.40 $\pm 0.10^{+2.47}_{-0.18}$	1,2 ABLIKIM	16N BES3
4.4 ± 0.4 ± 0.8	196	2 ABLIKIM
3.3 ± 0.8 ± 0.5		2 BAI
2.7 ± 0.6 ± 0.6		2 BAI
2.4 ± 1.5	3,4 BISELLO	89B DM2
		TECN
		J/ ψ $\rightarrow 4\pi\gamma$
1 From a partial wave analysis of J/ $\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).		
2 Includes unknown branching fraction to $\phi\phi$.		
3 Estimated by us from various fits.		
4 Includes unknown branching fraction to $\rho^0\rho^0$.		
$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$	Γ_{291}/Γ	NODE=M070R86 NODE=M070R86
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
1.63 ± 0.12 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.
		TECN
2.07 $\pm 0.16^{+0.02}_{-0.07}$	2.4k	1,2 DOBBS
		15
		TECN
		J/ $\psi \rightarrow \gamma\pi\pi$
1.63 $\pm 0.26^{+0.02}_{-0.05}$		3 ABLIKIM
		06v BES2
		e ⁺ e ⁻ $\rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.42 $\pm 0.21^{+0.02}_{-0.05}$		4 ABLIKIM
		06v BES2
		e ⁺ e ⁻ $\rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
1.33 $\pm 0.05 \pm 0.20$		5 AUGUSTIN
		87 DM2
		J/ $\psi \rightarrow \gamma\pi^+\pi^-$
1.36 $\pm 0.09 \pm 0.23$		5 BALTRUSAIT..87
		MRK3
		J/ $\psi \rightarrow \gamma\pi^+\pi^-$
1.48 $\pm 0.25 \pm 0.30$	178	EDWARDS
		82B CBAL
		e ⁺ e ⁻ $\rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER
		78 PLUT
		e ⁺ e ⁻
1.2 ± 0.6	30	6 BRANDELIK
		78B DASP
		e ⁺ e ⁻ $\rightarrow \pi^+\pi^-\gamma$
1 Using CLEO-c data but not authored by the CLEO Collaboration.		
2 DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
3 ABLIKIM 06v reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi)$		

$= (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270)) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

Γ_{292}/Γ

NODE=M070P68
NODE=M070P68

$\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.61 ± 0.08 OUR AVERAGE			

$0.69 \pm 0.16 \pm 0.20$

$0.61 \pm 0.04 \pm 0.21$

$0.45 \pm 0.09 \pm 0.17$

$0.625 \pm 0.063 \pm 0.103$

$0.70 \pm 0.08 \pm 0.16$

1 Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.

2 Assuming $\Gamma(f_1(1285) \rightarrow K\bar{K}\pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$.

3 Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi) / \Gamma_{\text{total}} = 0.5 \pm 0.18$.

4 Obtained summing the sequential decay channels

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.

5 Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

Γ_{293}/Γ

NODE=M070R88
NODE=M070R88

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\pi\pi) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
38±10	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

Γ_{294}/Γ

NODE=M070P92
NODE=M070P92

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$	Γ_{295}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.19 \pm 0.73 \pm 1.34$	478	1 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.3 ± 0.4	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{296}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM	18AA BES3		$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$	Γ_{297}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.5 ± 1.0	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{298}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.9 ± 0.3	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$	Γ_{299}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.79 ± 0.13 OUR AVERAGE				
0.68 $\pm 0.04 \pm 0.24$	BAI	00D BES		$J/\psi \rightarrow \gamma K_S^{\pm} K_S^0 \pi^{\mp}$
0.76 $\pm 0.15 \pm 0.21$	1,2 AUGUSTIN	92 DM2		$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.87 $\pm 0.14^{+0.14}_{-0.11}$	1 BAI	90C MRK3		$J/\psi \rightarrow \gamma K_S^0 K_S^{\pm} \pi^{\mp}$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K\bar{K}\pi$.

² From fit to the $K^*(892)K$ 1^{++} partial wave.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$	Γ_{300}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.09 ± 0.24 OUR AVERAGE				
1.21 $\pm 0.29 \pm 0.24$	174	1 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.00 $\pm 0.03 \pm 0.45$		2 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.02 $\pm 0.09 \pm 0.45$		2 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90 ± 0.17	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
5.7 ± 0.8	3.4 BUGG	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+\pi^-\pi^+\pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$	Γ_{301}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	1 ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.1 ± 0.4	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-------------------	--	--	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{302}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.59^{+0.16+0.18}_{-0.56}$	ABLIKIM	18AA BES3		$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.3	SARANTSEV 21 RVUE			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-------------------	--	--	---

NODE=M070R00
NODE=M070R00

NODE=M070R00;LINKAGE=A

NODE=M070P63
NODE=M070P63

NODE=M070P93
NODE=M070P93

NODE=M070P94
NODE=M070P94

NODE=M070S31
NODE=M070S31

OCCUR=2

NODE=M070S31;LINKAGE=A
NODE=M070S31;LINKAGE=D

NODE=M070S32
NODE=M070S32

OCCUR=2

NODE=M070S32;LINKAGE=C
NODE=M070S32;LINKAGE=AB
NODE=M070S32;LINKAGE=A
NODE=M070S32;LINKAGE=B

NODE=M070S83
NODE=M070S83

NODE=M070S83;LINKAGE=A

NODE=M070P64
NODE=M070P64

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.1 \pm 1.1^{+1.9}_{-1.3}$ ¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$

12 ± 5 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

 Γ_{303}/Γ

NODE=M070P99

NODE=M070P99

 $\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.5 \pm 1.0 \pm 0.7$ BAI 99 BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

 Γ_{304}/Γ

NODE=M070S36

NODE=M070S36

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

5.7 $^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.0 \pm 0.9 \pm 0.2$	750	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3 BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4 90 4 ⁴ BRANDELIK 79c DASP $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
 <2.3 90 3 ALEXANDER 78 PLUT $e^+ e^- \rightarrow K^+ K^- \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.


⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

 Γ_{305}/Γ

NODE=M070R87

NODE=M070R87

WEIGHTED AVERAGE
5.7+0.8-0.5 (Error scaled by 1.5)

 $\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.99 $^{+0.03}_{-0.04}$ $^{+0.69}_{-0.50}$ ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

 Γ_{306}/Γ

NODE=M070P69

NODE=M070P69

$\Gamma(\gamma f_2'(1525) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$	Γ_{307}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1565) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{308}/Γ		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.32 \pm 0.05^{+0.12}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta'$ P -wave.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$	Γ_{309}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$	Γ_{310}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT

3.8 ± 0.5 OUR AVERAGE

$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06v BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06v BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 \pm 0.2		³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
2.5 \pm 1.6 \pm 0.8		BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$	Γ_{311}/Γ				
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT

9.5 ± 1.0 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.00 \pm 0.12 \pm 1.24	1.24		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
11.76 \pm 0.54 \pm 0.94	1.2k		² DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
9.62 \pm 0.29 \pm 3.51			³ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
5.0 \pm 0.8 \pm 1.8			^{1,4} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
9.2 \pm 1.4 \pm 1.4			¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
10.4 \pm 1.2 \pm 1.6			¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
9.6 \pm 1.2 \pm 1.8			¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 \pm 0.8			⁵ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
1.6 \pm 0.2 \pm 0.6			1.6 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$

< 0.8	90		⁷ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
1.6 \pm 0.4 \pm 0.3			⁸ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 \pm 1.6			⁹ EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta\eta\gamma$

¹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.

⁵ There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.

⁶ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁷ Includes unknown branching fraction to $\rho^0\rho^0$.

⁸ Includes unknown branching fraction to $\pi^+\pi^-$.

⁹ Includes unknown branching fraction to $\eta\eta$.

NODE=M070S86
NODE=M070S86


NODE=M070Q31;LINKAGE=A
NODE=M070Q31
NODE=M070Q31

NODE=M070R02
NODE=M070R02
NODE=M070B01
NODE=M070B01

NODE=M070B01;LINKAGE=A
NODE=M070B01;LINKAGE=AB
NODE=M070B01;LINKAGE=B
OCCUR=2

OCCUR=2

OCCUR=2
OCCUR=2
NODE=M070R91;LINKAGE=B
NODE=M070R91;LINKAGE=D
NODE=M070R91;LINKAGE=K9
NODE=M070R91;LINKAGE=A1
NODE=M070R91;LINKAGE=E
NODE=M070R91;LINKAGE=A2
NODE=M070R91;LINKAGE=C
NODE=M070R91;LINKAGE=Z
NODE=M070R91;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

 Γ_{312}/Γ

NODE=M070R01
NODE=M070R01

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	1 ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

 Γ_{313}/Γ

NODE=M070S84
NODE=M070S84

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.4	2 SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	---------------------	--

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{314}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.5 ± 2.5	1 SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	---------------------	--

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi)/\Gamma_{\text{total}}$ Γ_{315}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.5 ± 0.6 OUR AVERAGE

2.00 $\pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
---------------------------------	------	---------	----------	---

2.61 $\pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$
--------------------------	----	---------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1 ± 0.1	1 SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	---------------------	--

¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

 $\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{316}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	-----------	---

 $\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{317}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

$5.40 \pm 0.60^{+3.42}_{-0.67-2.35}$	5.5k	1 ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$
--	------	-----------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070P65
NODE=M070P65

NODE=M070S87
NODE=M070S87

NODE=M070S87;LINKAGE=A

$\Gamma(\gamma\eta_1(1855) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{318}/Γ
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$2.70 \pm 0.41 \begin{array}{l} +0.16 \\ -0.35 \end{array}$	¹ ABLIKIM 22AI BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave. For analysis details see ABLIKIM 22AS.	
$\Gamma(\gamma f_0(1770) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{319}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$0.11 \pm 0.01 \begin{array}{l} +0.04 \\ -0.03 \end{array}$	¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.	
$\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$	Γ_{320}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.20 \pm 0.04 \pm 0.13$	151 ABLIKIM 06H BES $J/\psi \rightarrow \gamma\omega\omega$
$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_{321}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.7 \pm 0.1 \pm 0.2$	BAI 00B BES $J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
$\Gamma(\gamma f_2(2010) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{322}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$0.71 \pm 0.06 \begin{array}{l} +0.10 \\ -0.06 \end{array}$	¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.	
$\Gamma(\gamma f_0(2020) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$	Γ_{323}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
42 ± 10	SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$\Gamma(\gamma f_0(2020) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$	Γ_{324}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
55 ± 25	SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$	Γ_{325}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
10 ± 10	SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$	Γ_{326}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$2.63 \pm 0.06 \begin{array}{l} +0.31 \\ -0.46 \end{array}$	¹ ABLIKIM 22C BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.	
$\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$	Γ_{327}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$2.28 \pm 0.12 \begin{array}{l} +0.29 \\ -0.20 \end{array}$	¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.	

$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$				Γ_{328}/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	
2.7±0.5±0.5	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1 Assuming branching fraction $f_4(2050) \rightarrow \pi\pi/\text{total} = 0.167$.				
$\Gamma(\gamma f_4(2050) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$				Γ_{329}/Γ
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.06±0.01 ^{+0.03} _{-0.01}	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$	
1 From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta' P$ -wave.				
$\Gamma(\gamma f_0(2100) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$				Γ_{330}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.13^{+0.09+0.64}_{-0.10-0.28}	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.8 ± 1.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.				
$\Gamma(\gamma f_0(2100) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$				Γ_{331}/Γ
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
32±20	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$				Γ_{332}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.24^{+0.48+0.87}_{-0.48-0.87}	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.0 ± 0.8	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
1 Using CLEO-c data but not authored by the CLEO Collaboration.				
$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$				Γ_{333}/Γ
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.5	¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	
1 Includes unknown branching fraction to $K_S^0 K_S^0$.				
$\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$				Γ_{334}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.86^{+0.49+1.20}_{-0.49-1.20}	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.5 ± 0.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
1 Using CLEO-c data but not authored by the CLEO Collaboration.				
$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{335}/Γ
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	
2.72^{+0.08+0.17}_{-0.06-0.47}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	
$\Gamma(\gamma f_0(2200) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$				Γ_{336}/Γ
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5±2	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$\Gamma(\gamma f_0(2200) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$				Γ_{337}/Γ
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.7±0.4	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2220))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
>300			1 BAI	96B BES	$e^+e^- \rightarrow \gamma p\bar{p}, K\bar{K}$
>250	99.9		2 HASAN	96 SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+K^-$
< 1.6	95		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$	23		3 BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$	93		3 BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+K^-$

¹ Using BARNEs 93.

² Using BAI 96B.

³ Includes unknown branching fraction to K^+K^- or $K_S^0 K_S^0$.

Γ_{338}/Γ

NODE=M070R92
NODE=M070R92

OCCUR=2

OCCUR=2

NODE=M070R92;LINKAGE=A

NODE=M070R92;LINKAGE=M

NODE=M070R92;LINKAGE=W

$\Gamma(\gamma f_0(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{339}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

$14 \pm 8 \pm 4$ BAI 98H BES $J/\psi \rightarrow \gamma\pi^0\pi^0$

$8.4 \pm 2.6 \pm 3.0$ BAI 96B BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

NODE=M070B02
NODE=M070B02

NODE=M070B02;LINKAGE=A

NODE=M070B02;LINKAGE=DO

$\Gamma(\gamma f_0(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{340}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

< 3.6 ³ DEL-AMO-SA..100 BABR $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$

< 2.9 ³ DEL-AMO-SA..100 BABR $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

$6.6 \pm 2.9 \pm 2.4$ BAI 96B BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$

$10.8 \pm 4.0 \pm 3.2$ BAI 96B BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070B03
NODE=M070B03

OCCUR=2

OCCUR=2

NODE=M070B03;LINKAGE=A

NODE=M070B03;LINKAGE=DO

NODE=M070B03;LINKAGE=DE

$\Gamma(\gamma f_0(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

Γ_{341}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$

NODE=M070B04
NODE=M070B04

$\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{342}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.95 \pm 0.21^{+0.66}_{-0.72}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P67
NODE=M070P67

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

0.6 ± 0.1 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{343}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

NODE=M070Q07
NODE=M070Q07

4 ± 2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{344}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

NODE=M070Q08
NODE=M070Q08

1.5 ± 0.4 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{345}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.09 \pm 0.64^{+4.00}_{-1.68}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta'X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{346}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

 $0.10 \pm 0.02^{+0.01}_{-0.02}$ ¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta\eta'$ P -wave.

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{347}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{348}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.54^{+0.34+3.82}_{-0.40-1.49}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{349}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.67 \pm 0.70^{+0.61}_{-1.67}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta'X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2470) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{350}/Γ

<u>VALUE</u> (units 10^{-7})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.18 \pm 1.77^{+3.73}_{-2.23}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta'X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{351}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7^{+0.6}_{-0.8}$ OUR AVERAGE				Error includes scale factor of 1.6.

 $3.93 \pm 0.38^{+0.31}_{-0.84}$ ¹ ABLIKIM 16J BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$ $2.2 \pm 0.4 \pm 0.4$ 264 ABLIKIM 05R BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

 $2.87 \pm 0.09^{+0.49}_{-0.52}$ 4265 ² ABLIKIM 11C BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatté formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma\eta(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

NODE=M070Q26

NODE=M070Q26

NODE=M070Q26;LINKAGE=A

NODE=M070Q36

NODE=M070Q36

NODE=M070Q36;LINKAGE=A

NODE=M070S88

NODE=M070S88

NODE=M070S88;LINKAGE=A

NODE=M070P70

NODE=M070P70

NODE=M070Q29

NODE=M070Q29

NODE=M070Q29;LINKAGE=A

NODE=M070Q27

NODE=M070Q27

NODE=M070Q27;LINKAGE=A

NODE=M070R78

NODE=M070R78

NODE=M070R78;LINKAGE=A

NODE=M070R78;LINKAGE=AI

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$	Γ_{352}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070S71</u>
0.77$^{+0.15}_{-0.09}$ OUR AVERAGE					<u>NODE=M070S71</u>
0.90 $^{+0.04}_{-0.11}$ $^{+0.27}_{-0.55}$		¹ ABLIKIM	12D	BES3 $J/\psi \rightarrow \gamma p\bar{p}$	
1.14 $^{+0.43}_{-0.30}$ $^{+0.42}_{-0.26}$	231	² ALEXANDER	10	CLEO $J/\psi \rightarrow \gamma p\bar{p}$	
0.70 ± 0.04 $^{+0.19}_{-0.08}$		BAI	03F	BES2 $J/\psi \rightarrow \gamma p\bar{p}$	
1 From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A.					
2 From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.					
$\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$	Γ_{353}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070S96</u>
3.31$^{+0.33}_{-0.30}$$^{+1.96}_{-1.29}$		ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$	<u>NODE=M070S96</u>
$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\phi(1020))/\Gamma_{\text{total}}$	Γ_{354}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070P37</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
1.77 ± 0.35 ± 0.25	305	¹ ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$	
8.09 ± 1.99 ± 1.36	1.3k	² ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$	
1 Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.					
2 Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.					
$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{355}/Γ				
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070P41</u>
<3.56 $\times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$	<u>NODE=M070P41</u>
$\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_{356}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070S82</u>
2.44$\pm 0.36$$^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$	<u>NODE=M070S82</u>
$\Gamma(\gamma\eta(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$	Γ_{357}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		<u>NODE=M070P86</u>
1.79$\pm 0.23$$\pm 0.65$	ABLIKIM	20Q	BES3	$J/\psi \rightarrow \gamma K^+ K^- \eta'$	<u>NODE=M070P86</u>
$\Gamma(\gamma\eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$	Γ_{358}/Γ				
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		<u>NODE=M070P87</u>
1.18$\pm 0.32$$\pm 0.39$	ABLIKIM	20Q	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$	<u>NODE=M070P87</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
1.31 ± 0.22 $^{+2.85}_{-0.84}$		¹ ABLIKIM	24	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$	
1 Decaying via the intermediate $f_0(980)\eta'$, fitted together with $X(1835)$, a 600 MeV broad structure around 2.8 GeV, and the tail of the $\eta_c(1S)$.					
$\Gamma(\gamma\eta(2370) \rightarrow \gamma\eta\eta\eta')/\Gamma_{\text{total}}$	Γ_{359}/Γ				
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070P88</u>
<9.2 $\times 10^{-6}$	90	ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$	<u>NODE=M070P88</u>
$\Gamma(\gamma D^0 + c.c.)/\Gamma_{\text{total}}$	Γ_{360}/Γ				
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>NODE=M070Q79</u>
<9.1 $\times 10^{-8}$	90	ABLIKIM	24BZ	BES3 $e^+ e^- \rightarrow J/\psi(1S)$	<u>NODE=M070Q79</u>
$\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$	Γ_{361}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.38$\pm 0.07$$\pm 0.07$		49	EATON	84	MRK2 $e^+ e^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.11		PERUZZI	78	MRK1	$e^+ e^-$

$\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{362}/Γ	NODE=M070R93
value $< 7.9 \times 10^{-3}$	CL% 90	DOCUMENT ID EATON TECN MPK2 e^+e^-

$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$	Γ_{363}/Γ	NODE=M070S8			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	NODE=M070S8
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.16 \times 10^{-3}$	90	BAI	98G	BES $e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\gamma A^0 \rightarrow \gamma \text{ invisible})/\Gamma_{\text{total}}$	Γ_{364}/Γ	NODE=M070S68				
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M070S68
$< 1.7 \times 10^{-6}$	90	88M	1_ABUKIM	20k	BES3	$\psi(2S) \rightarrow U/\pi^+\pi^-$

We do not use the following data for our mass fits, limits etc.

$<6.3 \times 10^{-6}$ 90 3.7M 2 INSLER 10 CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

²The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$ reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{365}/Γ	NODE=M070Q68			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	NODE=M070Q68
$< 4.9 \times 10^{-7}$	95	1 ABUKIM	24AD.BFS3	$1/\sqrt{s} \rightarrow \gamma\gamma\gamma$	

~~2.3.3 We do not use the following data for averages, fits, limits, etc.~~

We do not use the following data for averages, fits, limits, etc.

$<1.8 \times 10^{-3}$ 95 - ABLIKIM 23E BES3 $J/\psi \rightarrow \gamma\gamma\gamma$

¹ For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.18–2.85 GeV, The measured 95% CL limit as a function of m_{A^0} ranges from 3.7×10^{-8} to 4.85×10^{-7}

²For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.165–2.84 GeV. The measured 95% CL limit as a function of m_{A^0} ranges from 8.3×10^{-8} to 1.8×10^{-6} .

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{366}/Γ
(narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE CL% DOCUMENT ID TECN COMMENT

$<7.8 \times 10^{-7}$	90	¹ ABLIKIM	22H	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.5 \times 10^{-5}$	90	² ABLIKIM	16E	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$<2.1 \times 10^{-5}$	90	³ ABLIKIM	12	BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.
² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The

³For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

— DALITZ DECAYS —

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$	Γ_{367}/Γ	NODE=M070S89			
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M070S89
7.56±1.32±0.50	39	ABLIKIM	14I	BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$	Γ_{368}/Γ	NODE=M070S90 NODE=M070S90			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.12 ± 0.04 ± 0.07	2,471	1,2 ARIKIM	101	REC3	/ \ + -

1.42±0.04±0.07 2.47 \pm ABLIKIM 19A BESS $J/\psi \rightarrow$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.
² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14l.

$\Gamma(\eta'(958)e^+e^-)/\Gamma_{\text{total}}$	Γ_{369}/Γ
VALUE (units 10^{-5})	EVTS DOCUMENT ID TECN COMMENT

6.59±0.07±0.17 8.9k 1 ABLIKIM 19H BES3 $J/\psi \rightarrow \eta'(958)e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.81\pm0.16\pm0.31$ 1.4k 1,2 ABLIKIM 14I BES3 $J/\psi \rightarrow \eta'(958)e^+e^-$

1 Using both $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \pi^+\pi^-\eta$ decays.

2 Superseded by ABLIKIM 19H.

NODE=M070S91
NODE=M070S91

$\Gamma(\eta(1405)e^+e^- \rightarrow f_0(980)\pi^0e^+e^- \rightarrow \pi^+\pi^-\pi^0e^+e^-)/\Gamma_{\text{total}}$	Γ_{370}/Γ
VALUE (units 10^{-7})	EVTS DOCUMENT ID TECN COMMENT

2.04±0.20±0.08 203 1 ABLIKIM 24I BES3 $J/\psi \rightarrow e^+e^-\eta(1405)$

1 With a significance of 9.8 σ .

NODE=M070S91;LINKAGE=A
NODE=M070S91;LINKAGE=B

$\Gamma(X(1835)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$	Γ_{371}/Γ
VALUE (units 10^{-6})	EVTS DOCUMENT ID TECN COMMENT

3.58±0.19±0.16 1364 1 ABLIKIM 22B BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

1 Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

NODE=M070Q75
NODE=M070Q75
OCCUR=2
NODE=M070Q75;LINKAGE=B

$\Gamma(X(2120)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$	Γ_{372}/Γ
VALUE (units 10^{-6})	EVTS DOCUMENT ID TECN COMMENT

0.82±0.12±0.06 310 ABLIKIM 22B BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

NODE=M070Q22
NODE=M070Q22

$\Gamma(\eta(2370)e^+e^-, \eta \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$	Γ_{373}/Γ
VALUE (units 10^{-6})	EVTS DOCUMENT ID TECN COMMENT

1.08±0.14±0.10 397 ABLIKIM 22B BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

NODE=M070Q23
NODE=M070Q23

$\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$	Γ_{374}/Γ
VALUE	CL% DOCUMENT ID TECN COMMENT

<9.11 × 10⁻⁷ 90 1 ABLIKIM 19A BES3 $J/\psi \rightarrow \eta e^+e^-$

1 For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

NODE=M070P42
NODE=M070P42

$\Gamma(\eta'(958)U \rightarrow \eta'(958)e^+e^-)/\Gamma_{\text{total}}$	Γ_{375}/Γ
VALUE	CL% DOCUMENT ID TECN COMMENT

<2.0 × 10⁻⁷ 90 1 ABLIKIM 19H BES3 $J/\psi \rightarrow \eta'(958)e^+e^-$

1 For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

NODE=M070P61
NODE=M070P61

$\Gamma(\phi e^+e^-)/\Gamma_{\text{total}}$	Γ_{376}/Γ
VALUE (units 10^{-7})	CL% DOCUMENT ID TECN COMMENT

<1.2 90 1 ABLIKIM 19AB BES3 $J/\psi \rightarrow \phi e^+e^-$

1 Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) = (34.49 \pm 0.30)\%$.

NODE=M070P82
NODE=M070P82

WEAK DECAYS

$\Gamma(D^-e^+\nu_e + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{377}/Γ
VALUE	CL% DOCUMENT ID TECN COMMENT

<7.1 × 10⁻⁸ 90 ABLIKIM 21Q BES3 $e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-5}$ 90 ABLIKIM 06M BES2 $e^+e^- \rightarrow J/\psi$

NODE=M070P82;LINKAGE=A

NODE=M070320

$\Gamma(D^-\mu^+\nu_\mu + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{378}/Γ
VALUE	CL% DOCUMENT ID TECN COMMENT

<5.6 × 10⁻⁷ 90 1 ABLIKIM 24AM BES3 $e^+e^- \rightarrow J/\psi$

1 Using $B(D^- \rightarrow K^+\pi^-\pi^-) = 9.38 \pm 0.16\%$.

NODE=M070Q67
NODE=M070Q67

NODE=M070Q67;LINKAGE=A

$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{379}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.5 \times 10^{-8}$	90	1 ABLIKIM	17AF BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S54 NODE=M070S54
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<1.1 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$	NODE=M070S54;LINKAGE=A
1 Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$.					
$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{380}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S55 NODE=M070S55
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<3.6 \times 10^{-5}$	90	1 ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$	NODE=M070S55;LINKAGE=AB
1 Using $B(D_s^- \rightarrow \phi\pi^-) = 4.4 \pm 0.5\%$.					
$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{381}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070B13 NODE=M070B13
$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{382}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.0 \times 10^{-8}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S61 NODE=M070S61
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<7.5 \times 10^{-5}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$	
$\Gamma(D^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{383}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.0 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070Q63 NODE=M070Q63
$\Gamma(\bar{D}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{384}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.7 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070Q64 NODE=M070Q64
$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{385}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-4}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$	NODE=M070S62 NODE=M070S62
$\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{386}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S93 NODE=M070S93
$\Gamma(\bar{D}^0 \eta + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{387}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.8 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070Q65 NODE=M070Q65
$\Gamma(\bar{D}^0 \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{388}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.2 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070Q66 NODE=M070Q66
$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{389}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-4}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$	NODE=M070S63 NODE=M070S63
$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{390}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S92 NODE=M070S92
— CHARGE CONJUGATION (C), PARITY (P), — — LEPTON FAMILY NUMBER (LF) VIOLATING MODES —					
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				Γ_{391}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.7 \times 10^{-7}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	NODE=M070R80 NODE=M070R80

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.5 \times 10^{-5}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$< 1.6 \times 10^{-4}$	90	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$< 50 \times 10^{-5}$	90	BARTEL	77	CNTR	$e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = 1.020 \times 10^{-3}$.

NODE=M070R80;LINKAGE=W1

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$		Γ_{392}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 1.4 \times 10^{-6}$	90	ABLIKIM	14Q	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070S95
NODE=M070S95

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$		Γ_{393}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 1.6 \times 10^{-7}$	90	ABLIKIM	13L	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S39
NODE=M070S39

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$		Γ_{394}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 7.5 \times 10^{-8}$	90	ABLIKIM	21M	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S40
NODE=M070S40

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$		Γ_{395}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.0 \times 10^{-6}$	90	ABLIKIM	04	BES	$e^+ e^- \rightarrow J/\psi$

NODE=M070S40;LINKAGE=A

$\Gamma(A_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$		Γ_{396}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 6.9 \times 10^{-8}$	90	ABLIKIM	19AF	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow pK^-\pi^+e^- (+ \text{c.c.})$

NODE=M070P74
NODE=M070P74

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$		Γ_{397}/Γ_5			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 6.6 \times 10^{-2}$	90	LEES	13I	BABR	$B \rightarrow K^{(*)} J/\psi$

NODE=M070S80
NODE=M070S80

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$		Γ_{397}/Γ_7			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070S60
NODE=M070S60

$\Gamma(\mu^+ \mu^- X^0 \rightarrow \mu^+ \mu^- + \text{invisible})/\Gamma_{\text{total}}$		Γ_{398}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 5.5 \times 10^{-7}$	90	¹ ABLIKIM	24G	BES3	$J/\psi \rightarrow \mu^+ \mu^- X^0 (\text{scalar}) \rightarrow \mu^+ \mu^- + \text{invisible}$
$< 9.6 \times 10^{-7}$	90	² ABLIKIM	24G	BES3	$J/\psi \rightarrow \mu^+ \mu^- X^0 (\text{vector}) \rightarrow \mu^+ \mu^- + \text{invisible}$

NODE=M070Q69
NODE=M070Q69

OCCUR=2

¹ For a light scalar, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 6.2×10^{-9} to 5.5×10^{-7} .

NODE=M070Q69;LINKAGE=A

² For a light vector, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 4.5×10^{-9} to 9.6×10^{-7} .

NODE=M070Q69;LINKAGE=B

J/ ψ (1S) REFERENCES

NODE=M070

AAIJ	24AE	JHEP 2412 062	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=63039
ABLIKIM	24	PRL 132 181901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62645
ABLIKIM	24AB	PR D110 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62681
ABLIKIM	24AD	PR D110 L031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62683
ABLIKIM	24AM	JHEP 2401 126	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62697
ABLIKIM	24BI	PR D110 032020	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62938
ABLIKIM	24BQ	PR D110 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=63020
ABLIKIM	24BZ	PR D110 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=63031
ABLIKIM	24CB	PR D110 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=63033
ABLIKIM	24G	PR D109 L031102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62657
ABLIKIM	24H	PR D109 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62658
ABLIKIM	24I	PR D109 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62659
ABLIKIM	24L	PR D109 052006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62662
HAYRAPETY...	24A	PR D109 L111101	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62675
ABLIKIM	23BD	PR D108 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62425
ABLIKIM	23BU	PR D108 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62517
ABLIKIM	23E	PL B838 137698	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62047
ABLIKIM	23S	PR D107 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62063
ANASHIN	23	JHEP 2306 196	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=62057
GONG	23	PR D107 072008	G. Gong <i>et al.</i>	(BELLE Collab.)	REFID=62073
LEES	23	PR D107 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=62061
LIAO	23	PR D107 112007	L. Liao <i>et al.</i>		REFID=62279
ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ABLIKIM	22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61881
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61902
ABLIKIM	22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61636
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61643
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=61894
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61445
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61463
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61030
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61118
ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61125
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61113
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61451
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60048
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60316
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60457
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=60512
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59517
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61449
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59846
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59850
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59857
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59890
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59604
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59747
ABLIKIM	19T	PR D122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59456
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58849
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59102
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58317
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57990
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57265
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57454
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57509
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM	16N	PR D93 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57512
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57522
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57566
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
AAIJ	15B1	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56776
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ABLIKIM	15T	PR D115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56785
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55900
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55902
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55905
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56238
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56388
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=55655

LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54954
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55300
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55392
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54265
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54267
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54268
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54863
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53220
DEL-AMO-SA..	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53533
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)	REFID=53359
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52718
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53099
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52047
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52128
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52253
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52255
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52256
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52571
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=52685
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52072
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 11900E (errat.)	(B. Aubert <i>et al.</i>)	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50986
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51058
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51128
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51130
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51036
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT	06B	PR D73 020005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50496
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50507
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)	REFID=50802
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	(BES Collab.)	REFID=51038
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49739
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49611
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49607
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49751
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49754
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
SETH	04	PR D69 097503	K.K. Seth	(KEDR Collab.)	REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(BES Collab.)	REFID=49579
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49403

BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47427
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46341
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)	REFID=46608
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45146
BAI	96B	PR 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44736
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45198
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)	REFID=45197
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44434
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)	REFID=43314
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)	REFID=43601
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41866
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)	REFID=41899
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41359
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)	REFID=41349
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=40346
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)	REFID=40002
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
BISELLLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40261
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)	REFID=40243
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)	REFID=21865
BISELLLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=22101
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22097
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)	REFID=22092
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22080
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22077
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22073
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
Translated from YAF 34 1471.					
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22065
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22066
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22067
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)	REFID=22068
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22058
BURMEISTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22060
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)	REFID=22063
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22054
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG	REFID=22056
BALDINI...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)	REFID=22026
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC	REFID=22030
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22036
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=22038
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)	REFID=22039