NODE=M079

$\chi_{b0}(2P)$

$I^G(J^{PC}) = 0^+(0^{++})$
J needs confirmation.

NODE=M079

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b0}(2P)$ MASS

NODE=M079M

NODE=M079M
→ UNCHECKED ←

VALUE (MeV)	*DOCUMENT ID*
10232.5±0.4±0.5 OUR EVALUATION	From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

NODE=M079M2
NODE=M079M2

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
23.8±1.7	LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$

γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M079DM

NODE=M079DM
→ UNCHECKED ←

VALUE (MeV)	*EVTS*	*DOCUMENT ID*		*TECN*	*COMMENT*
121.9 ±0.4 OUR EVALUATION		Treating systematic errors as correlated			
122.2 ±0.5 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.			
121.55±0.16±0.46		ARTUSO	05	CLEO	$\Upsilon(3S) \to \gamma X$
123.0 ±0.8	4959	[1] HEINTZ	92	CSB2	$e^+e^- \to \gamma X$
124.6 ±1.4	17	[2] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
122.3 ±0.3 ±0.6	9903	MORRISON	91	CLE2	$e^+e^- \to \gamma X$

OCCUR=2

[1] A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M079DM;LINKAGE=A

[2] A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M079DM;LINKAGE=B



$\chi_{b0}(2P)$ DECAY MODES

NODE=M079215;NODE=M079

	Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1	$\gamma\,\Upsilon(2S)$	(1.38 ± 0.30) %		DESIG=2
Γ_2	$\omega\,\Upsilon(1S)$	$(5.5 \pm2.0) \times 10^{-3}$		DESIG=18
Γ_3	$\gamma\,\Upsilon(1S)$	$(3.8 \pm1.7) \times 10^{-3}$		DESIG=1
Γ_4	$D^0 X$	< 8.2 %	90%	DESIG=3
Γ_5	$\pi^+\pi^-K^+K^-\pi^0$	$< 3.4 \times 10^{-5}$	90%	DESIG=4
Γ_6	$2\pi^+\pi^-K^-K^0_S$	$< 5 \times 10^{-5}$	90%	DESIG=5
Γ_7	$2\pi^+\pi^-K^-K^0_S 2\pi^0$	$< 2.2 \times 10^{-4}$	90%	DESIG=6
Γ_8	$2\pi^+2\pi^-2\pi^0$	$< 2.4 \times 10^{-4}$	90%	DESIG=7

Γ_9	$2\pi^+ 2\pi^- K^+ K^-$	< 1.5	$\times 10^{-4}$	90%	DESIG=8
Γ_{10}	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.2	$\times 10^{-4}$	90%	DESIG=9
Γ_{11}	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 1.1	$\times 10^{-3}$	90%	DESIG=10
Γ_{12}	$3\pi^+ 2\pi^- K^- K^0_S \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=11
Γ_{13}	$3\pi^+ 3\pi^-$	< 7	$\times 10^{-5}$	90%	DESIG=12
Γ_{14}	$3\pi^+ 3\pi^- 2\pi^0$	< 1.2	$\times 10^{-3}$	90%	DESIG=13
Γ_{15}	$3\pi^+ 3\pi^- K^+ K^-$	< 1.5	$\times 10^{-4}$	90%	DESIG=14
Γ_{16}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=15
Γ_{17}	$4\pi^+ 4\pi^-$	< 1.7	$\times 10^{-4}$	90%	DESIG=16
Γ_{18}	$4\pi^+ 4\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=17

$\chi_{b0}(2P)$ BRANCHING RATIOS

NODE=M079220

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ Γ_1/Γ

NODE=M079R2
NODE=M079R2

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
1.38±0.30 OUR AVERAGE					
$1.31\pm0.27^{+0.13}_{-0.12}$		[3,4] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$
$3.6 \pm1.6 \pm0.3$		[3,5] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.8	90	[6] LEES	11J	BABR	$\Upsilon(3S) \to X\gamma$
<8.9	90	[7] CRAWFORD	92B	CLE2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$

[3] Assuming B($\Upsilon(2S) \to \mu^+\mu^-$) = (1.93 ± 0.17)%. NODE=M079R2;LINKAGE=D

[4] LEES 14M reports $[\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] = (7.7 \pm 1.6) \times 10^{-4}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value. NODE=M079R2;LINKAGE=E

[5] Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\chi_{b0}(2P) \to \gamma\Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$ using $B(\Upsilon(2S) \to \mu^+\mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91. NODE=M079R2;LINKAGE=C

[6] LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$. NODE=M079R2;LINKAGE=LE

[7] Using $B(\Upsilon(2S) \to \mu^+\mu^-) = (1.37\pm0.26)\%$, $B(\Upsilon(3S) \to \gamma\gamma\Upsilon(2S))\times 2\, B(\Upsilon(2S) \to \mu^+\mu^-) < 1.19 \times 10^{-4}$, and $B(\Upsilon(3S) \to \chi_{b0}(2P)\gamma) = 0.049$. NODE=M079R2;LINKAGE=B

$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_2/Γ

NODE=M079R00
NODE=M079R00

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
0.55±0.19±0.07	32	[8] STOTTLER	25	BELL	$\Upsilon(3S) \to \gamma\omega\Upsilon(1S)$

[8] STOTTLER 25 reports 0.55 ± 0.19 ± 0.07 % from a measurement of $[\Gamma(\chi_{b0}(2P) \to \omega\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))]$ assuming $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. NODE=M079R00;LINKAGE=A

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

NODE=M079R1
NODE=M079R1

VALUE (%)	CL%	DOCUMENT ID		TECN	COMMENT
0.38±0.17 OUR AVERAGE					
$0.36\pm0.17\pm0.03$		[9,10,11] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$
$0.9 \pm0.7 \pm0.1$		[10,12] HEINTZ	92	CSB2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.2	90	[13] LEES	11J	BABR	$\Upsilon(3S) \to X\gamma$
<2.5	90	[14] CRAWFORD	92B	CLE2	$e^+e^- \to \ell^+\ell^-\gamma\gamma$

[9] LEES 14M quotes $\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{\text{total}} = (2.1 \pm 1.0) \times 10^{-4}$ combining the results from $\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$ samples with and without photon conversions. NODE=M079R1;LINKAGE=D

[10] Assuming $B(\Upsilon(1S) \to \mu^+\mu^-) = (2.48 \pm 0.05)\%$. NODE=M079R1;LINKAGE=E

[11] LEES 14M reports $[\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] = (2.1 \pm 1.0) \times 10^{-4}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value. NODE=M079R1;LINKAGE=F

[12] Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\chi_{b0}(2P) \to \gamma\Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \to \mu^+\mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91. NODE=M079R1;LINKAGE=C

[13] LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$. NODE=M079R1;LINKAGE=LE

[14] Using $B(\Upsilon(1S) \to \mu^+\mu^-) = (2.57\pm0.07)\%$, $B(\Upsilon(3S) \to \gamma\gamma\Upsilon(1S))\times 2\, B(\Upsilon(1S) \to \mu^+\mu^-) < 0.63 \times 10^{-4}$, and $B(\Upsilon(3S) \to \chi_{b0}(2P)\gamma) = 0.049$. NODE=M079R1;LINKAGE=B

$\Gamma(D^0 X)/\Gamma_{total}$ Γ_4/Γ

NODE=M079R01

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<8.2 × 10^{-2}	90	[15,16] BRIERE	08	CLEO	$\Upsilon(3S) \to \gamma D^0 X$

[15] For $p_{D^0} > 2.5$ GeV/c.

[16] The authors also present their result as $(4.1 \pm 3.0 \pm 0.4) \times 10^{-2}$.

NODE=M079R01;LINKAGE=BR
NODE=M079R01;LINKAGE=RI

$\Gamma(\pi^+\pi^- K^+ K^- \pi^0)/\Gamma_{total}$ Γ_5/Γ

NODE=M079R02

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.34	90	[17] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma\pi^+\pi^- K^+ K^- \pi^0$

[17] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to \pi^+\pi^- K^+ K^- \pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 2\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R02;LINKAGE=AS

$\Gamma(2\pi^+\pi^- K^- K^0_S)/\Gamma_{total}$ Γ_6/Γ

NODE=M079R03

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.5	90	[18] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+\pi^- K^- K^0_S$

[18] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+\pi^- K^- K^0_S)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 3\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R03;LINKAGE=AS

$\Gamma(2\pi^+\pi^- K^- K^0_S 2\pi^0)/\Gamma_{total}$ Γ_7/Γ

NODE=M079R04

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.2	90	[19] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+\pi^- K^- 2\pi^0$

[19] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+\pi^- K^- K^0_S 2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R04;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{total}$ Γ_8/Γ

NODE=M079R05

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.4	90	[20] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+ 2\pi^- 2\pi^0$

[20] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 14 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R05;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{total}$ Γ_9/Γ

NODE=M079R06

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.5	90	[21] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+ 2\pi^- K^+ K^-$

[21] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 9\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R06;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{total}$ Γ_{10}/Γ

NODE=M079R07

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<2.2	90	[22] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

[22] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R07;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{total}$ Γ_{11}/Γ

NODE=M079R08

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<11	90	[23] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

[23] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 63 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R08;LINKAGE=AS

$\Gamma(3\pi^+ 2\pi^- K^- K^0_S \pi^0)/\Gamma_{total}$ Γ_{12}/Γ

NODE=M079R09

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<7	90	[24] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 3\pi^+ 2\pi^- K^- K^0_S \pi^0$

[24] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 3\pi^+ 2\pi^- K^- K^0_S \pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 39 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R09;LINKAGE=AS

$\Gamma(3\pi^+3\pi^-)/\Gamma_{total}$ — Γ_{13}/Γ

NODE=M079R10

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<0.7	90	[25] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 3\pi^+ 3\pi^-$

[25] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 3\pi^+3\pi^-)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 4\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R10;LINKAGE=AS

$\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{total}$ — Γ_{14}/Γ

NODE=M079R11

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<12	90	[26] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 3\pi^+ 3\pi^- 2\pi^0$

[26] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 3\pi^+3\pi^-2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 72 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R11;LINKAGE=AS

$\Gamma(3\pi^+3\pi^-K^+K^-)/\Gamma_{total}$ — Γ_{15}/Γ

NODE=M079R12

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.5	90	[27] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 3\pi^+ 3\pi^- K^+ K^-$

[27] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 3\pi^+3\pi^-K^+K^-)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 9\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R12;LINKAGE=AS

$\Gamma(3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{total}$ — Γ_{16}/Γ

NODE=M079R13

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<7	90	[28] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

[28] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 43 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R13;LINKAGE=AS

$\Gamma(4\pi^+4\pi^-)/\Gamma_{total}$ — Γ_{17}/Γ

NODE=M079R14

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.7	90	[29] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 4\pi^+ 4\pi^-$

[29] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 4\pi^+4\pi^-)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 10\times10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R14;LINKAGE=AS

$\Gamma(4\pi^+4\pi^-2\pi^0)/\Gamma_{total}$ — Γ_{18}/Γ

NODE=M079R15

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<6	90	[30] ASNER	08A	CLEO	$\Upsilon(3S) \to \gamma 4\pi^+ 4\pi^- 2\pi^0$

[30] ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \to 4\pi^+4\pi^-2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))] < 38 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R15;LINKAGE=AS

$\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{total}$ — $\Gamma_3/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma^{\Upsilon(3S)}$

NODE=M079B01

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<8.2	90	[31] LEES	11J	BABR	$\Upsilon(3S) \to X\gamma$

[31] LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{total} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \to \gamma\Upsilon(1S)) < 1.2\%$ using $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.

NODE=M079B01;LINKAGE=LE

$B(\chi_{b0}(2P) \to \gamma\Upsilon(1S)) \times B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\Upsilon(1S) \to \ell^+\ell^-)$

NODE=M079A02

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
1.4±0.9 OUR AVERAGE				
$1.7^{+1.5+0.1}_{-1.4-1.2}$	[32] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$
$1.3\pm1.0\pm0.3$	[33] HEINTZ	92	CSB2	$\Upsilon(3S) \to \gamma\gamma\ell^+\ell^-$

[32] From a sample of $\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$ with one converted photon.

NODE=M079A02;LINKAGE=A

[33] Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\chi_{b0}(2P) \to \gamma\Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \to \mu^+\mu^-) = (2.57 \pm 0.05)\%$.

NODE=M079A02;LINKAGE=K

$[B(\chi_{b0}(2P) \to \gamma\Upsilon(1S)) \times B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))]$ / $[B(\chi_{b1}(2P) \to \gamma\Upsilon(1S)) \times B(\Upsilon(3S) \to \gamma\chi_{b1}(2P))]$

NODE=M079A00
NODE=M079A00

VALUE (%)	DOCUMENT ID		TECN	COMMENT
1.71±0.80	[34] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$

[34] From a sample of $\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$ without converted photons.

NODE=M079A00;LINKAGE=A

$\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(2S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{total}$ $\Gamma_1/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma^{\Upsilon(3S)}$

NODE=M079B02
NODE=M079B02

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<1.6	90	[35] LEES	11J	BABR	$\Upsilon(3S) \to X\gamma$

[35] LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \to \gamma\Upsilon(2S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \to \gamma\chi_{b0}(2P))/\Gamma_{total} = (-0.3 \pm 0.2^{+0.5}_{-0.4})$% and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \to \gamma\Upsilon(2S)) < 2.8$% using $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)$%.

NODE=M079B02;LINKAGE=LE

$B(\chi_{b0}(2P) \to \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\Upsilon(2S) \to \ell^+\ell^-)$

NODE=M079A03
NODE=M079A03

VALUE (units 10^{-5})	DOCUMENT ID		TECN	COMMENT
4.4±1.6 OUR AVERAGE				
$6.6^{+4.9}_{-4.0}{}^{+2.0}_{-0.3}$	[36] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$
4.0±1.7±0.3	[37] HEINTZ	92	CSB2	$\Upsilon(3S) \to \gamma\gamma\ell^+\ell^-$

[36] From a sample of $\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$ with one converted photon.

NODE=M079A03;LINKAGE=A

[37] Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \to \gamma\chi_{b0}(2P)) \times B(\chi_{b0}(2P) \to \gamma\Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)$% using $B(\Upsilon(2S) \to \mu^+\mu^-) = (1.44 \pm 0.10)$%.

NODE=M079A03;LINKAGE=B

$[B(\chi_{b0}(2P) \to \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \to \gamma\chi_{b0}(2P))]$ / $[B(\chi_{b1}(2P) \to \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \to \gamma\chi_{b1}(2P))]$

NODE=M079A01
NODE=M079A01

VALUE (%)	DOCUMENT ID		TECN	COMMENT
3.31±0.56	[38] LEES	14M	BABR	$\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$

[38] From a sample of $\Upsilon(3S) \to \gamma\gamma\mu^+\mu^-$ without converted photons.

NODE=M079A01;LINKAGE=A

$\chi_{b0}(2P)$ REFERENCES

NODE=M079

STOTTLER	25	PRL 135 121902	Z.S. Stottler *et al.*	(BELLE Collab.)	REFID=63606
LEES	14M	PR D90 112010	J.P. Lees *et al.*	(BABAR Collab.)	REFID=56343
LEES	11J	PR D84 072002	J.P. Lees *et al.*	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner *et al.*	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere *et al.*	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso *et al.*	(CLEO Collab.)	REFID=50454
CRAWFORD	92B	PL B294 139	G. Crawford *et al.*	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz *et al.*	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz *et al.*	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison *et al.*	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain *et al.*	(CUSB Collab.)	REFID=41586