

NODE=M176

 $\chi_{c1}(3872)$

$I^G(J^{PC}) = 0^+(1^{++})$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M176

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to < 4% at 95% CL.

See the review on "Spectroscopy of Mesons Containing Two Heavy Quarks."

 $\chi_{c1}(3872)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \operatorname{Im}(\sqrt{s})$.

NODE=M176PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$(3871.70 \pm 0.15 \pm 0.07)$	¹ ABLIKIM	24C	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
$- i(0.19 \pm 0.08 \pm 0.14)$			

NODE=M176PP

¹ From simultaneous line shape fits of $e^+e^- \rightarrow \gamma\chi_{c1}(3872) \rightarrow \gamma[D^0\bar{D}^0\pi^0]$ and $\gamma[J/\psi\pi^+\pi^-]$. The most prominent pole is reported at $7.04 \pm 0.15 \pm 0.07$ MeV above the $D^0\bar{D}^0\pi^0$ threshold of 3864.66 MeV on the first sheet with respect to the $D^{*0}\bar{D}^0$ channel. The uncertainty in the D^0 width is included in the uncertainty of the pole mass.

NODE=M176PP;LINKAGE=A

 $\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

NODE=M176M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.64 ± 0.06 OUR AVERAGE				
3870.2 ± 0.7 ± 0.3	24.6	ABLIKIM	23W BES3	$e^+e^- \rightarrow J/\psi(1S)\pi^+\pi^-\omega$
3871.64 ± 0.06 ± 0.01	19.8k	¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
3871.9 ± 0.7 ± 0.2	20	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95 ± 0.48 ± 0.12	0.6k	AAIJ	12H LHCb	$pp \rightarrow J/\psi\pi^+\pi^-X$
3871.85 ± 0.27 ± 0.19	170	² CHOI	11 BELL	$B \rightarrow K\pi^+\pi^-J/\psi$
3873 ± 1.8 ± 1.3	27	³ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61 ± 0.16 ± 0.19	6k	^{3,4} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3871.4 ± 0.6 ± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+J/\psi\pi^+\pi^-$
3868.7 ± 1.5 ± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.8 ± 3.1 ± 3.0	522	^{3,5} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3871.57 ± 0.09	155	⁶ AAIJ	23AP LHCb	$B_s^0 \rightarrow J/\psi 2(\pi^+\pi^-)$
3871.695 $\pm 0.067 \pm 0.068$	15.6k	⁷ AAIJ	20AD LHCb	$pp \rightarrow J/\psi\pi^+\pi^-X$
3871.59 ± 0.06 ± 0.03	4.2k	⁸ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
3873.3 ± 1.1 ± 1.0	45	⁹ ABLIKIM	19V BES	$e^+e^- \rightarrow \gamma\omega J/\psi$
3860.0 ± 10.4	13.6	^{3,10} AGHASYAN	18A COMP	$\gamma^*N \rightarrow X\pi^\pm N'$
3868.6 ± 1.2 ± 0.2	8	¹¹ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$

NODE=M176M

OCCUR=2

OCCUR=2

3871.3	± 0.6	± 0.1	61	11 AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi \pi^+ \pi^-$
3873.4	± 1.4		25	12 AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
3871.3	± 0.7	± 0.4	730	3,13 ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3872.0	± 0.6	± 0.5	36	14 CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
3836	± 13		58	3,15 ANTONIAZZI	94 E705	$300 \pi^\pm Li \rightarrow J/\psi \pi^\pm \pi^- X$

OCCUR=2

¹ Calculated from $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.54 \pm 0.06$ MeV obtained by combining the data with $\chi_{c1}(3872)$ produced in B^+ decays from AAIJ 20S and inclusive b -hadron decays from AAIJ 20AD and using $m_{\psi(2S)} = 3686.097$ MeV. Breit-Wigner parametrization.

² The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

³ Width consistent with detector resolution.

⁴ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/ c^2 is excluded at 95% CL.

⁵ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.

⁶ From a fit of a relativistic S-wave Breit-Wigner convolved with the detector resolution. The width of $\chi_{c1}(3872)$ is constrained to the PDG 22 value. Systematic errors not evaluated.

⁷ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays and $m_{\psi(2S)} = 3686.097 \pm 0.010$ MeV. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.

⁸ Using Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.

⁹ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

¹⁰ Could be a different state.

¹¹ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.

¹² Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.

¹³ Superseded by AALTONEN 09AU.

¹⁴ Superseded by CHOI 11.

¹⁵ A lower mass value can be due to an incorrect momentum scale for soft pions.

$\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3873.71 $^{+0.56}_{-0.50} \pm 0.13$		1 HIRATA	23 BELL	$B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
3872.9 $^{+0.6}_{-0.4} \pm 0.4$	50	2,3 AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
3875.1 $^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	3 AUBERT	08B BABR	$B \rightarrow \bar{D}^{*0} D^0 K$
3875.2 ± 0.7 $^{+0.9}_{-1.8}$	24 ± 6	3,4 GOKHROO	06 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

NODE=M176M;LINKAGE=D

NODE=M176M;LINKAGE=E

NODE=M176M;LINKAGE=B

NODE=M176M;LINKAGE=A

NODE=M176M;LINKAGE=AE

NODE=M176M;LINKAGE=AU

NODE=M176M;LINKAGE=AT

NODE=M176M;LINKAGE=CH

NODE=M176M;LINKAGE=AN

NODE=M176MD0

NODE=M176MD0

NODE=M176MD0;LINKAGE=A

NODE=M176MD0;LINKAGE=AS

NODE=M176MD0;LINKAGE=AU

NODE=M176MD0;LINKAGE=GO

NODE=M176DM

NODE=M176DM

NODE=M176DM2

NODE=M176DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
774.9 $\pm 3.1 \pm 3.0$	522	ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

$m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
185.598 $\pm 0.067 \pm 0.068$	15.6k	1 AAIJ	20AD LHCb	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
185.54 ± 0.06	19.8k	2 AAIJ	20S LHCb	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
187.4 ± 1.4	25	3 AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.

² Combining $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.49 \pm 0.06 \pm 0.03$ MeV from AAIJ 20S and the measured mass difference from AAIJ 20AD. Breit-Wigner parametrization.

³ Superseded by AUBERT 06.

$\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.19±0.21 OUR AVERAGE			Error includes scale factor of 1.1.		
1.39±0.24±0.10	15.6k	1 AAIJ	20AD LHCb	$p p \rightarrow J/\psi \pi^+ \pi^- X$	
0.96 ^{+0.19} _{-0.18} ±0.21	4.2k	2 AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4	90	ABLIKIM	14	BES3	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<1.2	90	CHOI	11	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
<3.3	90	AUBERT	08Y	BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
<4.1	90	69	AUBERT	06	BABR $B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36	3 CHOI	03	BELL $B \rightarrow K \pi^+ \pi^- J/\psi$

¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization.

² Using Breit-Wigner parametrization. Partially overlapping dataset with that of AAIJ 20AD.

3 Superseded by CHOI 11.

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ^{+2.2} _{-1.5} ±0.4		1 HIRATA	23	BELL $B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
3.9 ^{+2.8} _{-1.4} ^{+0.2} _{-1.1}	50	2 AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
3.0 ^{+1.9} _{-1.4} ±0.9	33 ± 6	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$

¹ From a fit of a Breit-Wigner function with energy dependent width.

² With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

$\chi_{c1}(3872)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	< 2.7 × 10 ⁻⁷	90%
$\Gamma_2 \pi^+ \pi^- \pi^0$	< 1.0 %	90%
$\Gamma_3 \pi^+ \pi^- J/\psi(1S)$	(4.3 ± 1.4) %	
$\Gamma_4 \pi^+ \pi^- \pi^0 J/\psi(1S)$	not seen	
$\Gamma_5 \omega \eta_c(1S)$	< 40 %	90%
$\Gamma_6 \rho(770)^0 J/\psi(1S)$	(3.4 ± 1.1) %	
$\Gamma_7 \omega J/\psi(1S)$	(5.0 ± 1.9) %	
$\Gamma_8 \phi \phi$	not seen	
$\Gamma_9 D^0 \bar{D}^0 \pi^0$	(55 ± 28) %	
$\Gamma_{10} \bar{D}^{*0} D^0$	(46 ± 16) %	
$\Gamma_{11} \gamma \gamma$	< 13 %	90%
$\Gamma_{12} D^0 \bar{D}^0$	< 32 %	90%
$\Gamma_{13} D^+ D^-$	< 22 %	90%
$\Gamma_{14} \pi^0 \chi_{c2}$	< 5 %	90%
$\Gamma_{15} \pi^0 \chi_{c1}$	(3.8 ± 1.9) %	
$\Gamma_{16} \pi^0 \chi_{c0}$	< 16 %	90%
$\Gamma_{17} \pi^+ \pi^- \eta_c(1S)$	< 16 %	90%
$\Gamma_{18} \pi^0 \pi^0 \chi_{c0}$	< 7 %	90%
$\Gamma_{19} \pi^0 \pi^0 \chi_{c1}$	< 5 %	90%
$\Gamma_{20} \pi^0 \pi^0 \chi_{c2}$	< 2.2 %	90%
$\Gamma_{21} \pi^+ \pi^- \chi_{c0}$	< 2.4 %	90%
$\Gamma_{22} \pi^+ \pi^- \chi_{c1}$	< 8 × 10 ⁻³	90%
$\Gamma_{23} p \bar{p}$	< 2.7 × 10 ⁻⁵	95%
$\Gamma_{24} \pi^+ \pi^- \eta$	< 5 × 10 ⁻³	90%

NODE=M176DM2;LINKAGE=A

NODE=M176DM2;LINKAGE=E

NODE=M176DM2;LINKAGE=AU

NODE=M176W

NODE=M176W

OCCUR=3

OCCUR=2

NODE=M176W;LINKAGE=E

NODE=M176W;LINKAGE=F

NODE=M176W;LINKAGE=CH

NODE=M176WD0

NODE=M176WD0

NODE=M176WD0;LINKAGE=B

NODE=M176WD0;LINKAGE=AU

NODE=M176215;NODE=M176

DESIG=1

DESIG=29

DESIG=2

DESIG=25

DESIG=24

DESIG=32

DESIG=13

DESIG=26

DESIG=8

DESIG=12

DESIG=5

DESIG=6

DESIG=7

DESIG=20

DESIG=18

DESIG=19

DESIG=14

DESIG=28

DESIG=36

DESIG=37

DESIG=27

DESIG=17

DESIG=16

DESIG=34

Radiative decays

Γ_{25}	$\gamma D^+ D^-$	< 4	%	90%
Γ_{26}	$\gamma \bar{D}^0 D^0$	< 7	%	90%
Γ_{27}	$\gamma J/\psi$	(10 \pm 4) $\times 10^{-3}$		
Γ_{28}	$\gamma \chi_{c1}$	< 1.0	%	90%
Γ_{29}	$\gamma \chi_{c2}$	< 4	%	90%
Γ_{30}	$\gamma \psi(2S)$	possibly seen		
Γ_{31}	$\gamma \psi_2(3823)$	< 3.3	$\times 10^{-3}$	90%

C-violating decays

Γ_{32}	$\eta J/\psi$	< 2.1	%	90%
---------------	---------------	-------	---	-----

 $\chi_{c1}(3872)$ PARTIAL WIDTHS **$\Gamma(e^+ e^-)$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_1
< 0.32	90	1 ABLIKIM	230 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 4.3	90	2 ABLIKIM	15V BES3	4.0–4.4 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
< 280	90	3 YUAN	04 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
1 Fit to cross section using a total width value of 1.19 ± 0.21 MeV and $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = (3.8 \pm 1.2)\%$ from PDG 20.					
2 ABLIKIM 15V reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-)/\Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma = 3\%$.					
3 Using BAI 98E data on $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$. Assuming that $\Gamma(\pi^+ \pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).					

 $\chi_{c1}(3872) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$ **$\Gamma(\pi^+ \pi^- J/\psi(1S)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$** **$\Gamma_3 \Gamma_1/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3 \Gamma_1/\Gamma$
< 7.5 $\times 10^{-3}$	90	1 ABLIKIM	230 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.13	90	ABLIKIM	15V BES3	4.0–4.4 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
< 6.2	90	2,3 AUBERT	05D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$	
< 8.3	90	3 DOBBS	05 CLE3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
< 10	90	4 YUAN	04 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
1 Fit to cross section using a total width value of 1.19 ± 0.21 MeV from PDG 20.					
2 Using $B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-) \cdot B(J/\psi \rightarrow \mu^+ \mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.					
3 Assuming $\chi_{c1}(3872)$ has $JPC = 1^{--}$.					
4 Using BAI 98E data on $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.88 \pm 0.10)\%$.					

 $\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(\pi^+ \pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$** **$\Gamma_3 \Gamma_{11}/\Gamma$**

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_3 \Gamma_{11}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
5.5 ± 4.1 ± 0.7	3	1 TERAMOTO	21 BELL	$e^+ e^- \rightarrow \gamma^* \gamma$ at $\Upsilon(nS)$		
< 12.9	90	2 DOBBS	05 CLE3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi \gamma$		
1 Measured in single-tag two-photon production assuming Q^2 dependence of a $c\bar{c}$ meson model. Here, $\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)$ is the reduced two-photon decay width, $\tilde{\Gamma}_{\gamma\gamma}$.						
2 Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.						

 $\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_7 \Gamma_{11}/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_7 \Gamma_{11}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.7	90	1 LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$	
1 Assuming $\chi_{c1}(3872)$ has spin 2.					

NODE=M176;CLUMP=B

DESIG=21

DESIG=23

DESIG=9

DESIG=3

DESIG=15

DESIG=11

DESIG=35

NODE=M176;CLUMP=A

DESIG=4

NODE=M176220

NODE=M176W1

NODE=M176W1

NODE=M176W1;LINKAGE=C

NODE=M176W1;LINKAGE=B

NODE=M176W1;LINKAGE=A

NODE=M176230

NODE=M176G1

NODE=M176G1

NODE=M176G1;LINKAGE=B

NODE=M176G1;LINKAGE=AU

NODE=M176G1;LINKAGE=DO

NODE=M176G1;LINKAGE=A

NODE=M176232

NODE=M176H1

NODE=M176H1

NODE=M176H1;LINKAGE=A

NODE=M176H1;LINKAGE=DO

NODE=M176G01

NODE=M176G01

NODE=M176G01;LINKAGE=LE

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{17}\Gamma_{11}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<11.1	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

$\chi_{c1}(3872)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	1,2 YIN	23 BELL	$B^+ \rightarrow \chi_{c1}(3872)K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3 90 2,3 YIN 23 BELL $B^0 \rightarrow \chi_{c1}(3872)K^0$

1 YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

2 Assuming the decay products, $\pi^+\pi^-\pi^0$, are uniformly distributed in phase space. The limit is the 90% "credible" upper limit (i.e. Bayesian).

3 YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \chi_{c1}(3872)K^0)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^0 \rightarrow \chi_{c1}(3872)K^0) = 1.1 \times 10^{-4}$.

$\Gamma(\pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}$	Γ_3/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

0.043±0.014 OUR AVERAGE

0.043±0.002±0.013 1 AAIJ 20S LHCb $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$

0.047±0.005±0.015 2 CHOI 11 BELL $B^+ \rightarrow \pi^+\pi^-J/\psi K^+$

0.045±0.009±0.014 93 3,4 AUBERT 08Y BABR $B \rightarrow \chi_{c1}(3872)K$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 151 5 BALA 15 BELL $B \rightarrow \chi_{c1}(3872)K\pi$

0.069±0.022±0.022 30 6 AUBERT 05R BABR $B^+ \rightarrow K^+\pi^+\pi^-J/\psi$

0.074±0.016±0.023 36 7 CHOI 03 BELL $B^+ \rightarrow K^+\pi^+\pi^-J/\psi$

1 AAIJ 20S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (7.95 \pm 0.15 \pm 0.33) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 CHOI 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.63 \pm 0.82 \pm 0.52) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

4 superseded by LEES 20C

5 BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872)K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872)K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

6 Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

7 CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] / [B(B^+ \rightarrow \psi(2S)K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S)K^+) = (6.24 \pm 0.21) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\pi^+\pi^-\pi^0 J/\psi(1S))/\Gamma_{\text{total}}$	Γ_4/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

not seen 1 WANG 11B BELL $\gamma(2S) \rightarrow \gamma X$

not seen 2 SHEN 10A BELL $\gamma(1S) \rightarrow \gamma X$

1 WANG 11B reports $B(\gamma(2S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.4 \times 10^{-6}$ at 95% CL.

2 SHEN 10A reports $B(\gamma(1S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.8 \times 10^{-6}$ at 95% CL.

NODE=M176G02
NODE=M176G02

NODE=M176235

NODE=M176R30
NODE=M176R30

OCCUR=2

NODE=M176R30;LINKAGE=A

NODE=M176R30;LINKAGE=E

NODE=M176R30;LINKAGE=D

NODE=M176R6
NODE=M176R6

SYCLP=A

SYCLP=A

SYCLP=A

SYCLP=A

NODE=M176R6;LINKAGE=E

NODE=M176R6;LINKAGE=F

NODE=M176R6;LINKAGE=AB

NODE=M176R6;LINKAGE=C

NODE=M176R6;LINKAGE=A

NODE=M176R6;LINKAGE=AE

NODE=M176R6;LINKAGE=CH

NODE=M176R25
NODE=M176R25

NODE=M176R25;LINKAGE=B

NODE=M176R25;LINKAGE=A

$\Gamma(\omega\eta_c(1S))/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.4	90	1 VINO KUROVA 15	BELL	$B^+ \rightarrow \omega\eta_c K^+$	

¹ VINO KUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 6.9 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$\Gamma(\rho(770)^0 J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$					Γ_6/Γ_3
VALUE (%)	DOCUMENT ID	TECN	COMMENT		
78.6±2.3±2.0	1 AAIJ	23S LHC B	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$		

1 Assuming pure ρ contribution only, i.e. excluding the contribution from $\rho\omega$ interference.
Using $B(\rho^0 \rightarrow \pi^+ \pi^-) = 100\%$.

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

0.032±0.012±0.010	21±7	1 DEL-AMO-SA..10B BABR	$B^+ \rightarrow \omega J/\psi K^+$		
1 DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872) K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi \omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.					

$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$					Γ_7/Γ_3
VALUE	DOCUMENT ID	TECN	COMMENT		

1.16±0.24 OUR AVERAGE	Error includes scale factor of 1.2.				
1.24±0.33±0.10	1,2 AAIJ	23S LHC B	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$		
1.6 ^{+0.4} _{-0.3} ±0.2	3 ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma \omega J/\psi$		
0.8 ±0.3	4 DEL-AMO-SA..10B BABR	B	$B \rightarrow \omega J/\psi K$		

¹ AAIJ 23S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(\omega(782) \rightarrow \pi^+\pi^-)] = (1.9 \pm 0.4 \pm 0.3) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-) = (1.53 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 Excluding $\rho\omega$ interference effects.

3 Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

4 Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	1 AAIJ	17BB LHC B	pp at 7, 8 TeV		

¹ AAIJ 17BB reports $B(b \rightarrow \chi_{c1}(3872) \text{anything}) \times B(\chi_{c1}(3872) \rightarrow \phi\phi) < 4.5 \times 10^{-7}$ at 95% CL.

$\Gamma(D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.55^{+0.20}_{-0.23}±0.17	17	1 GOKHROO	06	BELL	$B^+ \rightarrow D^0\bar{D}^0\pi^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.32	90	2 CHISTOV	04	BELL	Sup. by GOKHROO 06
1 GOKHROO 06	reports	$[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

² CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 0.6 \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$\Gamma(D^0\bar{D}^0\pi^0)/\Gamma(\pi^+\pi^- J/\psi(1S))$					Γ_9/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.16	90	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

NODE=M176R24
NODE=M176R24

NODE=M176R24;LINKAGE=A

NODE=M176R33
NODE=M176R33

NODE=M176R33;LINKAGE=A

NODE=M176R14
NODE=M176R14

NODE=M176R14;LINKAGE=DE

NODE=M176R15
NODE=M176R15

OCCUR=3

NODE=M176R15;LINKAGE=G

NODE=M176R15;LINKAGE=H

NODE=M176R15;LINKAGE=A

NODE=M176R15;LINKAGE=DE

NODE=M176R26
NODE=M176R26

NODE=M176R26;LINKAGE=A

NODE=M176R12
NODE=M176R12

NODE=M176R12;LINKAGE=GO

NODE=M176R12;LINKAGE=A

NODE=M176R17
NODE=M176R17

$\Gamma(\bar{D}^{*0} D^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{10}/Γ
0.46±0.16 OUR AVERAGE					
$0.52^{+0.13}_{-0.11} \pm 0.16$		1,2 HIRATA	23 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$0.42 \pm 0.10 \pm 0.13$	41^{+9}_{-8}	3 AUSHEV	10 BELL	$B^+ \rightarrow D^{*0} \bar{D}^0 K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.90 \pm 0.32 \pm 0.28$	27 ± 6	4 AUBERT	08B BABR	$B^+ \rightarrow \bar{D}^{*0} D^0 K^+$	
$1 \text{ HIRATA } 23 \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.97^{+0.21}_{-0.18} \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					
3 AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$4 \text{ AUBERT } 08B \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M176R13
NODE=M176R13

SYCLP=A

SYCLP=A

NODE=M176R13;LINKAGE=A

NODE=M176R13;LINKAGE=B

NODE=M176R13;LINKAGE=AS

NODE=M176R13;LINKAGE=AU

 $\Gamma(\bar{D}^{*0} D^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{10}/Γ_3
11.77±3.09	50	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

NODE=M176R16
NODE=M176R16 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{11}/Γ
<0.13	90	1 WICHT	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$1 \text{ WICHT } 08 \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 2.4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.					

NODE=M176R09
NODE=M176R09

NODE=M176R09;LINKAGE=A

 $\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ
<0.32	90	1 CHISTOV	04 BELL	$B \rightarrow K D^0 \bar{D}^0$	
$1 \text{ CHISTOV } 04 \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 6 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.					

NODE=M176R3
NODE=M176R3

NODE=M176R3;LINKAGE=A

 $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ
<0.22	90	1 CHISTOV	04 BELL	$B \rightarrow K D^+ D^-$	
$1 \text{ CHISTOV } 04 \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow D^+ D^-)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.					

NODE=M176R4
NODE=M176R4

NODE=M176R4;LINKAGE=A

 $\Gamma(\pi^0 \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ_3
<1.1	90	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

NODE=M176R06
NODE=M176R06 $\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.04	90	1 BHARDWAJ	19 BELL	$B^\pm \rightarrow \pi^0 \chi_{c1} K^\pm$	
$1 \text{ BHARDWAJ } 19 \text{ reports } [\Gamma(\chi_{c1}(3872) \rightarrow \pi^0 \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 8.1 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.					

NODE=M176R23
NODE=M176R23

NODE=M176R23;LINKAGE=A

 $\Gamma(\pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ_3
88⁺³³₋₂₇±10	10.8	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

NODE=M176R05
NODE=M176R05

$\Gamma(\pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<19	90	ABLIKIM	19U	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{16}/Γ_3 NODE=M176R04
NODE=M176R04 $\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	1 VINOKUROVA	15	BELL $B^+ \rightarrow \pi^+ \pi^- \eta_c K^+$
1 VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 3.0 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.				

 Γ_{17}/Γ NODE=M176R22
NODE=M176R22

NODE=M176R22;LINKAGE=A

 $\Gamma(\pi^0 \pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{18}/Γ_3 NODE=M176R28
NODE=M176R28 $\Gamma(\pi^0 \pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	24BU	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{19}/Γ_3 NODE=M176R37
NODE=M176R37 $\Gamma(\pi^0 \pi^0 \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	ABLIKIM	24BU	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{20}/Γ_3 NODE=M176R38
NODE=M176R38 $\Gamma(\pi^+ \pi^- \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.56	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{21}/Γ_3 NODE=M176R29
NODE=M176R29 $\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 × 10 ⁻³	90	1 BHARDWAJ	16	BELL $B^+ \rightarrow \pi^+ \pi^- \chi_{c1} K^+$
1 BHARDWAJ 16 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.				

 Γ_{22}/Γ NODE=M176R00
NODE=M176R00

NODE=M176R00;LINKAGE=A

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.7 × 10 ⁻⁵	95	1 AAIJ	17AD	LHCb $B^+ \rightarrow p\bar{p} K^+$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<9 × 10 ⁻⁵	95	2 AAIJ	13S	LHCb $B^+ \rightarrow p\bar{p} K^+$
1 AAIJ 17AD reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 0.5 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.				
2 AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.				

 Γ_{23}/Γ NODE=M176R03
NODE=M176R03

SYCLP=A

SYCLP=A

NODE=M176R03;LINKAGE=C

NODE=M176R03;LINKAGE=B

 $\Gamma(\pi^+ \pi^- \eta)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.12	90	ABLIKIM	24K	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{24}/Γ_3 NODE=M176R35
NODE=M176R35

NODE=M176410

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.99	90	ABLIKIM	20W	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{25}/Γ_3 NODE=M176R20
NODE=M176R20 $\Gamma(\gamma D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.58	90	ABLIKIM	20W	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 Γ_{26}/Γ_3 NODE=M176R21
NODE=M176R21

$\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{27}/Γ
9.6\pm2.7\pm3.0		1 BHARDWAJ	11 BELL	$B^\pm \rightarrow \gamma J/\psi K^\pm$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

15 \pm 4 \pm 5	20	2 AUBERT	09B BABR	$B^+ \rightarrow \gamma J/\psi K^+$	
18 \pm 6 \pm 6	19	3 AUBERT,BE	06M BABR	$B^+ \rightarrow \gamma J/\psi K^+$	

1 BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 Superseded by AUBERT 09B. AUBERT,BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\gamma J/\psi)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{27}/Γ_3
0.52\pm0.19 OUR AVERAGE				Error includes scale factor of 1.2.	
0.38 \pm 0.20 \pm 0.01	8 \pm 4	ABLIKIM	24X BES3	$e^+ e^- \rightarrow \omega \chi_{c1}(3872)$	
0.79 \pm 0.28		ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{28}/Γ
<0.010	90	1 BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$	
1 BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.					

 $\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{28}/Γ_3
<0.89	90	CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$	

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{29}/Γ
<0.04	90	1 BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2} \gamma K^\pm$	
1 BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.					

 $\Gamma(\gamma \psi(2S))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ
possibly seen	36 \pm 9	1 AAIJ	14AH LHCb	$B^+ \rightarrow \gamma \psi(2S) K^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		2 BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma \psi(2S) K^+$
0.051 \pm 0.015 \pm 0.016	25 \pm 7	3 AUBERT	09B BABR	$B^+ \rightarrow \gamma \psi(2S) K^+$

1 From 36.4 \pm 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi \gamma$ decays with a statistical significance of 4.4σ .

2 BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma \psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

3 AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\gamma \psi(2S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ_3
<0.42	90	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

 $\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ_3
<0.18	90	ABLIKIM	24S BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

NODE=M176R7

NODE=M176R7

SYCLP=A

SYCLP=A

SYCLP=A

NODE=M176R7;LINKAGE=AB

NODE=M176R7;LINKAGE=AU

NODE=M176R18
NODE=M176R18

NODE=M176R08
NODE=M176R08

NODE=M176R08;LINKAGE=B

NODE=M176R1
NODE=M176R1

NODE=M176R01
NODE=M176R01

NODE=M176R01;LINKAGE=B

NODE=M176R10
NODE=M176R10

NODE=M176R10;LINKAGE=A

NODE=M176R10;LINKAGE=BH

NODE=M176R10;LINKAGE=AU

NODE=M176R19
NODE=M176R19

NODE=M176R34
NODE=M176R34

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ_{27}
1.67±0.21±0.13			1 AAIJ	24AD LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$ and $\gamma J/\psi K^+$	
<0.59	90		ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma\chi_{c1}(3872)$	
2.46±0.64±0.29	36 ± 9		2 AAIJ	14AH LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$	
<2.1	90		BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$	
3.4 ± 1.4			AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c}K'$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.59	90	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma\chi_{c1}(3872)$
2.46±0.64±0.29	36 ± 9	2 AAIJ	14AH LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$
<2.1	90	BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$
3.4 ± 1.4		AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c}K'$

¹ AAIJ 24AD reports this ratio as $1.67 \pm 0.21 \pm 0.12 \pm 0.04$ where the last uncertainty is due to the uncertainties of the branching fractions of $\psi(2S)$ and J/ψ mesons. We have added the last two uncertainties in quadrature.

² From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ . Superseded by AAIJ 24AD.

 $\Gamma(\gamma\psi_2(3823))/\Gamma(\pi^+\pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ_3
<0.075	90	1 ABLIKIM	24Z BES3	$e^+ e^- \rightarrow \gamma\psi_2(3823)$	

¹ Using $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.343 \pm 0.010$. We have assumed that $B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = 1$.

C-violating decays $\Gamma(\eta J/\psi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{32}/Γ
<0.021	90	1,2 IWASHITA	14 BELL	$B \rightarrow K\eta J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	3 AUBERT	04Y BABR	$B \rightarrow K\eta J/\psi$
-------	----	----------	----------	------------------------------

¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.

³ AUBERT 04Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 7.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

 $\chi_{c1}(3872)$ REFERENCES

AAIJ	24AD	JHEP 2411 121	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	24BU	PR D110 072015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24C	PRL 132 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24K	PR D109 L011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24S	PR D109 L071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24X	PR D110 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24Z	PR D110 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AP	JHEP 2307 084	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	23S	PR D108 L011103	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23O	PR D107 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23W	PRL 130 151904	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HIRATA	23	PR D107 112011	H. Hirata <i>et al.</i>	(BELLE Collab.)
YIN	23	PR D107 052004	J.H. Yin <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22D	PR D105 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
TERAMOTO	21	PRL 126 122001	Y. Teramoto <i>et al.</i>	(BELLE Collab.)
AAIJ	20AD	PR D102 092005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	20W	PRL 124 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	20C	PRL 124 152001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AGHASIYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M176R11
NODE=M176R11

NODE=M176R11;LINKAGE=C

NODE=M176R11;LINKAGE=A

NODE=M176R36
NODE=M176R36

NODE=M176R36;LINKAGE=A

NODE=M176R2
NODE=M176R2

NODE=M176R2;LINKAGE=A

NODE=M176R2;LINKAGE=C

NODE=M176R2;LINKAGE=D

NODE=M176

REFID=63038
REFID=63026
REFID=62652
REFID=62661
REFID=62670
REFID=62677
REFID=62679
REFID=62412
REFID=62287
REFID=62058
REFID=62068
REFID=62281
REFID=61909
REFID=61638
REFID=61634
REFID=61098
REFID=60673
REFID=60526
REFID=60570
REFID=60554
REFID=60676
REFID=58191
REFID=57272
REFID=56771
REFID=56787
REFID=56408
REFID=56706
REFID=57795
REFID=55897
REFID=55647
REFID=55925
REFID=54985
REFID=55008

BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=53779
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=53934
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53098
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52722
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52265
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT,BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51454
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)	REFID=51432
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50498
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50627
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=50458
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50200
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49742
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49997
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>		REFID=49677
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=49628
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074