

CHARMED, STRANGE MESONS ($C = \pm 1, S = \pm 1$) (including possibly non- $q\bar{q}$ states)

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \quad \text{similarly for } D_s^{*'}s$$

D_s^\pm

$$I(J^P) = 0(0^-)$$

$$\text{Mass } m = 1968.35 \pm 0.07 \text{ MeV}$$

$$m_{D_s^\pm} - m_{D^\pm} = 98.69 \pm 0.05 \text{ MeV}$$

$$\text{Mean life } \tau = (501.2 \pm 2.2) \times 10^{-15} \text{ s} \quad (S = 1.3)$$

$$c\tau = 150.3 \mu\text{m}$$

CP-violating decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (-0.2 \pm 2.5)\%$$

$$A_{CP}(\tau^\pm \nu) \text{ in } D_s^+ \rightarrow \tau^+ \nu_\tau, D_s^- \rightarrow \tau^- \bar{\nu}_\tau = (3 \pm 5)\%$$

$$A_{CP}(K^\pm K_S^0) = (0.07 \pm 0.24)\%$$

$$A_{CP}(K^\pm K_L^0) \text{ in } D_s^\pm \rightarrow K^\pm K_L^0 = (-1.1 \pm 2.7) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.35 \pm 0.34)\%$$

$$A_{CP}(\phi \pi^\pm) = (-0.38 \pm 0.27)\%$$

$$A_{CP}(K^\pm K_S^0 \pi^0) = (-0.9 \pm 1.9)\%$$

$$A_{CP}(2K_S^0 \pi^\pm) = (1.3 \pm 1.6)\%$$

$$A_{CP}(K^+ K^- \pi^\pm \pi^0) = (-0.6 \pm 0.9)\%$$

$$A_{CP}(K^\pm K_S^0 \pi^+ \pi^-) = (0.7 \pm 2.9)\% \quad (S = 1.3)$$

$$A_{CP}(K_S^0 K^\mp 2\pi^\pm) = (0.7 \pm 1.8)\% \quad (S = 1.3)$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-0.9 \pm 1.1)\%$$

$$A_{CP}(\pi^\pm \eta) = (0.24 \pm 0.29)\%$$

$$A_{CP}(\pi^\pm \pi^+ \pi^- \eta) = (2.4 \pm 3.0)\%$$

$$A_{CP}(\pi^\pm \eta') = (-0.08 \pm 0.17)\% \quad (S = 1.2)$$

$$A_{CP}(\eta \pi^\pm \pi^0) = (0.9 \pm 1.5)\%$$

$$A_{CP}(\eta' \pi^\pm \pi^0) = (-1.5 \pm 2.5)\%$$

$$A_{CP}(K^\pm \pi^0) = (2 \pm 4)\% \quad (S = 1.2)$$

$$A_{CP}(\bar{K}^0 / K^0 \pi^\pm) = (0.4 \pm 0.5)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (0.20 \pm 0.18)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^-) = (2.2 \pm 1.9)\%$$

$$A_{CP}(K_S^0 \pi^+ \pi^0) \text{ in } D_s^\pm \rightarrow K_S^0 \pi^\pm \pi^0 = (-2 \pm 5)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^- \pi^0) \text{ in } D_s^\pm \rightarrow K^\pm \pi^+ \pi^- \pi^0 = (7 \pm 5) \times 10^{-2}$$

$$A_{CP}(K^\pm \eta) = (1.8 \pm 1.9)\%$$

$$A_{CP}(K^\pm \eta'(958)) = (6 \pm 19)\%$$

CP violating asymmetries of P-odd (T-odd) moments

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-8 \pm 6) \times 10^{-3} \text{ [a]}$$

NODE=MXXX040

NODE=S034

NODE=S034M;DTYPE=M

NODE=S034DM;DTYPE=D

NODE=S034T;DTYPE=T

NODE=S034CTA;DTYPE=C;OUR EVAL

CLUMP=P

NODE=S034A13;DTYPE=v;CLUMP=P

NODE=S034A30;DTYPE=v;CLUMP=P

NODE=S034A05;DTYPE=v;CLUMP=P

NODE=S034A00;DTYPE=v;CLUMP=P

NODE=S034A06;DTYPE=v;CLUMP=P

NODE=S034A20;DTYPE=v;CLUMP=P

NODE=S034A15;DTYPE=v;CLUMP=P

NODE=S034A16;DTYPE=v;CLUMP=P

NODE=S034A07;DTYPE=v;CLUMP=P

NODE=S034A14;DTYPE=v;CLUMP=P

NODE=S034A08;DTYPE=v;CLUMP=P

NODE=S034A09;DTYPE=v;CLUMP=P

NODE=S034A10;DTYPE=v;CLUMP=P

NODE=S034A48;DTYPE=v;CLUMP=P

NODE=S034A11;DTYPE=v;CLUMP=P

NODE=S034A19;DTYPE=v;CLUMP=P

NODE=S034A17;DTYPE=v;CLUMP=P

NODE=S034A01;DTYPE=v;CLUMP=P

NODE=S034A21;DTYPE=v;CLUMP=P

NODE=S034A02;DTYPE=v;CLUMP=P

NODE=S034A12;DTYPE=v;CLUMP=P

NODE=S034A28;DTYPE=v;CLUMP=P

NODE=S034A39;DTYPE=v;CLUMP=P

NODE=S034A03;DTYPE=v;CLUMP=P

NODE=S034A04;DTYPE=v;CLUMP=P

CLUMP=T

NODE=S034TV0;DTYPE=t;CLUMP=T

$D_s^+ \rightarrow \phi \ell^+ \nu_\ell$ form factors

$$\begin{aligned}
r_2 &= 0.83 \pm 0.08 \quad (S = 1.8) \\
r_V &= 1.76 \pm 0.07 \quad (S = 1.1) \\
\Gamma_L/\Gamma_T &= 0.72 \pm 0.18 \\
f_+(0) |V_{cs}| \text{ in } D_s^+ &\rightarrow \eta e^+ \nu_e = 0.449 \pm 0.009 \\
f_+(0) |V_{cs}| \text{ in } D_s^+ &\rightarrow \eta \mu^+ \nu_\mu = 0.452 \pm 0.012 \\
r_1 \equiv a_1/a_0 \text{ in } D_s^+ &\rightarrow \eta \mu^+ \nu_\mu = -2.9 \pm 0.6 \\
\langle A_{FB}^\eta \rangle \text{ in } D_s^+ &\rightarrow \eta \mu^+ \nu_\mu = (-5.9 \pm 3.1) \times 10^{-2} \\
f_+(0) |V_{cs}| \text{ in } D_s^+ &\rightarrow \eta' e^+ \nu_e = 0.527 \pm 0.024 \\
r_1 \equiv a_1/a_0 \text{ in } D_s^+ &\rightarrow \eta' \mu^+ \nu_\mu = -11 \pm 5 \\
\langle A_{FB}^{\eta'} \rangle \text{ in } D_s^+ &\rightarrow \eta' \mu^+ \nu_\mu = (-6 \pm 8) \times 10^{-2} \\
f_+(0) |V_{cs}| \text{ in } D_s^+ &\rightarrow \eta' \mu^+ \nu_\mu = 0.50 \pm 0.04 \\
f_+(0) |V_{cs}| \text{ in } D_s^+ &\rightarrow f_0(980) e^+ \nu_e = 0.50 \pm 0.04 \\
f_+(0) |V_{cd}| \text{ in } D_s^+ &\rightarrow K^0 e^+ \nu_e = 0.145 \pm 0.010 \\
r_V \equiv V(0)/A_1(0) \text{ in } D_s^+ &\rightarrow K^*(892)^0 e^+ \nu_e = 1.7 \pm 0.4 \\
r_2 \equiv A_2(0)/A_1(0) \text{ in } D_s^+ &\rightarrow K^*(892)^0 e^+ \nu_e = 0.77 \pm 0.29 \\
f_{D_s^+} |V_{cs}| \text{ in } D_s^+ &\rightarrow \mu^+ \nu_\mu = 242.7 \pm 3.0 \text{ MeV} \\
f_{D_s^+} |V_{cs}| \text{ in } D_s^+ &\rightarrow \tau^+ \nu_\tau = 247.5 \pm 2.3 \text{ MeV}
\end{aligned}$$

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

CLUMP=F
NODE=S034FR2;DTYPE=f;CLUMP=F
NODE=S034FRV;DTYPE=f;CLUMP=F
NODE=S034GLT;DTYPE=f;CLUMP=F
NODE=S034A26;DTYPE=f;CLUMP=F
NODE=S034A42;DTYPE=f;CLUMP=F
NODE=S034A44;DTYPE=f;CLUMP=F
NODE=S034A46;DTYPE=f;CLUMP=F
NODE=S034A27;DTYPE=f;CLUMP=F
NODE=S034A45;DTYPE=f;CLUMP=F
NODE=S034A47;DTYPE=f;CLUMP=F
NODE=S034A43;DTYPE=f;CLUMP=F
NODE=S034A50;DTYPE=f;CLUMP=F
NODE=S034A22;DTYPE=t;CLUMP=F
NODE=S034A23;DTYPE=t;CLUMP=F
NODE=S034A24;DTYPE=t;CLUMP=F
NODE=S034A25;DTYPE=t;CLUMP=F
NODE=S034A29;DTYPE=t;CLUMP=F
NODE=S034215;NODE=S034

D_s^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	[b] (6.33 \pm 0.15) %		—
π^+ anything	(119.3 \pm 1.4) %		—
π^- anything	(43.2 \pm 0.9) %		—
π^0 anything	(123 \pm 7) %		—
K^- anything	(18.7 \pm 0.5) %		—
K^+ anything	(28.9 \pm 0.7) %		—
K_S^0 anything	(19.0 \pm 1.1) %		—
η anything	[c] (29.9 \pm 2.8) %		—
ω anything	(6.1 \pm 1.4) %		—
η' anything	[d] (10.3 \pm 1.4) %	S=1.1	—
$f_0(980)$ anything, $f_0 \rightarrow \pi^+ \pi^-$	< 1.3 %	CL=90%	—
ϕ anything	(15.7 \pm 1.0) %		—
$K^+ K^-$ anything	(15.8 \pm 0.7) %		—
$K_S^0 K^+$ anything	(5.8 \pm 0.5) %		—
$K_S^0 K^-$ anything	(1.9 \pm 0.4) %		—
$2K_S^0$ anything	(1.70 \pm 0.32) %		—
$2K^+$ anything	< 2.6 $\times 10^{-3}$	CL=90%	—
$2K^-$ anything	< 6 $\times 10^{-4}$	CL=90%	—
$2\pi^+ \pi^- +$ anything	(32.8 \pm 0.7) %		—

NODE=S034;CLUMP=D
DESIG=25
DESIG=133
DESIG=134
DESIG=135
DESIG=32
DESIG=33
DESIG=138
DESIG=2
DESIG=136
DESIG=117
DESIG=137
DESIG=92
DESIG=139
DESIG=140
DESIG=141
DESIG=142
DESIG=143
DESIG=144
DESIG=236

Leptonic and semileptonic modes

$e^+ \nu_e$	< 8.3	$\times 10^{-5}$ CL=90%	984	NODE=S034;CLUMP=C DESIG=118
$\mu^+ \nu_\mu$	(5.37 \pm 0.11)	$\times 10^{-3}$	981	DESIG=7
$\tau^+ \nu_\tau$	(5.39 \pm 0.09)	%	182	DESIG=89
$\gamma e^+ \nu_e$	< 1.3	$\times 10^{-4}$ CL=90%	984	DESIG=169
$K^+ K^- e^+ \nu_e$	—		851	DESIG=125;OUR EVAL;→ UNCHECKED ←
$K_S^0 K_S^0 e^+ \nu_e$	< 3.8	$\times 10^{-4}$ CL=90%	849	DESIG=206
$\phi e^+ \nu_e$	[e] (2.34 \pm 0.12)	% S=1.2	720	DESIG=30
$K_1(1270)^0 e^+ \nu_e$	< 4.1	$\times 10^{-4}$ CL=90%	585	DESIG=243
$b_1(1235)^0 e^+ \nu_e, b_1^0 \rightarrow \omega \pi^0$	< 6.4	$\times 10^{-4}$ CL=90%	—	DESIG=244
$\phi \mu^+ \nu_\mu$	(2.24 \pm 0.11)	%	715	DESIG=166
$\eta e^+ \nu_e$	[e] (2.27 \pm 0.06)	%	908	DESIG=86
$\eta'(958) e^+ \nu_e$	[e] (8.1 \pm 0.4)	$\times 10^{-3}$	751	DESIG=87
$\eta \mu^+ \nu_\mu$	(2.24 \pm 0.07)	%	905	DESIG=167
$\eta'(958) \mu^+ \nu_\mu$	(8.0 \pm 0.6)	$\times 10^{-3}$	747	DESIG=168
$\omega e^+ \nu_e$	[f] < 2.0	$\times 10^{-3}$ CL=90%	829	DESIG=153
$K^0 e^+ \nu_e$	(2.88 \pm 0.26)	$\times 10^{-3}$ S=1.2	921	DESIG=145
$K^*(892)^0 e^+ \nu_e$	[e] (2.05 \pm 0.20)	$\times 10^{-3}$	782	DESIG=146
$f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0$	< 7.3	$\times 10^{-4}$ CL=90%	—	DESIG=205
$f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$	< 3.3	$\times 10^{-4}$ CL=90%	—	DESIG=265
$f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0$	(7.9 \pm 1.5)	$\times 10^{-4}$	—	DESIG=204
$f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$	(1.64 \pm 0.13)	$\times 10^{-3}$	—	DESIG=147
$f_0(980) \mu^+ \nu_\mu, f_0 \rightarrow K^+ K^-$	< 5.45	$\times 10^{-4}$ CL=90%	—	DESIG=246
$a_0(980)^0 e^+ \nu_e, a_0^0 \rightarrow \pi^0 \eta$	< 1.2	$\times 10^{-4}$ CL=90%	—	DESIG=175
$\pi^0 e^+ \nu_e$	< 6.4	$\times 10^{-5}$ CL=90%	980	DESIG=216

Hadronic modes with a $K\bar{K}$ pair

$K^+ K_S^0$	(1.500 \pm 0.014)	%	850	NODE=S034;CLUMP=A DESIG=22
$K^+ K_L^0$	(1.49 \pm 0.06)	%	850	DESIG=170
$K^+ \bar{K}^0$	(2.95 \pm 0.14)	%	850	DESIG=164
$K^+ K^- \pi^+$	[g] (5.45 \pm 0.08)	% S=1.3	805	DESIG=40
$\phi \pi^+$	[e,h] (4.5 \pm 0.4)	%	712	DESIG=6
$\phi \pi^+, \phi \rightarrow K^+ K^-$	[h] (2.25 \pm 0.05)	%	712	DESIG=114
$K^+ \bar{K}^*(892)^0$	(12.7 \pm 4.0 -3.1)	%	685	DESIG=232
$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	(2.61 \pm 0.05)	%	416	DESIG=115
$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$ $K_S^0 \pi^0$	(4.8 \pm 0.4)	$\times 10^{-3}$	—	DESIG=227
$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$	(1.12 \pm 0.19)	%	732	DESIG=79
$f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$	(7.2 \pm 3.0)	$\times 10^{-4}$	—	DESIG=148
$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$	(6.8 \pm 2.8)	$\times 10^{-4}$	198	DESIG=88
$a_0(980)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$	(1.1 \pm 0.4)	$\times 10^{-3}$	—	DESIG=229
$a_0(1710)^+ \pi^0, a_0^+ \rightarrow$ $K^+ K_S^0$	(3.5 \pm 0.6)	$\times 10^{-3}$	—	DESIG=231
$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow$ $K^- \pi^+$	(1.79 \pm 0.26)	$\times 10^{-3}$	218	DESIG=80
$K^+ \bar{K}_0^*(1410)^0, \bar{K}_0^* \rightarrow$ $K_S^0 \pi^0$	(8.8 \pm 2.8)	$\times 10^{-4}$	—	DESIG=230
$K^+ K_S^0 \pi^0$	(1.471 \pm 0.028)	%	805	DESIG=162
$K^*(892)^+ K_S^0, K^{*+} \rightarrow$ $K^+ \pi^0$	(2.04 \pm 0.32)	$\times 10^{-3}$	—	DESIG=228
$2K_S^0 \pi^+$	(7.32 \pm 0.14)	$\times 10^{-3}$	802	DESIG=163
$f_0(980) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	< 1.8	$\times 10^{-4}$ CL=90%	—	DESIG=212
$f_0(1710) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	(3.39 \pm 0.31)	$\times 10^{-3}$	—	DESIG=210
$K^*(892)^+ K_S^0, K^{*+} \rightarrow$ $K_S^0 \pi^+$	(3.19 \pm 0.29)	$\times 10^{-3}$	683	DESIG=211
$K^0 \bar{K}^0 \pi^+$	—		802	DESIG=46;OUR EVAL;→ UNCHECKED ←
$K^*(892)^+ \bar{K}^0$	[e] (5.4 \pm 1.2)	%	683	DESIG=24

$K^+ K^- \pi^+ \pi^0$	(5.53 ±0.15) %	S=1.3	748	DESIG=41
$\phi \rho^+$	[e] (5.59 ±0.34) %		401	DESIG=5
$\bar{K}_1(1270)^0 K^+,$ $\bar{K}_1(1270)^0 \rightarrow K^- \rho^+$	(5.7 ±0.6) × 10 ⁻³		-	DESIG=177
$\bar{K}_1(1270)^0 K^+,$ $\bar{K}_1(1270)^0 \rightarrow K^*(892)\pi$	(1.31 ±0.25) %		-	DESIG=176
$\bar{K}_1(1400)^0 K^+,$ $\bar{K}_1(1400)^0 \rightarrow K^*(892)\pi$	(2.0 ±0.4) %		-	DESIG=178
$a_0(980)^0 \rho^+, a_0^0 \rightarrow K^+ K^-$	(1.9 ±0.4) × 10 ⁻³		-	DESIG=179
$f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow$ $K^*(892)^\mp K^\pm$	(3.9 ±0.7) × 10 ⁻³		-	DESIG=180
$f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow$ $a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow$ $K^+ K^-$	(4.0 ±1.4) × 10 ⁻⁴		-	DESIG=181
$\eta(1475)\pi^+, \eta(1475) \rightarrow$ $a_0(980)^0 \pi^0, a_0(980)^0 \rightarrow$ $K^+ K^-$	(7.0 ±2.8) × 10 ⁻⁴		-	DESIG=182
$K_S^0 K^- 2\pi^+$	(1.569±0.028) %		744	DESIG=26
$K^+ K^- K_S^0 \pi^+$	(1.27 ±0.15) × 10 ⁻⁴		527	DESIG=245
$K^*(892)^+ \bar{K}^*(892)^0$	[e] (5.64 ±0.35) %		417	DESIG=27
$\eta(1475) K_S^0, \eta \rightarrow$ $K^*(892)^0 \pi^+, K^{*0} \rightarrow$ $K^- \pi^+$	(3.5 ±1.0) × 10 ⁻⁴		-	DESIG=199
$\eta(1475)\pi^+, \eta \rightarrow$ $\bar{K}^*(892)^+ K^-, \bar{K}^{*+} \rightarrow$ $K_S^0 \pi^+$	(3.5 ±1.0) × 10 ⁻⁴		-	DESIG=200
$\eta(1475)\pi^+, \eta \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(1.7 ±0.9) × 10 ⁻³		-	DESIG=196
$f_1(1285)\pi^+, f_1 \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(3.5 ±0.8) × 10 ⁻⁴		-	DESIG=202
$K^+ K_S^0 \pi^+ \pi^-$	(9.34 ±0.22) × 10 ⁻³		744	DESIG=43
$K^+ K^- 2\pi^+ \pi^-$	(6.6 ±0.6) × 10 ⁻³		673	DESIG=42
$\phi 2\pi^+ \pi^-$	[e] (1.21 ±0.16) %		640	DESIG=8
$\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-$	(4.9 ±0.7) × 10 ⁻³		181	DESIG=105
$\phi a_1(1260)^+, \phi \rightarrow$ $K^+ K^-, a_1^+ \rightarrow$ $\rho^0 \pi^+$	(7.5 ±1.2) × 10 ⁻³		†	DESIG=106
$\phi 2\pi^+ \pi^- \text{ non-}\rho, \phi \rightarrow$ $K^+ K^-$	(1.4 ±0.5) × 10 ⁻³		-	DESIG=174
$K^+ K^- \rho^0 \pi^+ \text{ non-}\phi$	< 2.0 × 10 ⁻⁴ CL=90%		249	DESIG=104
$K^+ K^- 2\pi^+ \pi^- \text{ nonresonant}$	(1.0 ±0.4) × 10 ⁻³		673	DESIG=107
$2K_S^0 2\pi^+ \pi^-$	(8.0 ±3.3) × 10 ⁻⁴		669	DESIG=112

Hadronic modes without K's

$\pi^+ \pi^0$	< 1.2 × 10 ⁻⁴ CL=90%		975	NODE=S034;CLUMP=B DESIG=119
$2\pi^+ \pi^-$	(1.090±0.014) %		959	DESIG=15
$\rho^0 \pi^+$	(1.14 ±0.16) × 10 ⁻⁴		825	DESIG=10
$\omega \pi^+, \omega \rightarrow \pi^+ \pi^-$	(3.9 ±0.4) × 10 ⁻⁵		-	DESIG=233
$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	[i] (9.23 ±0.13) × 10 ⁻³		959	DESIG=113
$f_2(1270)\pi^+, f_2 \rightarrow \pi^+ \pi^-$	(1.42 ±0.10) × 10 ⁻³		559	DESIG=90
$f_2'(1525)^0 \pi^+, f_2' \rightarrow \pi^+ \pi^-$	(5.8 ±2.0) × 10 ⁻⁶		-	DESIG=235
$\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	(1.8 ±0.6) × 10 ⁻⁴		421	DESIG=103
$\rho(1700)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	(4 ±4) × 10 ⁻⁵		-	DESIG=234
$\pi^+ 2\pi^0$	(5.2 ±0.5) × 10 ⁻³	S=1.1	961	DESIG=149
$f_0(980)\pi^+, f_0 \rightarrow \pi^0 \pi^0$	(2.9 ±0.6) × 10 ⁻³		-	DESIG=213
$f_0(1370)\pi^+, f_0 \rightarrow \pi^0 \pi^0$	(1.3 ±0.6) × 10 ⁻³		-	DESIG=214
$f_2(1270)\pi^+, f_2 \rightarrow \pi^0 \pi^0$	(5.0 ±3.5) × 10 ⁻⁴		-	DESIG=215
$2\pi^+ \pi^- \pi^0$	—		935	DESIG=18;OUR EVAL;→ UNCHECKED ←

$\omega\pi^+$	[e]	$(1.93 \pm 0.30) \times 10^{-3}$	822	DESIG=19
$\eta\pi^+$	[e]	$(1.686 \pm 0.027) \%$	902	DESIG=1
$(2\pi^+\pi^-\pi^0)_{\text{non-}\eta}$		$(2.04 \pm 0.09) \%$	—	DESIG=253
$f_0(1370)^0\rho^+$,		$(5.1 \pm 0.9) \times 10^{-3}$	—	DESIG=254
$f_0(1370)^0\rho^+ \rightarrow$				
$2\pi^+\pi^-\pi^0$				
$f_0(980)^0\rho^+$,		$(2.6 \pm 0.5) \times 10^{-3}$	—	DESIG=255
$f_0(980)^0\rho^+ \rightarrow$				
$2\pi^+\pi^-\pi^0$				
$f_2(1270)^0\rho^+$,		$(1.9 \pm 0.4) \times 10^{-3}$	—	DESIG=256
$f_2(1270)^0\rho^+ \rightarrow$				
$2\pi^+\pi^-\pi^0$				
$(\rho^+\rho^0)_{S\text{-wave}} \rightarrow$		$(7.1 \pm 2.8) \times 10^{-4}$	—	DESIG=257
$2\pi^+\pi^-\pi^0$				
$(\rho(1450)^+\rho^0)_{S\text{-wave}} \rightarrow$		$(9.4 \pm 3.1) \times 10^{-4}$	—	DESIG=258
$2\pi^+\pi^-\pi^0$				
$(\rho^+\rho(1450)^0)_{P\text{-wave}} \rightarrow$		$(1.75 \pm 0.29) \times 10^{-3}$	—	DESIG=259
$2\pi^+\pi^-\pi^0$				
$\phi\pi^+$, $\phi \rightarrow \rho\pi$		$(5.08 \pm 0.35) \times 10^{-3}$	—	DESIG=260
$\omega\pi^+$, $\omega \rightarrow \rho\pi$		$(1.41 \pm 0.19) \times 10^{-3}$	—	DESIG=261
$a_1(1260)^+\pi^0$, $a_1^+ \rightarrow$		$(2.6 \pm 0.4) \times 10^{-3}$	—	DESIG=262
$(\rho^0\pi^+)_{S\text{-wave}}$				
$a_1(1260)^0\pi^+$, $a_1^0 \rightarrow$		$(1.3 \pm 0.5) \times 10^{-3}$	—	DESIG=263
$(\rho\pi)_{S\text{-wave}}$				
$\pi(1300)^0\pi^+$, $\pi(1300)^0 \rightarrow$		$(2.4 \pm 0.7) \times 10^{-3}$	—	DESIG=264
$(\rho\pi)_{P\text{-wave}}$				
$3\pi^+2\pi^-$		$(8.0 \pm 0.8) \times 10^{-3}$	899	DESIG=17
$2\pi^+\pi^-2\pi^0$		—	902	DESIG=57;OUR EVAL;→ UNCHECKED ←
$\eta\rho^+$	[e]	$(8.9 \pm 0.8) \%$	724	DESIG=58
$\eta\pi^+\pi^0$		$(9.10 \pm 0.17) \%$	885	DESIG=160
$\eta(\pi^+\pi^0)_{P\text{-wave}}$		$(4.9 \pm 3.0) \times 10^{-3}$	885	DESIG=172
$a_0(980)^+\pi^0$,		$(2.1 \pm 0.4) \%$	—	DESIG=173
$a_0(980)^+\pi^0 \rightarrow \eta\pi^+\pi^0$				
$\omega\pi^+\pi^0$	[e]	$(2.8 \pm 0.7) \%$	802	DESIG=126
$2\pi^+\pi^-\eta$		$(3.08 \pm 0.08) \%$	855	DESIG=183
$a_1(1260)^+\eta$, $a_1^+ \rightarrow$		$(1.71 \pm 0.14) \%$	—	DESIG=184
$\rho(770)^0\pi^+$, $\rho^0 \rightarrow$				
$\pi^+\pi^-$				
$a_1(1260)^+\eta$, $a_1^+ \rightarrow$		$(2.5 \pm 0.9) \times 10^{-3}$	—	DESIG=185
$f_0(500)\pi^+$, $f_0 \rightarrow \pi^+\pi^-$				
$a_0(980)^+\rho(770)^0$, $a_0^+ \rightarrow$		$(2.1 \pm 0.9) \times 10^{-3}$	—	DESIG=186
$\eta\pi^+$				
$\eta(1405)\pi^+$, $\eta(1405) \rightarrow$		$(2.2 \pm 0.7) \times 10^{-4}$	—	DESIG=187
$a_0(980)^-\pi^+$, $a_0^- \rightarrow$				
$\eta\pi^-$				
$\eta(1405)\pi^+$, $\eta(1405) \rightarrow$		$(2.2 \pm 0.7) \times 10^{-4}$	—	DESIG=188
$a_0(980)^+\pi^-$, $a_0^+ \rightarrow$				
$\eta\pi^+$				
$f_1(1420)\pi^+$, $f_1 \rightarrow$		$(5.9 \pm 1.8) \times 10^{-4}$	—	DESIG=195
$a_0(980)^-\pi^+$, $a_0^- \rightarrow$				
$\eta\pi^-$				
$f_1(1420)\pi^+$, $f_1 \rightarrow$		$(5.2 \pm 1.8) \times 10^{-4}$	—	DESIG=190
$a_0(980)^+\pi^-$, $a_0^+ \rightarrow$				
$\eta\pi^+$				
$3\pi^+2\pi^-\pi^0$		$(4.9 \pm 3.2) \%$	856	DESIG=59
$\omega 2\pi^+\pi^-$	[e]	$(1.6 \pm 0.5) \%$	766	DESIG=127
$\eta'(958)\pi^+$	[d,e]	$(3.95 \pm 0.08) \%$	743	DESIG=13
$3\pi^+2\pi^-2\pi^0$		—	803	DESIG=61;OUR EVAL;→ UNCHECKED ←
$\omega\eta\pi^+$	[e]	$(5.4 \pm 1.3) \times 10^{-3}$	654	DESIG=128
$\eta'(958)\rho^+$	[d,e]	$(5.8 \pm 1.5) \%$	465	DESIG=62
$\eta'(958)\pi^+\pi^0$		$(6.14 \pm 0.18) \%$	720	DESIG=161
$\eta'(958)\pi^+\pi^0$ nonresonant	<	5.1 %	720	DESIG=165
				CL=90%

Modes with one or three K's

				NODE=S034;CLUMP=K
$K^+ \pi^0$	(7.5 ±0.5) × 10 ⁻⁴	917	DESIG=120	
$K_S^0 \pi^+$	(1.22 ±0.04) × 10 ⁻³	916	DESIG=23	
$K^+ \eta$	[e] (1.76 ±0.08) × 10 ⁻³	835	DESIG=121	
$K^+ \omega$	[e] (9.9 ±1.5) × 10 ⁻⁴	741	DESIG=129	
$K^+ \eta'(958)$	[e] (2.68 ±0.24) × 10 ⁻³	646	DESIG=122	
$K^+ \pi^+ \pi^-$	(6.23 ±0.10) × 10 ⁻³	900	DESIG=45	
$K^+ \rho^0$	(2.18 ±0.25) × 10 ⁻³	745	DESIG=82	
$K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-$	(7.3 ±1.7) × 10 ⁻⁴	-	DESIG=108	
$K^+ f_0(500), f_0 \rightarrow \pi^+ \pi^-$	(4.5 ±3.0) × 10 ⁻⁴	-	DESIG=207	
$K^+ f_0(980), f_0 \rightarrow \pi^+ \pi^-$	(2.8 ±1.1) × 10 ⁻⁴	-	DESIG=209	
$K^+ f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	(1.2 ±0.6) × 10 ⁻³	-	DESIG=208	
$K^*(892)^0 \pi^+, K^{*0} \rightarrow$	(1.68 ±0.26) × 10 ⁻³	775	DESIG=83	
$K^+ \pi^-$				
$K^*(1410)^0 \pi^+, K^{*0} \rightarrow$	(6 ±4) × 10 ⁻⁴	-	DESIG=109	
$K^+ \pi^-$				
$K^*(1430)^0 \pi^+, K^{*0} \rightarrow$	(9.4 ±3.2) × 10 ⁻⁴	-	DESIG=110	
$K^+ \pi^-$				
$K^+ \pi^+ \pi^-$ nonresonant	(9.9 ±3.2) × 10 ⁻⁴	900	DESIG=111	
$K_S^0 \pi^+ \pi^0$	(5.09 ±0.22) × 10 ⁻³	899	DESIG=242	
$K_S^0 \rho(770)^+, \rho^+ \rightarrow \pi^+ \pi^0$	(2.6 ±0.4) × 10 ⁻³	-	DESIG=237	
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	(1.04 ±0.32) × 10 ⁻³	-	DESIG=238	
$K^*(892)^0 \pi^+, K^{*0} \rightarrow$	(4.3 ±1.2) × 10 ⁻⁴	-	DESIG=239	
$K_S^0 \pi^0$				
$K^*(892)^+ \pi^0, K^{*+} \rightarrow$	(2.3 ±0.7) × 10 ⁻⁴	-	DESIG=240	
$K_S^0 \pi^+$				
$K^*(1410)^0 \pi^+, K^{*0} \rightarrow$	(1.7 ±0.9) × 10 ⁻⁴	-	DESIG=241	
$K_S^0 \pi^0$				
$K_S^0 2\pi^+ \pi^-$	(2.8 ±1.0) × 10 ⁻³	870	DESIG=123	
$K^+ \pi^+ \pi^- \pi^0$	(9.7 ±0.6) × 10 ⁻³	873	DESIG=217	
$K^*(892)^0 \rho^+, K^{*0} \rightarrow$	(3.9 ±0.4) × 10 ⁻³	-	DESIG=218	
$K^+ \pi^-$				
$K^*(892)^+ \rho^0, K^{*+} \rightarrow$	(4.2 ±1.2) × 10 ⁻⁴	-	DESIG=219	
$K^+ \pi^0$				
$K_1(1270)^0 \pi^+, K_1^0 \rightarrow$	(3.9 ±1.3) × 10 ⁻⁴	†	DESIG=220	
$K^+ \rho^-$				
$K_1(1400)^0 \pi^+, K_1^0 \rightarrow$	(5.4 ±0.9) × 10 ⁻⁴	-	DESIG=221	
$K^*(890)^+ \pi^-, K^{*+} \rightarrow$				
$K^+ \pi^0$				
$K_1(1400)^0 \pi^+, K_1^0 \rightarrow$	(5.9 ±1.0) × 10 ⁻⁴	-	DESIG=222	
$K^*(890)^0 \pi^0, K^{*0} \rightarrow$				
$K^+ \pi^-$				
$K^+ a_1(1260)^0, a_1 \rightarrow \rho^+ \pi^-$	(1.8 ±1.1) × 10 ⁻⁴	-	DESIG=223	
$K^+ a_1(1260)^0, a_1 \rightarrow \rho^- \pi^+$	(1.8 ±1.1) × 10 ⁻⁴	-	DESIG=224	
$K^+ \pi^+ \pi^- \pi^0$ nonresonant	(9.2 ±2.4) × 10 ⁻⁴	873	DESIG=225	
$(K^+ \pi^0) P\text{-wave } \rho^0$	(1.01 ±0.21) × 10 ⁻³	688	DESIG=226	
$K^+ \omega \pi^0$	[e] < 8.2 × 10 ⁻³ CL=90%	684	DESIG=130	
$K^+ \omega \pi^+ \pi^-$	[e] < 5.4 × 10 ⁻³ CL=90%	603	DESIG=131	
$K^+ \omega \eta$	[e] < 7.9 × 10 ⁻³ CL=90%	366	DESIG=132	
$2K^+ K^-$	(2.18 ±0.20) × 10 ⁻⁴	628	DESIG=67	
$\phi K^+, \phi \rightarrow K^+ K^-$	(8.9 ±2.0) × 10 ⁻⁵	-	DESIG=154	

Radiative decays

$\rho(770)^+ \gamma$	< 6.1 × 10 ⁻⁴ CL=90%	-	NODE=S034;CLUMP=H DESIG=252
----------------------	---------------------------------	---	--------------------------------

Doubly Cabibbo-suppressed modes

$2K^+ \pi^-$	(1.293 ±0.027) × 10 ⁻⁴ S=1.1	805	NODE=S034;CLUMP=F DESIG=116
$K^+ K^*(892)^0, K^{*0} \rightarrow$	(6.1 ±3.4) × 10 ⁻⁵	-	DESIG=155
$K^+ \pi^-$			
$2K^+ \pi^- \pi^0$	< 1.7 × 10 ⁻⁴ CL=90%	748	DESIG=247

Baryon-antibaryon mode

$p\bar{n}$	(1.22 ± 0.11) × 10 ⁻³	295
$p\bar{p}e^+\nu_e$	< 2.0 × 10 ⁻⁴ CL=90%	296

NODE=S034;CLUMP=G

DESIG=124

DESIG=171

**ΔC = 1 weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

NODE=S034;CLUMP=E

$\pi^+ e^+ e^-$	[j] < 5.5 × 10 ⁻⁶ CL=90%	979	DESIG=93
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[k] (1.17 ± 0.22) × 10 ⁻⁵	-	DESIG=152
$\pi^+ \pi^0 e^+ e^-$	< 7.0 × 10 ⁻⁵ CL=90%	974	DESIG=249
$\rho^+ \phi, \phi \rightarrow e^+ e^-$	(2.4 ± 0.7) × 10 ⁻⁵	-	DESIG=248
$\pi^+ \mu^+ \mu^-$	[j] < 1.8 × 10 ⁻⁷ CL=90%	968	DESIG=73
$K^+ e^+ e^-$	C1 < 3.7 × 10 ⁻⁶ CL=90%	922	DESIG=94
$K^+ \pi^0 e^+ e^-$	< 7.1 × 10 ⁻⁵ CL=90%	917	DESIG=250
$K_S^0 \pi^+ e^+ e^-$	< 8.1 × 10 ⁻⁵ CL=90%	916	DESIG=251
$K^+ \mu^+ \mu^-$	C1 < 1.4 × 10 ⁻⁷ CL=90%	909	DESIG=74
$K^*(892)^+ \mu^+ \mu^-$	C1 < 1.4 × 10 ⁻³ CL=90%	765	DESIG=75
$\pi^+ e^+ \mu^-$	LF < 1.1 × 10 ⁻⁶ CL=90%	976	DESIG=156
$\pi^+ e^- \mu^+$	LF < 9.4 × 10 ⁻⁷ CL=90%	976	DESIG=157
$K^+ e^+ \mu^-$	LF < 7.9 × 10 ⁻⁷ CL=90%	919	DESIG=158
$K^+ e^- \mu^+$	LF < 5.6 × 10 ⁻⁷ CL=90%	919	DESIG=159
$\pi^- 2e^+$	L < 1.4 × 10 ⁻⁶ CL=90%	979	DESIG=97
$\pi^- 2\mu^+$	L < 8.6 × 10 ⁻⁸ CL=90%	968	DESIG=76
$\pi^- e^+ \mu^+$	L < 6.3 × 10 ⁻⁷ CL=90%	976	DESIG=98
$K^- 2e^+$	L < 7.7 × 10 ⁻⁷ CL=90%	922	DESIG=99
$K^- 2\mu^+$	L < 2.6 × 10 ⁻⁸ CL=90%	909	DESIG=77
$K^- e^+ \mu^+$	L < 2.6 × 10 ⁻⁷ CL=90%	919	DESIG=100
$K^*(892)^- 2\mu^+$	L < 1.4 × 10 ⁻³ CL=90%	765	DESIG=78

 $D_s^{*\pm}$

$$I(J^P) = 0(1^-)$$

NODE=S074

Mass $m = 2112.2 \pm 0.4$ MeV

NODE=S074M;DTYPE=M

 $m_{D_s^{*\pm}} - m_{D_s^\pm} = 143.8 \pm 0.4$ MeVFull width $\Gamma < 1.9$ MeV, CL = 90%NODE=S074DM;DTYPE=D
NODE=S074W;DTYPE=G D_s^{*-} modes are charge conjugates of the modes below.

NODE=S074215;NODE=S074

D_s^{*+} DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)	
$D_s^+ \gamma$	(93.6 ± 0.4) %	139	DESIG=1
$D_s^+ \pi^0$	(5.77 ± 0.35) %	48	DESIG=2
$D_s^+ e^+ e^-$	(6.7 ± 1.6) × 10 ⁻³	139	DESIG=3
$e^+ \nu_e$	(2.1 $^{+1.2}_{-0.9}$) × 10 ⁻⁵	1056	DESIG=4

 $D_{s0}^*(2317)^\pm$

$$I(J^P) = 0(0^+)$$

 J, P need confirmation.

NODE=M172

 J^P is natural, low mass consistent with 0^+ .See the review on "Heavy Non- $q\bar{q}$ Mesons."Mass $m = 2317.8 \pm 0.5$ MeV

NODE=M172M;DTYPE=M

 $m_{D_{s0}^*(2317)^\pm} - m_{D_s^\pm} = 349.4 \pm 0.5$ MeV

NODE=M172DM;DTYPE=D

Full width $\Gamma < 3.8$ MeV, CL = 95%

NODE=M172W;DTYPE=G

$D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

NODE=M172215;NODE=M172

$D_{s0}^*(2317)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$D_s^+ \pi^0$	$(100^{+0}_{-20})\%$		298
$D_s^+ \gamma$	$< 5\%$	90%	323
$D_s^*(2112)^+ \gamma$	$< 6\%$	90%	—
$D_s^+ \gamma \gamma$	$< 18\%$	95%	323
$D_s^*(2112)^+ \pi^0$	$< 11\%$	90%	—
$D_s^+ \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%	194
$D_s^+ \pi^0 \pi^0$	not seen		205

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7;OUR EVAL;→ UNCHECKED ←

$D_{s1}(2460)^\pm$

$$I(J^P) = 0(1^+)$$

NODE=M173

See the review on "Heavy Non- $q\bar{q}$ Mesons."

Mass $m = 2459.5 \pm 0.6$ MeV ($S = 1.1$)

$m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}} = 347.3 \pm 0.7$ MeV ($S = 1.2$)

$m_{D_{s1}(2460)^\pm} - m_{D^\pm} = 491.1 \pm 0.6$ MeV ($S = 1.1$)

Full width $\Gamma < 3.5$ MeV, CL = 95%

NODE=M173M;DTYPE=M

NODE=M173MD;DTYPE=D

NODE=M173DM;DTYPE=D

NODE=M173W;DTYPE=G

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

NODE=M173215;NODE=M173

$D_{s1}(2460)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$D_s^{*+} \pi^0$	$(48 \pm 11)\%$		297
$D_s^+ \gamma$	$(18 \pm 4)\%$		442
$D_s^+ \pi^+ \pi^-$	$(4.3 \pm 1.3)\%$	$S=1.1$	363
$D_s^{*+} \gamma$	$< 8\%$	CL=90%	323
$D_{s0}^*(2317)^+ \gamma$	$(3.7^{+5.0}_{-2.4})\%$		138

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

$D_{s1}(2536)^\pm$

$$I(J^P) = 0(1^+)$$

J, P need confirmation.

NODE=M121

Mass $m = 2535.11 \pm 0.06$ MeV

NODE=M121M;DTYPE=M

$m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)} = 422.9 \pm 0.4$ MeV

NODE=M121DM;DTYPE=D

$m_{D_{s1}(2536)^\pm} - m_{D^*(2010)^\pm} = 524.85 \pm 0.04$ MeV

NODE=M121DN;DTYPE=D

$m_{D_{s1}(2536)^\pm} - m_{D^*(2007)^0} = 528.26 \pm 0.05$ MeV ($S = 1.1$)

NODE=M121DP;DTYPE=D

Full width $\Gamma = 0.92 \pm 0.05$ MeV

NODE=M121W;DTYPE=G

Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M121215;NODE=M121

$D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

$D_{s1}(2536)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$D^*(2010)^+ K^0$	$(31 \pm 7)\%$		149
$(D^*(2010)^+ K^0)_{S-wave}$	$(22 \pm 5)\%$		149
$K_S^0 D^*(2010)^+$	$(17 \pm 4)\%$		149
$D_s^+ \pi^- K^+$	$(10.0 \pm 2.5) \times 10^{-3}$		176
$D^*(2007)^0 K^+$	$(36 \pm 6)\%$		167
$D^+ K^0$	$< 12\%$	90%	381
$D^0 K^+$	$< 4\%$	90%	391
$D_s^{*+} \gamma$	possibly seen		388
$D_s^+ \pi^+ \pi^-$	seen		437

DESIG=1

DESIG=7

DESIG=10

DESIG=8

DESIG=4

DESIG=2

DESIG=5

DESIG=3

DESIG=6

$D_{s2}^*(2573)$

$$I(J^P) = 0(2^+)$$

NODE=M148

Mass $m = 2569.1 \pm 0.8$ MeV ($S = 2.4$)

NODE=M148M;DTYPE=M

$m_{D_{s2}^*(2573)} - m_{D^0} = 704 \pm 3.2$ MeV

NODE=M148DM;DTYPE=D

Full width $\Gamma = 16.9 \pm 0.7$ MeV

NODE=M148W;DTYPE=G

$D_{s2}^*(2573)^-$ modes are charge conjugates of the modes below.

NODE=M148215;NODE=M148

$D_{s2}^*(2573)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	(37±6) %	431
$D^*(2007)^0 K^+$	not seen	238
$D^+ K_S^0$	seen	422
$D^{*+} K_S^0$	seen	225

DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=5;OUR EVAL;→ UNCHECKED ←

$D_{s1}^*(2700)^\pm$

$$I(J^P) = 0(1^-)$$

Mass $m = 2714 \pm 5$ MeV ($S = 1.5$)

Full width $\Gamma = 122 \pm 10$ MeV

NODE=M182

NODE=M182M;DTYPE=M

NODE=M182W;DTYPE=G

$D_{s1}^*(2700)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	seen	579
$D^+ K_S^0$	seen	573
$D^{*0} K^+$	seen	438
$D^{*+} K_S^0$	seen	431

NODE=M182215;DESIG=1;OUR EVAL;
→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=6;OUR EVAL;→ UNCHECKED ←

$D_{s3}^*(2860)^\pm$

$$I(J^P) = 0(3^-)$$

Mass $m = 2860 \pm 7$ MeV

Full width $\Gamma = 53 \pm 10$ MeV

NODE=M226

NODE=M226M;DTYPE=M

NODE=M226W;DTYPE=G

$D_{s3}^*(2860)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	seen	710
$D^+ K_S^0$	seen	704
$D^{*0} K^+$	seen	589
$D^{*+} K_S^0$	seen	584

NODE=M226215;DESIG=2;OUR EVAL;
→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=6;OUR EVAL;→ UNCHECKED ←

NOTES

- [a] See the Particle Listings for the (complicated) definition of this quantity. LINKAGE=DEF
- [b] This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+ exclusive fractions — an $e^+ \nu_e$ with an η , η' , ϕ , K^0 , or K^{*0} — is 5.99 ± 0.31 % . LINKAGE=SLE
- [c] This fraction includes η from η' decays. LINKAGE=EFR
- [d] The sum of our exclusive η' fractions — $\eta' e^+ \nu_e$, $\eta' \mu^+ \nu_\mu$, $\eta' \pi^+$, $\eta' \rho^+$, and $\eta' K^+$ — is 11.8 ± 1.6 %. LINKAGE=INC
- [e] This branching fraction includes all the decay modes of the final-state resonance. LINKAGE=DFR
- [f] A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and ω - ϕ mixing is an unlikely explanation for any fraction above about 2×10^{-4} . LINKAGE=MAR
- [g] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings. LINKAGE=SDQ
- [h] We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491. LINKAGE=DBF
- [i] This is the average of a model-independent and a K -matrix parametrization of the $\pi^+ \pi^-$ S -wave and is a sum over several f_0 mesons. LINKAGE=KMP
- [j] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay. LINKAGE=FIX
- [k] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state. LINKAGE=NTC