b' (4^{th} Generation) Quark, Searches for

NODE=Q008

$b'(-1/3)$-quark/hadron mass limits in $p\overline{p}$ and pp collisions

NODE=Q008BPP
NODE=Q008BPP;CHECK LIMITS

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	
>1530	95	[1] AAD 24BP	ATLS	$B(b' \to W u) = 1$	
>1540	95	[2] HAYRAPETY... 24AQ	CMS	$B(b' \to Z b) = 1$	
>1570	95	[2] HAYRAPETY... 24AQ	CMS	$B(b' \to H b) = 1$	OCCUR=2
>1560	95	[3] TUMASYAN 23V	CMS	$B(b' \to W t) = 1$	
>1570	95	[4] SIRUNYAN 20BI	CMS	$B(b' \to H b) = 1$	
>1000	95	[5] AABOUD 18CE	ATLS	$\geq 2\ell + \not{E}_T + \geq 1bj$	
> 950	95	[6] AABOUD 18CL	ATLS	$W t$, $Z b$, $h b$ modes	
>1010	95	[7,8] AABOUD 18CP	ATLS	2,3ℓ, singlet model	
>1140	95	[6,9] AABOUD 18CP	ATLS	2,3ℓ, doublet model	OCCUR=2
>1220	95	[10,11] AABOUD 18CR	ATLS	singlet b'. ATLAS Combination	
>1370	95	[10,12] AABOUD 18CR	ATLS	b' in a weak isospin doublet (t',b'). ATLAS combination.	OCCUR=2
> 730	95	[13] SIRUNYAN 17AU	CMS		
> 810	95	[14] AAD 15Z	ATLS		
> 190	95	[15] ABAZOV 08X	D0	$c\tau$= 200mm	
> 190	95	[16] ACOSTA 03	CDF	quasi-stable b'	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>1460	95	[17] AAD 23AG	ATLS	$B(b' \to W t) = 1$	
>1420	95	[18] AAD 23AV	ATLS	$B(b' \to Z b) = 1$	
>1390	95	[4] SIRUNYAN 20BI	CMS	$B(b' \to Z b) = 1$	OCCUR=2
>1130	95	[19] SIRUNYAN 19AQ	CMS	$B(b' \to Z b) = 1$	
>1230	95	[20] SIRUNYAN 19BW	CMS	$B(b' \to W t) = 1$	
>1350	95	[21] AABOUD 18AW	ATLS	$B(b' \to W t) = 1$	
> 910	95	[22] SIRUNYAN 18BM	CMS	$W t$, $Z b$, $h b$ modes	
> 845	95	[23] SIRUNYAN 18Q	CMS	$B(b' \to W u) = 1$	
> 880	95	[24] KHACHATRY... 16AN	CMS	$B(b' \to W t) = 1$	
<350, 580–635, >700	95	[25] AAD 15AR	ATLS	$B(b' \to H b) = 1$	
> 620	95	[26] AAD 15BY	ATLS	$W t$, $Z b$, $h b$ modes	
> 730	95	[27] AAD 15BY	ATLS	$B(b' \to W t) = 1$	OCCUR=2
> 690	95	[28] AAD 15CN	ATLS	$B(b' \to W q) = 1$ (q=u)	
> 755	95	[29] AAD 14AZ	ATLS	$B(b' \to W t) = 1$	
> 675	95	[30] CHATRCHYAN 13I	CMS	$B(b' \to W t) = 1$	
> 480	95	[31] AAD 12AT	ATLS	$B(b' \to W t) = 1$	
> 400	95	[32] AAD 12AU	ATLS	$B(b' \to Z b) = 1$	
> 350	95	[33] AAD 12BC	ATLS	$B(b' \to W q) = 1$ (q=u,c)	
> 450	95	[34] AAD 12BE	ATLS	$B(b' \to W t) = 1$	
> 685	95	[35] CHATRCHYAN 12BH	CMS	$m_{t'} = m_{b'}$	
> 611	95	[36] CHATRCHYAN 12X	CMS	$B(b' \to W t) = 1$	
> 372	95	[37] AALTONEN 11J	CDF	$b' \to W t$	
> 361	95	[38] CHATRCHYAN 11L	CMS	Repl. by CHATRCHYAN 12X	
> 338	95	[39] AALTONEN 10H	CDF	$b' \to W t$	
> 380–430	95	[40] FLACCO 10	RVUE	$m_{b'} > m_{t'}$	
> 268	95	[41,42] AALTONEN 07C	CDF	$B(b' \to Zb) = 1$	
> 199	95	[43] AFFOLDER 00	CDF	NC: $b' \to Z b$	
> 148	95	[44] ABE 98N	CDF	NC: $b' \to Z b$ + vertex	
> 96	95	[45] ABACHI 97D	D0	NC: $b' \to b\gamma$	
> 128	95	[46] ABACHI 95F	D0	$\ell\ell$ + jets, ℓ + jets	
> 75	95	[47] MUKHOPAD... 93	RVUE	NC: $b' \to b\ell\ell$	
> 85	95	[48] ABE 92	CDF	CC: $\ell\ell$	
> 72	95	[49] ABE 90B	CDF	CC: $e + \mu$	
> 54	95	[50] AKESSON 90	UA2	CC: e + jets + $\not{E}_T$	
> 43	95	[51] ALBAJAR 90B	UA1	CC: μ + jets	
> 34	95	[52] ALBAJAR 88	UA1	CC: e or μ + jets	

[1] AAD 24BP based on 140 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Limit on pair-production of heavy vectorlike quarks where each decays into a W boson and a light quark. Used events with ℓ + $\not{E}_T$ + multiple jets + 1 boosted-W-jet. NODE=Q008BPP;LINKAGE=JA

[2] HAYRAPETYAN 24AQ based on 138 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like b' is searched for in the fully hadronic final states and those containing $\ell^+ \ell^-$ from a Z boson. The data are consistent with the SM background predictions and limits are obtained for different branching ratios B($b' \rightarrow Hb$), B($b' \rightarrow Zb$), B($b' \rightarrow Wt$). NODE=Q008BPP;LINKAGE=KA

[3] TUMASYAN 23V based on 138 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like b' is seached for in the single-lepton, same-sign charge dilepton and multi-lepton channels. The data are consistent with the SM background predictions and limits are obtained for different branching ratios. NODE=Q008BPP;LINKAGE=EA

[4] SIRUNYAN 20BI based on 137 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like b' is seached for with each b' decaying into Zb or hb. Analysis focuses on final states consisting of jets from six quarks. Mass limits are obtained for a variety of branching ratios of b' decays. NODE=Q008BPP;LINKAGE=DA

[5] AABOUD 18CE based on 36.1 fb^{-1} of proton-proton data taken at $\sqrt{s}$ = 13 TeV. Events including a same-sign lepton pair are used. The limit is for a singlet model, assuming the branching ratios of b' into Zb, Wt and Hb as predicted by the model. NODE=Q008BPP;LINKAGE=Y

[6] AABOUD 18CL, AABOUD 18CP based on 36.1 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. The limit is for the pair-produced vector-like b' using all-hadronic final state. The analysis is particularly powerful for the $b' \rightarrow hb$ mode. Assuming the pure decay only in this mode sets a limit $m_{b'} > 1010$ GeV. NODE=Q008BPP;LINKAGE=Q

[7] AABOUD 18CP based on 36.1 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair and single production of vector-like b' are seached for with at least one b' decaying into Zb. In the case of B($b' \rightarrow Zb$) = 1, the limit is $m_{b'} > 1220$ GeV. NODE=Q008BPP;LINKAGE=R

[8] The limit is for the singlet model, assuming that the branching ratios into Wt, Zb, hb add up to one. NODE=Q008BPP;LINKAGE=T

[9] The limit is for the doublet model, assuming that the branching ratios into Wt, Zb, hb add up to one. NODE=Q008BPP;LINKAGE=S

[10] AABOUD 18CR based on 36.1 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. A combination of searches for the pair-produced vector-like b' in various decay channels ($b' \rightarrow Wt$, Zb, hb). Also a model-independent limit is obtained as $m_{b'} > 1.03$ TeV, assuming that the branching ratios into Zb, Wt, and hb add up to one. NODE=Q008BPP;LINKAGE=V

[11] The limit is for the singlet b'. NODE=Q008BPP;LINKAGE=W

[12] The limit is for b' in a weak isospin doublet (t',b') and $|V_{t'b}| \ll |V_{tb'}|$. For a b' in a doublet with a charge $-4/3$ vector-like quark, the limit $m_{b'} > 1.14$ TeV is obtained. NODE=Q008BPP;LINKAGE=X

[13] SIRUNYAN 17AU based on 2.3–2.6 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Limit on pair-produced singlet vector-like b' using one lepton and several jets. The mass bound is given for a b' transforming as a singlet under the electroweak symmetry group, assumed to decay through W, Z or Higgs boson (which decays to jets) and to a third generation quark. NODE=Q008BPP;LINKAGE=M

[14] AAD 15Z based on 20.3 fb^{-1} of pp data at $\sqrt{s}$ = 8 TeV. Used events with ℓ + $\not{E}_T$ + $\geq$ 6j ($\geq$ 1 b) and at least one pair of jets from weak boson decay, primarily designed to select the signature $b' \overline{b}' \rightarrow WWt\overline{t} \rightarrow WWWWb\overline{b}$. This is a limit on pair-produced vector-like b'. The lower mass limit is 640 GeV for a vector-like singlet b'. NODE=Q008BPP;LINKAGE=K

[15] Result is based on 1.1 fb^{-1} of data. No signal is found for the search of long-lived particles which decay into final states with two electrons or photons, and upper bound on the cross section times branching fraction is obtained for $2 < c\tau < 7000$ mm; see Fig. 3. 95% CL excluded region of b' lifetime and mass is shown in Fig. 4. NODE=Q008BPP;LINKAGE=AA

[16] ACOSTA 03 looked for long-lived fourth generation quarks in the data sample of 90 pb^{-1} of $\sqrt{s}$=1.8 TeV $p\overline{p}$ collisions by using the muon-like penetration and anomalously high ionization energy loss signature. The corresponding lower mass bound for the charge (2/3)e quark (t') is 220 GeV. The t' bound is higher than the b' bound because t' is more likely to produce charged hadrons than b'. The 95% CL upper bounds for the production cross sections are given in their Fig. 3. NODE=Q008BPP;LINKAGE=CS

[17] AAD 23AG based on 139 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like top or bs is searched for in the mode 1ℓ + $\geq$ 4j($\geq$ 1b-tagged) + $\not{E}_T$. The data are consistent with the SM background predictions and limits are obtained for different branching ratios. Masses below 1.59 TeV are excluded assuming a mass-degenerate vector-like doublet (t',b') model. NODE=Q008BPP;LINKAGE=FA

[18] AAD 23AV based on 139 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like b' is searched for in the mode $\ell^{\pm}\ell^{\mp}$ + $\geq$ 2j ($\geq$ 1b-tagged) + $\not{E}_T$ or with 3ℓ. The data are consistent with the SM background predictions and limits are obtained for different branching ratios. NODE=Q008BPP;LINKAGE=IA

[19] SIRUNYAN 19AQ based on 35.9 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Pair production of vector-like b' is seached for with one b' decaying into Zb and the other b' decaying into Wt, Zb, hb. Events with an opposite-sign lepton pair consistent with coming from Z and jets are used. Mass limits are obtained for a variety of branching ratios of b'. NODE=Q008BPP;LINKAGE=Z

[20] SIRUNYAN 19BW based on 35.9 fb^{-1} of pp data at $\sqrt{s}=13$ TeV. The limit is for the pair-produced vector-like b' using all-hadronic final state. The analysis is made for the Zb, Wt, hb modes and mass limits are obtained for a variety of branching ratios. NODE=Q008BPP;LINKAGE=BA

[21] AABOUD 18AW based on 36.1 fb^{-1} of pp data at $\sqrt{s}=13$ TeV. The limit is for the pair-produced vector-like b' using lepton-plus-jets final state. The search is also sensitive to the decays into Zb and Hb final states. NODE=Q008BPP;LINKAGE=O

[22] SIRUNYAN 18BM based on 35.9 fb^{-1} of pp data at $\sqrt{s}=13$ TeV. The limit is for the pair-produced vector-like b'. Three channels (single lepton, same-charge 2 leptons, or at least 3 leptons) are considered for various branching fraction combinations. Assuming B(tW) = 1, the limit is 1240 GeV and for B(bZ) = 1 it is 960 GeV. NODE=Q008BPP;LINKAGE=N

[23] SIRUNYAN 18Q based on 19.7 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. The limit is for the pair-produced vector-like b' that couple only to light quarks. Upper cross section limits on the single production of a b' and constraints for other decay channels (Zq and Hq) are also given in the paper. NODE=Q008BPP;LINKAGE=P

[24] KHACHATRYAN 16AN based on 19.7 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. Limit on pair-produced vector-like b' using 1, 2, and >2 leptons as well as fully hadronic final states. Other limits depending on the branching fractions to tW, bZ, and bH are given in Table IX. NODE=Q008BPP;LINKAGE=L

[25] AAD 15AR based on 20.3 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. Used lepton-plus-jets final state. See Fig. 24 for mass limits in the plane of B($b' \to Wt$) vs. B($b' \to Hb$) from $b'\overline{b}' \to Hb + X$ searches. NODE=Q008BPP;LINKAGE=G

[26] AAD 15BY based on 20.3 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. Limit on pair-produced vector-like b' assuming the branching fractions to W, Z, and h modes of the singlet model. Used events containing $\geq 2\ell + \not\!\!E_T + \geq 2$j ($\geq 1$ b) and including a same-sign lepton pair. NODE=Q008BPP;LINKAGE=H

[27] AAD 15BY based on 20.3 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. Limit on pair-produced chiral b'-quark. Used events containing $\geq 2\ell + \not\!\!E_T + \geq 2$j ($\geq 1$ b) and including a same-sign lepton pair. NODE=Q008BPP;LINKAGE=I

[28] AAD 15CN based on 20.3 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. Limit on pair-production of chiral b'-quark. Used events with $\ell + \not\!\!E_T + \geq 4$j (non-$b$-tagged). Limits on a heavy vector-like quark, which decays into Wq, Zq, hq, are presented in the plane B($Q \to Wq$) vs. B($Q \to hq$) in Fig. 12. NODE=Q008BPP;LINKAGE=J

[29] Based on 20.3 fb^{-1} of pp data at $\sqrt{s}=8$ TeV. No significant excess over SM expectation is found in the search for pair production or single production of b' in the events with dilepton from a high p_T Z and additional jets ($\geq$ 1 b-tag). If instead of B($b' \to Wt$) = 1 an electroweak singlet with B($b' \to Wt$) $\sim$ 0.45 is assumed, the limit reduces to 685 GeV. NODE=Q008BPP;LINKAGE=E

[30] Based on 5.0 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. CHATRCHYAN 13I looked for events with one isolated electron or muon, large $\not\!\!E_T$, and at least four jets with large transverse momenta, where one jet is likely to originate from the decay of a bottom quark. NODE=Q008BPP;LINKAGE=B

[31] Based on 1.04 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. No signal is found for the search of heavy quark pair production that decay into W and a t quark in the events with a high p_T isolated lepton, large $\not\!\!E_T$, and at least 6 jets in which one, two or more dijets are from W. NODE=Q008BPP;LINKAGE=GD

[32] Based on 2.0 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. No $b' \to Zb$ invariant mass peak is found in the search of heavy quark pair production that decay into Z and a b quark in events with $Z \to e^+e^-$ and at least one b-jet. The lower mass limit is 358 GeV for a vector-like singlet b' mixing solely with the third SM generation. NODE=Q008BPP;LINKAGE=DG

[33] Based on 1.04 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. No signal is found for the search of heavy quark pair production that decay into W and a quark in the events with dileptons, large $\not\!\!E_T$, and ≥ 2 jets. NODE=Q008BPP;LINKAGE=GA

[34] Based on 1.04 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. AAD 12BE looked for events with two isolated like-sign leptons and at least 2 jets, large $\not\!\!E_T$ and $H_T > 350$ GeV. NODE=Q008BPP;LINKAGE=AD

[35] Based on 5 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. CHATRCHYAN 12BH searched for QCD and EW production of single and pair of degenerate 4'th generation quarks that decay to bW or tW. Absence of signal in events with one lepton, same-sign dileptons or tri-leptons gives the bound. With a mass difference of 25 GeV/c^2 between $m_{t'}$ and $m_{b'}$, the corresponding limit shifts by about ± 20 GeV/c^2. NODE=Q008BPP;LINKAGE=CT

[36] Based on 4.9 fb^{-1} of pp data at $\sqrt{s}=7$ TeV. CHATRCHYAN 12X looked for events with trileptons or same-sign dileptons and at least one b jet. NODE=Q008BPP;LINKAGE=CA

[37] Based on 4.8 fb^{-1} of data in $p\overline{p}$ collisions at 1.96 TeV. AALTONEN 11J looked for events with $\ell + \not\!\!E_T + \geq 5$j ($\geq 1$ b or c). No signal is observed and the bound $\sigma(b'\overline{b}')$ < 30 fb for $m_{b'} > 375$ GeV is found for B($b' \to Wt$) = 1. NODE=Q008BPP;LINKAGE=AO

[38] Based on 34 pb^{-1} of data in pp collisions at 7 TeV. CHATRCHYAN 11L looked for multi-jet events with trileptons or same-sign dileptons. No excess above the SM background excludes $m_{b'}$ between 255 and 361 GeV at 95% CL for B($b' \to Wt$) = 1. NODE=Q008BPP;LINKAGE=CH

[39] Based on 2.7 fb^{-1} of data in $p\overline{p}$ collisions at $\sqrt{s}=1.96$ TeV. AALTONEN 10H looked for pair production of heavy quarks which decay into tW^- or tW^+, in events with same sign dileptons (e or μ), several jets and large missing E_T. The result is obtained for b' which decays into tW^-. For the charge 5/3 quark ($T_{5/3}$) which decays into tW^+, NODE=Q008BPP;LINKAGE=AT

$m_{T_{5/3}} > 365$ GeV (95% CL) is found when it has the charge $-1/3$ partner B of the same mass.

[40] FLACCO 10 result is obtained from AALTONEN 10H result of $m_{b'} > 338$ GeV, by relaxing the condition B($b' \rightarrow W t$) = 100% when $m_{b'} > m_{t'}$. NODE=Q008BPP;LINKAGE=FL

[41] Result is based on 1.06 fb^{-1} of data. No excess from the SM Z+jet events is found when Z decays into $e e$ or $\mu\mu$. The $m_{b'}$ bound is found by comparing the resulting upper bound on $\sigma(b'\overline{b}')$ [1-(1-B($b' \rightarrow Z b$))^2] and the LO estimate of the b' pair production cross section shown in Fig. 38 of the article. NODE=Q008BPP;LINKAGE=AL

[42] HUANG 08 reexamined the b' mass lower bound of 268 GeV obtained in AALTONEN 07C that assumes B($b' \rightarrow Z b$) = 1, which does not hold for $m_{b'} > 255$ GeV. The lower mass bound is given in the plane of $\sin^2(\theta_{t b'})$ and $m_{b'}$. NODE=Q008BPP;LINKAGE=HU

[43] AFFOLDER 00 looked for b' that decays in to b+Z. The signal searched for is $b b Z Z$ events where one Z decays into $e^+ e^-$ or $\mu^+ \mu^-$ and the other Z decays hadronically. The bound assumes B($b' \rightarrow Z b$)= 100%. Between 100 GeV and 199 GeV, the 95%CL upper bound on $\sigma(b' \rightarrow \overline{b}')\times$B^2($b' \rightarrow Z b$) is also given (see their Fig. 2). NODE=Q008BPP;LINKAGE=EB

[44] ABE 98N looked for $Z \rightarrow e^+ e^-$ decays with displaced vertices. Quoted limit assumes B($b' \rightarrow Z b$)=1 and $c\tau_{b'}$=1 cm. The limit is lower than m_Z+m_b ($\sim$ 96 GeV) if $c\tau >$ 22 cm or $c\tau <$ 0.009 cm. See their Fig. 4. NODE=Q008BPP;LINKAGE=AN

[45] ABACHI 97D searched for b' that decays mainly via FCNC. They obtained 95%CL upper bounds on B($b'\overline{b}' \rightarrow \gamma+$ 3 jets) and B($b'\overline{b}' \rightarrow 2\gamma+$ 2 jets), which can be interpreted as the lower mass bound $m_{b'} > m_Z + m_b$. NODE=Q008BPP;LINKAGE=K2

[46] ABACHI 95F bound on the top-quark also applies to b' and t' quarks that decay predominantly into W. See FROGGATT 97. NODE=Q008BPP;LINKAGE=1K

[47] MUKHOPADHYAYA 93 analyze CDF dilepton data of ABE 92G in terms of a new quark decaying via flavor-changing neutral current. The above limit assumes B($b' \rightarrow b\ell^+\ell^-$)=1%. For an exotic quark decaying only via virtual Z [B($b\ell^+\ell^-$) = 3%], the limit is 85 GeV. NODE=Q008BPP;LINKAGE=C

[48] ABE 92 dilepton analysis limit of >85 GeV at CL=95% also applies to b' quarks, as discussed in ABE 90B. NODE=Q008BPP;LINKAGE=U

[49] ABE 90B exclude the region 28–72 GeV. NODE=Q008BPP;LINKAGE=AB

[50] AKESSON 90 searched for events having an electron with $p_T > 12$ GeV, missing momentum > 15 GeV, and a jet with $E_T > 10$ GeV, $|\eta| < 2.2$, and excluded $m_{b'}$ between 30 and 69 GeV. NODE=Q008BPP;LINKAGE=F

[51] For the reduction of the limit due to non-charged-current decay modes, see Fig. 19 of ALBAJAR 90B. NODE=Q008BPP;LINKAGE=A

[52] ALBAJAR 88 study events at $E_{\rm cm}$ = 546 and 630 GeV with a muon or isolated electron, accompanied by one or more jets and find agreement with Monte Carlo predictions for the production of charm and bottom, without the need for a new quark. The lower mass limit is obtained by using a conservative estimate for the $b'\overline{b}'$ production cross section and by assuming that it cannot be produced in W decays. The value quoted here is revised using the full $O(\alpha_s^3)$ cross section of ALTARELLI 88. NODE=Q008BPP;LINKAGE=D

$b'(-1/3)$ mass limits from single production in $p\overline{p}$ and $p p$ collisions

NODE=Q008BPS
NODE=Q008BPS

VALUE (GeV)	*CL%*	*DOCUMENT ID*	*TECN*	*COMMENT*	
>3000	95	[1] TUMASYAN 22O	CMS	$g b \rightarrow b' \rightarrow t W$, B($b' \rightarrow t W$)=1	
> 693	95	[2] ABAZOV 11F	D0	$q u \rightarrow q' b' \rightarrow q'(W u)$ $\widetilde{\kappa}_{u b'}$=1, B($b' \rightarrow W u$)=1	
> 430	95	[2] ABAZOV 11F	D0	$q d \rightarrow q b' \rightarrow q(Z d)$ $\widetilde{\kappa}_{d b'}=\sqrt{2}$, B($b' \rightarrow Z d$)=1	OCCUR=2
• • • We do not use the following data for averages, fits, limits, etc. • • •					
		[3] AAD 23CQ	ATLS	$b' \rightarrow b h$ ($h \rightarrow b\overline{b}$)	
>2600	95	[4] SIRUNYAN 21AG	CMS	$g b \rightarrow b' \rightarrow t W$, B($b' \rightarrow t W$)=1	
		[5] SIRUNYAN 19AI	CMS	$b Z / t W \rightarrow b' \rightarrow t W$	
>1500	95	[6] AAD 16AH	ATLS	$g b \rightarrow b' \rightarrow t W$, B($b' \rightarrow t W$)=1	
>1390	95	[7] KHACHATRY...16I	CMS	$g b \rightarrow b'_L \rightarrow t W$, B($b'_L \rightarrow t W$)=1	
>1430	95	[8] KHACHATRY...16I	CMS	$g b \rightarrow b'_R \rightarrow t W$, B($b'_R \rightarrow t W$)=1	OCCUR=2
>1530	95	[9] KHACHATRY...16I	CMS	$g b \rightarrow b' \rightarrow t W$, B($b' \rightarrow t W$)=1	OCCUR=3

[1] TUMASYAN 22O based on 138 fb^{-1} of data in pp collisions at 13 TeV. No significant excess over SM expectation is found in the search for a left-handed b' assuming 100% decay to tW using a t-tagged jet and a lepton from W. The model assumes that the b' has the excited quark couplings. The bound is from a statistical combination with an earlier analysis by SIRUNYAN 21AG. The 95% CL bounds are also set as 3.0, 3.0, and 3.2 TeV, respectively, for left-handed, right-handed, and vector-like couplings. NODE=Q008BPS;LINKAGE=F

[2] ABAZOV 11F based on 5.4 fb^{-1} of data in ppbar collisions at 1.96 TeV. ABAZOV 11F looked for single production of b' via the W or Z coupling to the first generation up or down quarks, respectively. Model independent cross section limits for the single production processes $p\overline{p} \rightarrow b' q \rightarrow W u q$, and $p\overline{p} \rightarrow b' q \rightarrow Z d q$ are given in Figs. 3 and 4, respectively, and the mass limits are obtained for the model of ATRE 09 with degenerate bi-doublets of vector-like quarks. NODE=Q008BPS;LINKAGE=AB

[3] AAD 23CQ based on 139 fb^{-1} of data in pp collisions at 13 TeV. No significant excess over SM expectation is found. Limits on mass and production cross section of a vector-like b' are obtained in several theoretical scenarios determined by the couplings betwen b' and W, Z, h. NODE=Q008BPS;LINKAGE=H

[4] SIRUNYAN 21AG based on 137 fb^{-1} of data in pp collisions at 13 TeV. No significant excess over SM expectation is found in the search for a left-handed b' assuming 100% decay to tW using all hadronic final states, where t and W are tagged as single jets, respectively. The model assumes that the b' has the excited quark couplings. The 95% CL bounds are also set as 2.8 and 3.1 TeV, respectively, for the right-handed and vector-like couplings. NODE=Q008BPS;LINKAGE=G

[5] SIRUNYAN 19AI based on 35.9 fb^{-1} of pp data at $\sqrt{s}$ = 13 TeV. Exclusion limits are set on the product of the production cross section and branching fraction for the $b'(-1/3) +\ b$ and $b'(-1/3) +\ t$ modes as a function of the vector-like quark mass in Figs. 7 and 8 and in Tab. 2 for relative vector-like quark widths between 1 and 30% for left- and right-handed vector-like quark couplings. No significant deviation from the SM prediction is observed. NODE=Q008BPS;LINKAGE=E

[6] AAD 16AH based on 20.3 fb^{-1} of data in pp collisions at 8 TeV. No significant excess over SM expectation is found in the search for a vector-like b' in the single-lepton and dilepton channels (ℓ or $\ell\ell$) + 1,2,3 j ($\geq 1b$). The model assumes that the b' has the excited quark couplings. NODE=Q008BPS;LINKAGE=D

[7] Based on 19.7 fb^{-1} of data in pp collisions at 8 TeV. Limit on left-handed b' assuming 100% decay to tW and using all-hadronic, lepton + jets, and dilepton final states. NODE=Q008BPS;LINKAGE=A

[8] Based on 19.7 fb^{-1} of data in pp collisions at 8 TeV. Limit on right-handed b' assuming 100% decay to tW and using all-hadronic, lepton + jets, and dilepton final states. NODE=Q008BPS;LINKAGE=B

[9] Based on 19.7 fb^{-1} of data in pp collisions at 8 TeV. Limit on vector-like b' assuming 100% decay to tW and using all-hadronic, lepton+jets, and dilepton final states. NODE=Q008BPS;LINKAGE=C

MASS LIMITS for b' (4th Generation) Quark or Hadron in $e^+ e^-$ Collisions

NODE=Q008BPE

Search for hadrons containing a fourth-generation $-1/3$ quark denoted b'. NODE=Q008BPE

The last column specifies the assumption for the decay mode (CC denotes the conventional charged-current decay) and the event signature which is looked for.

NODE=Q008BPE

VALUE (GeV)	*CL%*	*DOCUMENT ID*		*TECN*	*COMMENT*	
>46.0	95	[1] DECAMP	90F	ALEP	any decay	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
none 96–103	95	[2] ABDALLAH	07	DLPH	$b' \rightarrow bZ$, cW	
		[3] ADRIANI	93G	L3	Quarkonium	
>44.7	95	ADRIANI	93M	L3	$\Gamma(Z)$	
>45	95	ABREU	91F	DLPH	$\Gamma(Z)$	
none 19.4–28.2	95	ABE	90D	VNS	Any decay; event shape	
>45.0	95	ABREU	90D	DLPH	B(CC) = 1; event shape	
>44.5	95	[4] ABREU	90D	DLPH	$b' \rightarrow cH^-$, $H^- \rightarrow \overline{c}s$, $\tau^-\nu$	OCCUR=2
>40.5	95	[5] ABREU	90D	DLPH	$\Gamma(Z \rightarrow$ hadrons)	OCCUR=3
>28.3	95	ADACHI	90	TOPZ	B(FCNC)=100%; isol. γ or 4 jets	
>41.4	95	[6] AKRAWY	90B	OPAL	Any decay; acoplanarity	
>45.2	95	[6] AKRAWY	90B	OPAL	B(CC) = 1; acoplanarity	OCCUR=2
>46	95	[7] AKRAWY	90J	OPAL	$b' \rightarrow \gamma$ + any	
>27.5	95	[8] ABE	89E	VNS	B(CC) =1; μ, e	
none 11.4–27.3	95	[9] ABE	89G	VNS	B($b' \rightarrow b\gamma$) > 10%; isolated γ	
>44.7	95	[10] ABRAMS	89C	MRK2	B(CC)= 100%; isol. track	
>42.7	95	[10] ABRAMS	89C	MRK2	B(bg)= 100%; event shape	OCCUR=2

>42.0	95	[10] ABRAMS	89C	MRK2	Any decay; event shape	OCCUR=3
>28.4	95	[11,12] ADACHI	89C	TOPZ	B($C C$) =1; μ	
>28.8	95	[13] ENO	89	AMY	B($C C$) $\gtrsim$ 90%; μ, e	
>27.2	95	[13,14] ENO	89	AMY	any decay; event shape	OCCUR=2
>29.0	95	[13] ENO	89	AMY	B($b' \rightarrow\ b g$) $\gtrsim$ 85%; event shape	OCCUR=3
>24.4	95	[15] IGARASHI	88	AMY	μ,e	
>23.8	95	[16] SAGAWA	88	AMY	event shape	
>22.7	95	[17] ADEVA	86	MRKJ	μ	
>21		[18] ALTHOFF	84C	TASS	R, event shape	
>19		[19] ALTHOFF	84I	TASS	Aplanarity	

[1] DECAMP 90F looked for isolated charged particles, for isolated photons, and for four-jet final states. The modes $b' \rightarrow\ b g$ for B($b' \rightarrow\ b g$) > 65% $b' \rightarrow\ b\gamma$ for B($b' \rightarrow\ b\gamma$) > 5% are excluded. Charged Higgs decay were not discussed. NODE=Q008BPE;LINKAGE=DC

[2] ABDALLAH 07 searched for b' pair production at $E_{\rm cm}$=196–209 GeV, with 420 pb^{-1}. No signal leads to the 95% CL upper limits on B($b' \rightarrow\ b Z$) and B($b' \rightarrow\ c W$) for $m_{b'}$ = 96 to 103 GeV. NODE=Q008BPE;LINKAGE=DA

[3] ADRIANI 93G search for vector quarkonium states near Z and give limit on quarkonium-Z mixing parameter δm^2 <(10–30) GeV^2 (95%CL) for the mass 88–94.5 GeV. Using Richardson potential, a 1S ($b'\overline{b}'$) state is excluded for the mass range 87.7–94.7 GeV. This range depends on the potential choice. NODE=Q008BPE;LINKAGE=TB

[4] ABREU 90D assumed $m_{H^-} < m_{b'} - 3$ GeV. NODE=Q008BPE;LINKAGE=AB

[5] Superseded by ABREU 91F. NODE=Q008BPE;LINKAGE=AF

[6] AKRAWY 90B search was restricted to data near the Z peak at $E_{\rm cm}$ = 91.26 GeV at LEP. The excluded region is between 23.6 and 41.4 GeV if no H^+ decays exist. For charged Higgs decays the excluded regions are between (m_{H^+} + 1.5 GeV) and 45.5 GeV. NODE=Q008BPE;LINKAGE=AK

[7] AKRAWY 90J search for isolated photons in hadronic Z decay and derive B($Z \rightarrow\ b'\overline{b}'$)·B($b' \rightarrow\ \gamma$X)/B($Z \rightarrow$ hadrons) < 2.2×10^{-3}. Mass limit assumes B($b' \rightarrow\ \gamma$X) > 10%. NODE=Q008BPE;LINKAGE=T

[8] ABE 89E search at $E_{\rm cm}$ = 56–57 GeV at TRISTAN for multihadron events with a spherical shape (using thrust and acoplanarity) or containing isolated leptons. NODE=Q008BPE;LINKAGE=A

[9] ABE 89G search was at $E_{\rm cm}$ = 55–60.8 GeV at TRISTAN. NODE=Q008BPE;LINKAGE=B

[10] If the photonic decay mode is large (B($b' \rightarrow\ b\gamma$) > 25%), the ABRAMS 89C limit is 45.4 GeV. The limit for for Higgs decay ($b' \rightarrow\ c H^-$, $H^- \rightarrow\ \overline{c} s$) is 45.2 GeV. NODE=Q008BPE;LINKAGE=G

[11] ADACHI 89C search was at $E_{\rm cm}$ = 56.5–60.8 GeV at TRISTAN using multi-hadron events accompanying muons. NODE=Q008BPE;LINKAGE=C

[12] ADACHI 89C also gives limits for any mixture of $C C$ and $b g$ decays. NODE=Q008BPE;LINKAGE=F

[13] ENO 89 search at $E_{\rm cm}$ = 50–60.8 at TRISTAN. NODE=Q008BPE;LINKAGE=D

[14] ENO 89 considers arbitrary mixture of the charged current, $b g$, and $b\gamma$ decays. NODE=Q008BPE;LINKAGE=E

[15] IGARASHI 88 searches for leptons in low-thrust events and gives $\Delta R(b')$ < 0.26 (95% CL) assuming charged current decay, which translates to $m_{b'}$ > 24.4 GeV. NODE=Q008BPE;LINKAGE=S

[16] SAGAWA 88 set limit σ(top) < 6.1 pb at CL=95% for top-flavored hadron production from event shape analyses at $E_{\rm cm}$ = 52 GeV. By using the quark parton model cross-section formula near threshold, the above limit leads to lower mass bounds of 23.8 GeV for charge −1/3 quarks. NODE=Q008BPE;LINKAGE=Q

[17] ADEVA 86 give 95%CL upper bound on an excess of the normalized cross section, ΔR, as a function of the minimum c.m. energy (see their figure 3). Production of a pair of 1/3 charge quarks is excluded up to $E_{\rm cm}$ = 45.4 GeV. NODE=Q008BPE;LINKAGE=J

[18] ALTHOFF 84C narrow state search sets limit $\Gamma(e^+ e^-)$B(hadrons) <2.4 keV CL = 95% and heavy charge 1/3 quark pair production m >21 GeV, CL = 95%. NODE=Q008BPE;LINKAGE=K

[19] ALTHOFF 84I exclude heavy quark pair production for 7 < m <19 GeV (1/3 charge) using aplanarity distributions (CL = 95%). NODE=Q008BPE;LINKAGE=L

REFERENCES FOR Searches for (Fourth Generation) b' Quark

NODE=Q008

AAD 24BP PR D110 052009 G. Aad *et al.* (ATLAS Collab.) REFID=63079
HAYRAPETY... 24AQ PR D110 052004 A. Hayrapetyan *et al.* (CMS Collab.) REFID=63077
AAD 23AG EPJ C83 719 G. Aad *et al.* (ATLAS Collab.) REFID=62172
AAD 23AV PL B843 138019 G. Aad *et al.* (ATLAS Collab.) REFID=62364
AAD 23CQ JHEP 2311 168 G. Aad *et al.* (ATLAS Collab.) REFID=62612
TUMASYAN 23V JHEP 2307 020 A. Tumasyan *et al.* (CMS Collab.) REFID=62165
TUMASYAN 22O JHEP 2204 048 A. Tumasyan *et al.* (CMS Collab.) REFID=61788
SIRUNYAN 21AG JHEP 2112 106 A.M. Sirunyan *et al.* (CMS Collab.) REFID=61565
SIRUNYAN 20BI PR D102 112004 A.M. Sirunyan *et al.* (CMS Collab.) REFID=60740
SIRUNYAN 19AI EPJ C79 90 A.M. Sirunyan *et al.* (CMS Collab.) REFID=59702
SIRUNYAN 19AQ EPJ C79 364 A.M. Sirunyan *et al.* (CMS Collab.) REFID=59721
SIRUNYAN 19BW PR D100 072001 A.M. Sirunyan *et al.* (CMS Collab.) REFID=60010
AABOUD 18AW JHEP 1808 048 M. Aaboud *et al.* (ATLAS Collab.) REFID=59145
AABOUD 18CE JHEP 1812 039 M. Aaboud *et al.* (ATLAS Collab.) REFID=59369
AABOUD 18CL PR D98 092005 M. Aaboud *et al.* (ATLAS Collab.) REFID=59472
AABOUD 18CP PR D98 112010 M. Aaboud *et al.* (ATLAS Collab.) REFID=59500
AABOUD 18CR PRL 121 211801 M. Aaboud *et al.* (ATLAS Collab.) REFID=59532
SIRUNYAN 18BM JHEP 1808 177 A.M. Sirunyan *et al.* (CMS Collab.) REFID=59137

SIRUNYAN	18Q	PR D97 072008	A.M. Sirunyan *et al.*	(CMS Collab.)	REFID=58920
SIRUNYAN	17AU	JHEP 1711 085	A.M. Sirunyan *et al.*	(CMS Collab.)	REFID=58344
AAD	16AH	JHEP 1602 110	G. Aad *et al.*	(ATLAS Collab.)	REFID=57318
KHACHATRY...	16AN	PR D93 112009	V. Khachatryan *et al.*	(CMS Collab.)	REFID=57295
KHACHATRY...	16I	JHEP 1601 166	V. Khachatryan *et al.*	(CMS Collab.)	REFID=57141
AAD	15AR	JHEP 1508 105	G. Aad *et al.*	(ATLAS Collab.)	REFID=56648
AAD	15BY	JHEP 1510 150	G. Aad *et al.*	(ATLAS Collab.)	REFID=56863
AAD	15CN	PR D92 112007	G. Aad *et al.*	(ATLAS Collab.)	REFID=57013
AAD	15Z	PR D91 112011	G. Aad *et al.*	(ATLAS Collab.)	REFID=56592
AAD	14AZ	JHEP 1411 104	G. Aad *et al.*	(ATLAS Collab.)	REFID=56201
CHATRCHYAN	13I	JHEP 1301 154	S. Chatrchyan *et al.*	(CMS Collab.)	REFID=54941
AAD	12AT	PRL 109 032001	G. Aad *et al.*	(ATLAS Collab.)	REFID=54229
AAD	12AU	PRL 109 071801	G. Aad *et al.*	(ATLAS Collab.)	REFID=54230
AAD	12BC	PR D86 012007	G. Aad *et al.*	(ATLAS Collab.)	REFID=54358
AAD	12BE	JHEP 1204 069	G. Aad *et al.*	(ATLAS Collab.)	REFID=54458
CHATRCHYAN	12BH	PR D86 112003	S. Chatrchyan *et al.*	(CMS Collab.)	REFID=54772
CHATRCHYAN	12X	JHEP 1205 123	S. Chatrchyan *et al.*	(CMS Collab.)	REFID=54460
AALTONEN	11J	PRL 106 141803	T. Aaltonen *et al.*	(CDF Collab.)	REFID=16439
ABAZOV	11F	PRL 106 081801	V.M. Abazov *et al.*	(D0 Collab.)	REFID=16469
CHATRCHYAN	11L	PL B701 204	S. Chatrchyan *et al.*	(CMS Collab.)	REFID=16643
AALTONEN	10H	PRL 104 091801	T. Aaltonen *et al.*	(CDF Collab.)	REFID=53271
FLACCO	10	PRL 105 111801	C.J. Flacco *et al.*	(UCI, HAIF)	REFID=53412
ATRE	09	PR D79 054018	A. Atre *et al.*		REFID=54081
ABAZOV	08X	PRL 101 111802	V.M. Abazov *et al.*	(D0 Collab.)	REFID=52402
HUANG	08	PR D77 037302	P.Q. Hung, M. Sher	(UVA, WILL)	REFID=52505
AALTONEN	07C	PR D76 072006	T. Aaltonen *et al.*	(CDF Collab.)	REFID=51994
ABDALLAH	07	EPJ C50 507	J. Abdallah *et al.*	(DELPHI Collab.)	REFID=51764
ACOSTA	03	PRL 90 131801	D. Acosta *et al.*	(CDF Collab.)	REFID=49298
AFFOLDER	00	PRL 84 835	A. Affolder *et al.*	(CDF Collab.)	REFID=47308
ABE	98N	PR D58 051102	F. Abe *et al.*	(CDF Collab.)	REFID=46140
ABACHI	97D	PRL 78 3818	S. Abachi *et al.*	(D0 Collab.)	REFID=45459
FROGGATT	97	ZPHY C73 333	C.D. Froggatt, D.J. Smith, H.B. Nielsen	(GLAS+)	REFID=45376
ABACHI	95F	PR D52 4877	S. Abachi *et al.*	(D0 Collab.)	REFID=44482
ADRIANI	93G	PL B313 326	O. Adriani *et al.*	(L3 Collab.)	REFID=43472
ADRIANI	93M	PRPL 236 1	O. Adriani *et al.*	(L3 Collab.)	REFID=43644
MUKHOPAD...	93	PR D48 2105	B. Mukhopadhyaya, D.P. Roy	(TATA)	REFID=43481
ABE	92	PRL 68 447	F. Abe *et al.*	(CDF Collab.)	REFID=41874
Also		PR D45 3921	F. Abe *et al.*	(CDF Collab.)	REFID=42068
ABE	92G	PR D45 3921	F. Abe *et al.*	(CDF Collab.)	REFID=42068
ABREU	91F	NP B367 511	P. Abreu *et al.*	(DELPHI Collab.)	REFID=41840
ABE	90B	PRL 64 147	F. Abe *et al.*	(CDF Collab.)	REFID=40986
ABE	90D	PL B234 382	K. Abe *et al.*	(VENUS Collab.)	REFID=41105
ABREU	90D	PL B242 536	P. Abreu *et al.*	(DELPHI Collab.)	REFID=41317
ADACHI	90	PL B234 197	I. Adachi *et al.*	(TOPAZ Collab.)	REFID=41106
AKESSON	90	ZPHY C46 179	T. Akesson *et al.*	(UA2 Collab.)	REFID=41051
AKRAWY	90B	PL B236 364	M.Z. Akrawy *et al.*	(OPAL Collab.)	REFID=40987
AKRAWY	90J	PL B246 285	M.Z. Akrawy *et al.*	(OPAL Collab.)	REFID=41336
ALBAJAR	90B	ZPHY C48 1	C. Albajar *et al.*	(UA1 Collab.)	REFID=41312
DECAMP	90F	PL B236 511	D. Decamp *et al.*	(ALEPH Collab.)	REFID=41035
ABE	89E	PR D39 3524	K. Abe *et al.*	(VENUS Collab.)	REFID=40844
ABE	89G	PRL 63 1776	K. Abe *et al.*	(VENUS Collab.)	REFID=40951
ABRAMS	89C	PRL 63 2447	G.S. Abrams *et al.*	(Mark II Collab.)	REFID=40966
ADACHI	89C	PL B229 427	I. Adachi *et al.*	(TOPAZ Collab.)	REFID=40952
ENO	89	PRL 63 1910	S. Eno *et al.*	(AMY Collab.)	REFID=40953
ALBAJAR	88	ZPHY C37 505	C. Albajar *et al.*	(UA1 Collab.)	REFID=40464
ALTARELLI	88	NP B308 724	G. Altarelli *et al.*	(CERN, ROMA, ETH)	REFID=40899
IGARASHI	88	PRL 60 2359	S. Igarashi *et al.*	(AMY Collab.)	REFID=40606
SAGAWA	88	PRL 60 93	H. Sagawa *et al.*	(AMY Collab.)	REFID=40453
ADEVA	86	PR D34 681	B. Adeva *et al.*	(Mark-J Collab.)	REFID=40171
ALTHOFF	84C	PL 138B 441	M. Althoff *et al.*	(TASSO Collab.)	REFID=12195
ALTHOFF	84I	ZPHY C22 307	M. Althoff *et al.*	(TASSO Collab.)	REFID=12196