



$$I(J^P) = 1(\frac{3}{2}^+) \text{ Status: } ***$$

I, J, P need confirmation.

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

NODE=S062

NODE=S062

Σ_b^* MASS

NODE=S062205

Σ_b^{*+} MASS

NODE=S062M+
NODE=S062M+

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5830.32±0.27 OUR AVERAGE			
5830.28±0.14±0.24	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
5832.1 ±0.7 $\begin{smallmatrix} +1.7 \\ -1.8 \end{smallmatrix}$	² AALTONEN	12F CDF	$p\bar{p}$ at 1.96 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			
² Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow K^- \pi^+$ decays.			

NODE=S062M+;LINKAGE=A
NODE=S062M+;LINKAGE=AL

Σ_b^{*-} MASS

NODE=S062M-
NODE=S062M-

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5834.74±0.30 OUR AVERAGE			
5834.73±0.17±0.25	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
5835.1 ±0.6 $\begin{smallmatrix} +1.7 \\ -1.8 \end{smallmatrix}$	² AALTONEN	12F CDF	$p\bar{p}$ at 1.96 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			
² Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow K^- \pi^+$ decays.			

NODE=S062M-;LINKAGE=A
NODE=S062M-;LINKAGE=AL

$m_{\Sigma_b^{*+}} - m_{\Sigma_b^{*-}}$

NODE=S062DMI
NODE=S062DMI

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-4.37±0.33 OUR AVERAGE	Error includes scale factor of 1.6.		
-4.45±0.22±0.01	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
-3.0 $\begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$ ±0.1	² AALTONEN	12F CDF	$p\bar{p}$ at 1.96 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			
² Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow K^- \pi^+$ decays.			

NODE=S062DMI;LINKAGE=A
NODE=S062DMI;LINKAGE=AL

$m_{\Sigma_b^{*+}} - m_{\Sigma_b^+}$

NODE=S062DMP
NODE=S062DMP

VALUE	DOCUMENT ID	TECN	COMMENT
19.73±0.18±0.01	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			

NODE=S062DMP;LINKAGE=A

$m_{\Sigma_b^{*-}} - m_{\Sigma_b^-}$

NODE=S062DMM
NODE=S062DMM

VALUE	DOCUMENT ID	TECN	COMMENT
19.09±0.22±0.02	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			

NODE=S062DMM;LINKAGE=A

Σ_b^* WIDTH

NODE=S062230

Σ_b^{*+} WIDTH

NODE=S062W+
NODE=S062W+

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9.4 ±0.5 OUR AVERAGE			
9.34±0.47±0.26	¹ AAIJ	19A LHCb	pp at 7, 8 TeV
11.5 $\begin{smallmatrix} +2.7 \\ -2.2 \end{smallmatrix}$ $\begin{smallmatrix} +1.0 \\ -1.5 \end{smallmatrix}$	² AALTONEN	12F CDF	$p\bar{p}$ at 1.96 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			
² Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow K^- \pi^+$ decays.			

NODE=S062W+;LINKAGE=A
NODE=S062W+;LINKAGE=AL

Σ_b^{*-} WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10.4 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.3.		
10.68 ± 0.60 ± 0.33	¹ AAIJ	19A LHCb	$p p$ at 7, 8 TeV
7.5 $^{+2.2}_{-1.8}$ $^{+0.9}_{-1.4}$	² AALTONEN	12F CDF	$p \bar{p}$ at 1.96 TeV
¹ Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays.			
² Measured using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\Lambda_c^+ \rightarrow K^- \pi^+$ decays.			

NODE=S062W-
NODE=S062W-

NODE=S062W-;LINKAGE=A
NODE=S062W-;LINKAGE=AL

$$m_{\Sigma_b^*} - m_{\Sigma_b}$$

NODE=S062DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$21.2^{+2.0+0.4}_{-1.9-0.3}$	¹ AALTONEN	07K CDF	$p\bar{p}$ at 1.96 TeV
¹ Observed four $\Lambda_b^0 \pi^\pm$ resonances in the fully reconstructed decay mode $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ where $\Lambda_c^+ \rightarrow p K^- \pi^+$. Assumes $m_{\Sigma_b^{*+}} - m_{\Sigma_b^+} = m_{\Sigma_b^{*-}} - m_{\Sigma_b^-}$			

NODE=S062DM

NODE=S062DM;LINKAGE=AA

Σ_b^* DECAY MODES

	Mode	Fraction (Γ_i/Γ)
Γ_1	$\Lambda_b^0 \pi$	dominant

NODE=S062215;NODE=S062

DESIG=1

Σ_b^* BRANCHING RATIOS

$\Gamma(\Lambda_b^0 \pi)/\Gamma_{\text{total}}$				Γ_1/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT	
dominant	AALTONEN	07κ	CDF $p\bar{p}$ at 1.96 TeV	

NODE=S062225

NODE=S062R01
NODE=S062R01

Σ_b^* REFERENCES

AAIJ	19A	PRL 122 012001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	12F	PR D85 092011	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	07K	PRL 99 202001	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=S062

REFID=59550
REFID=54118
REFID=52023