B^0–$\overline{B}^0$ MIXING

Updated April 2008 by O. Schneider (Ecole Polytechnique Fédérale de Lausanne).

There are two neutral B^0–$\overline{B}^0$ meson systems, B^0_d–$\overline{B}^0_d$ and B^0_s–$\overline{B}^0_s$ (generically denoted B^0_q–$\overline{B}^0_q$, $q = s, d$), which exhibit particle-antiparticle mixing [1]. This mixing phenomenon is described in Ref. 2. In the following, we adopt the notation introduced in Ref. 2, and assume CPT conservation throughout. In each system, the light (L) and heavy (H) mass eigenstates,

$$|B_{\rm L,H}\rangle = p|B^0_q\rangle \pm q|\overline{B}^0_q\rangle\,, \tag{1}$$

have a mass difference $\Delta m_q = m_{\rm H} - m_{\rm L} > 0$, and a total decay width difference $\Delta\Gamma_q = \Gamma_{\rm L} - \Gamma_{\rm H}$. In the absence of CP violation in the mixing, $|q/p| = 1$, these differences are given by $\Delta m_q = 2|M_{12}|$ and $|\Delta\Gamma_q| = 2|\Gamma_{12}|$, where M_{12} and Γ_{12} are the off-diagonal elements of the mass and decay matrices [2]. The evolution of a pure $|B^0_q\rangle$ or $|\overline{B}^0_q\rangle$ state at $t = 0$ is given by

$$|B^0_q(t)\rangle = g_+(t)\,|B^0_q\rangle + \frac{q}{p}\,g_-(t)\,|\overline{B}^0_q\rangle\,, \tag{2}$$

$$|\overline{B}^0_q(t)\rangle = g_+(t)\,|\overline{B}^0_q\rangle + \frac{p}{q}g_-(t)\,|B^0_q\rangle\,, \tag{3}$$

which means that the flavor states remain unchanged (+) or oscillate into each other (−) with time-dependent probabilities proportional to

$$|g_\pm(t)|^2 = \frac{e^{-\Gamma_q t}}{2}\left[\cosh\left(\frac{\Delta\Gamma_q}{2}\,t\right) \pm \cos(\Delta m_q\,t)\right]\,, \tag{4}$$

where $\Gamma_q = (\Gamma_{\rm H} + \Gamma_{\rm L})/2$. In the absence of CP violation, the time-integrated mixing probability $\int |g_-(t)|^2\,dt/(\int |g_-(t)|^2\,dt + \int |g_+(t)|^2\,dt)$ is given by

$$\chi_q = \frac{x_q^2 + y_q^2}{2(x_q^2+1)}\,, \qquad \text{where} \qquad x_q = \frac{\Delta m_q}{\Gamma_q}\,, \qquad y_q = \frac{\Delta\Gamma_q}{2\Gamma_q}\,. \tag{5}$$

Standard Model predictions and phenomenology

In the Standard Model, the transitions $B^0_q \to \overline{B}^0_q$ and $\overline{B}^0_q \to B^0_q$ are due to the weak interaction. They are described, at the lowest order, by box diagrams involving two W bosons and two



Figure 1: Dominant box diagrams for the $B_q^0 \to \overline{B}_q^0$ transitions ($q = d$ or s). Similar diagrams exist where one or both t quarks are replaced with c or u quarks.

up-type quarks (see Fig. 1), as is the case for K^0–$\overline{K}^0$ mixing. However, the long range interactions arising from intermediate virtual states are negligible for the neutral B meson systems, because the large B mass is off the region of hadronic resonances. The calculation of the dispersive and absorptive parts of the box diagrams yields the following predictions for the off-diagonal element of the mass and decay matrices [3],

$$M_{12} = -\frac{G_F^2 m_W^2 \eta_B m_{B_q} B_{B_q} f_{B_q}^2}{12\pi^2} S_0(m_t^2/m_W^2)\,(V_{tq}^* V_{tb})^2 \,, \tag{6}$$

$$\Gamma_{12} = \frac{G_F^2 m_b^2 \eta_B' m_{B_q} B_{B_q} f_{B_q}^2}{8\pi} \times \left[(V_{tq}^* V_{tb})^2 + V_{tq}^* V_{tb} V_{cq}^* V_{cb}\, \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right) + (V_{cq}^* V_{cb})^2\, \mathcal{O}\left(\frac{m_c^4}{m_b^4}\right) \right] , \tag{7}$$

where G_F is the Fermi constant, m_W the W boson mass, and m_i the mass of quark i; m_{B_q}, f_{B_q} and B_{B_q} are the B_q^0 mass, weak decay constant and bag parameter, respectively. The known function $S_0(x_t)$ can be approximated very well by $0.784\, x_t^{0.76}$ [4], and V_{ij} are the elements of the CKM matrix [5]. The QCD corrections η_B and η_B' are of order unity. The only non-negligible contributions to M_{12} are from box diagrams involving two top quarks. The phases of M_{12} and Γ_{12} satisfy

$$\phi_M - \phi_\Gamma = \pi + \mathcal{O}\left(\frac{m_c^2}{m_b^2}\right) , \tag{8}$$

implying that the mass eigenstates have mass and width differences of opposite signs. This means that, like in the K^0–$\overline{K}^0$ system, the heavy state is expected to have a smaller decay width

than that of the light state: $\Gamma_{\rm H} < \Gamma_{\rm L}$. Hence, $\Delta\Gamma = \Gamma_{\rm L} - \Gamma_{\rm H}$ is expected to be positive in the Standard Model.

Furthermore, the quantity

$$\left|\frac{\Gamma_{12}}{M_{12}}\right| \simeq \frac{3\pi}{2}\frac{m_b^2}{m_W^2}\frac{1}{S_0(m_t^2/m_W^2)} \sim \mathcal{O}\left(\frac{m_b^2}{m_t^2}\right) \tag{9}$$

is small, and a power expansion of $|q/p|^2$ yields

$$\left|\frac{q}{p}\right|^2 = 1 + \left|\frac{\Gamma_{12}}{M_{12}}\right| \sin(\phi_M - \phi_\Gamma) + \mathcal{O}\left(\left|\frac{\Gamma_{12}}{M_{12}}\right|^2\right) . \tag{10}$$

Therefore, considering both Eqs. (8) and (9), the CP-violating parameter

$$1 - \left|\frac{q}{p}\right|^2 \simeq \mathrm{Im}\left(\frac{\Gamma_{12}}{M_{12}}\right) \tag{11}$$

is expected to be very small: $\sim \mathcal{O}(10^{-3})$ for the B_d^0–$\overline{B}_d^0$ system and $\lesssim \mathcal{O}(10^{-4})$ for the B_s^0–$\overline{B}_s^0$ system [6].

In the approximation of negligible CP violation in mixing, the ratio $\Delta\Gamma_q/\Delta m_q$ is equal to the small quantity $|\Gamma_{12}/M_{12}|$ of Eq. (9); it is hence independent of CKM matrix elements, *i.e.*, the same for the B_d^0–$\overline{B}_d^0$ and B_s^0–$\overline{B}_s^0$ systems. Recent calculations [7] yield $\sim 5 \times 10^{-3}$ with a $\sim 20\%$ uncertainty. Given the current experimental knowledge on the mixing parameter x_q

$$\begin{cases} x_d = 0.776 \pm 0.008 & (B_d^0\text{–}\overline{B}_d^0 \text{ system}) \\ x_s = 26.1 \quad \pm 0.5 & (B_s^0\text{–}\overline{B}_s^0 \text{ system}) \end{cases} , \tag{12}$$

the Standard Model thus predicts that $\Delta\Gamma_d/\Gamma_d$ is very small (below 1%), but $\Delta\Gamma_s/\Gamma_s$ considerably larger ($\sim 10\%$). These width differences are caused by the existence of final states to which both the B_q^0 and $\overline{B}_q^0$ mesons can decay. Such decays involve $b \to c\overline{c}q$ quark-level transitions, which are Cabibbo-suppressed if $q = d$ and Cabibbo-allowed if $q = s$.

A recent and complete set of Standard Model predictions for all mixing parameters in both the B_d^0–$\overline{B}_d^0$ and B_s^0–$\overline{B}_s^0$ systems can be found in Ref. 7.

Experimental issues and methods for oscillation analyses

Time-integrated measurements of B^0–$\overline{B}^0$ mixing were published for the first time in 1987 by UA1 [8] and ARGUS [9], and

since then by many other experiments. These measurements are typically based on counting same-sign and opposite-sign lepton pairs from the semileptonic decay of the produced $b\overline{b}$ pairs. Such analyses cannot easily separate the contributions from the different b-hadron species, therefore, the clean environment of $\Upsilon(4S)$ machines (where only B_d^0 and charged B_u mesons are produced) is in principle best suited to measure χ_d.

However, better sensitivity is obtained from time-dependent analyses aiming at the direct measurement of the oscillation frequencies Δm_d and Δm_s, from the proper time distributions of B_d^0 or B_s^0 candidates identified through their decay in (mostly) flavor-specific modes, and suitably tagged as mixed or unmixed. This is particularly true for the B_s^0–$\overline{B}_s^0$ system, where the large value of x_s implies maximal mixing, *i.e.*, $\chi_s \simeq 1/2$. In such analyses, the B_d^0 or B_s^0 mesons are either fully reconstructed, partially reconstructed from a charm meson, selected from a lepton with the characteristics of a $b \to \ell^-$ decay, or selected from a reconstructed displaced vertex. At high-energy colliders (LEP, SLC, Tevatron), the proper time $t = \dfrac{m_B}{p} L$ is measured from the distance L between the production vertex and the B decay vertex, and from an estimate of the B momentum p. At asymmetric B factories (KEKB, PEP-II), producing $e^+e^- \to \Upsilon(4S) \to B_d^0\overline{B}_d^0$ events with a boost $\beta\gamma$ (= 0.425, 0.55), the proper time difference between the two B candidates is estimated as $\Delta t \simeq \dfrac{\Delta z}{\beta\gamma c}$, where Δz is the spatial separation between the two B decay vertices along the boost direction. In all cases, the good resolution needed on the vertex positions is obtained with silicon detectors.

The average statistical significance $\mathcal{S}$ of a B_d^0 or B_s^0 oscillation signal can be approximated as [10]

$$\mathcal{S} \approx \sqrt{N/2}\, f_{\rm sig}\, (1 - 2\eta)\, e^{-(\Delta m\, \sigma_t)^2/2}\,, \tag{13}$$

where N is the number of selected and tagged candidates, $f_{\rm sig}$ is the fraction of signal in that sample, η is the total mistag probability, and σ_t is the resolution on proper time (or proper time difference). The quantity $\mathcal{S}$ decreases very quickly as Δm increases; this dependence is controlled by σ_t, which is therefore

a critical parameter for Δm_s analyses. At high-energy colliders, the proper time resolution $\sigma_t \sim \frac{m_B}{\langle p \rangle}\sigma_L \oplus t\frac{\sigma_p}{p}$ includes a constant contribution due to the decay length resolution σ_L (typically 0.05–0.3 ps), and a term due to the relative momentum resolution σ_p/p (typically 10–20% for partially reconstructed decays), which increases with proper time. At B factories, the boost of the B mesons is estimated from the known beam energies, and the term due to the spatial resolution dominates (typically 1–1.5 ps because of the much smaller B boost).

In order to tag a B candidate as mixed or unmixed, it is necessary to determine its flavor both in the initial state and in the final state. The initial and final state mistag probabilities, η_i and η_f, degrade $\mathcal{S}$ by a total factor $(1-2\eta) = (1-2\eta_i)(1-2\eta_f)$. In lepton-based analyses, the final state is tagged by the charge of the lepton from $b \to \ell^-$ decays; the largest contribution to η_f is then due to $\overline{b} \to \overline{c} \to \ell^-$ decays. Alternatively, the charge of a reconstructed charm meson (D^{*-} from B_d^0 or D_s^- from B_s^0), or that of a kaon hypothesized to come from a $b \to c \to s$ decay [11], can be used. For fully-inclusive analyses based on topological vertexing, final-state tagging techniques include jet-charge [12] and charge-dipole [13,14] methods. At high-energy colliders, the methods to tag the initial state (*i.e.*, the state at production), can be divided into two groups: the ones that tag the initial charge of the $\overline{b}$ quark contained in the B candidate itself (same-side tag), and the ones that tag the initial charge of the other b quark produced in the event (opposite-side tag). On the same side, the charge of a track from the primary vertex is correlated with the production state of the B if that track is a decay product of a B^{**} state or the first particle in the fragmentation chain [15,16]. Jet- and vertex-charge techniques work on both sides and on the opposite side, respectively. Finally, the charge of a lepton from $b \to \ell^-$ or of a kaon from $b \to c \to s$ can be used as opposite side tags, keeping in mind that their performance is degraded due to integrated mixing. At SLC, the beam polarization produced a sizeable forward-backward asymmetry in the $Z \to b\overline{b}$ decays, and provided another very interesting and effective initial state

tag based on the polar angle of the B candidate [13]. Initial state tags have also been combined to reach $\eta_i \sim 26\%$ at LEP [16,17], or even 22% at SLD [13] with full efficiency. In the case $\eta_f = 0$, this corresponds to an effective tagging efficiency $Q = \epsilon D^2 = \epsilon(1-2\eta)^2$, where ϵ is the tagging efficiency, in the range $23-31\%$. The equivalent figure achieved by CDF during Tevatron Run I was $\sim 3.5\%$ [18], reflecting the fact that tagging is more difficult at hadron colliders. The current CDF and DØ analyses of Tevatron Run II data reach $\epsilon D^2 = (1.8 \pm 0.1)\%$ [19] and $(2.5 \pm 0.2)\%$ [20] for opposite-side tagging, while same-side kaon tagging (for B_s^0 oscillation analyses) is contributing an additional $3.7-4.8\%$ at CDF [19], and pushes the combined performance to $(4.5 \pm 0.9)\%$ at DØ [21].

At B factories, the flavor of a B_d^0 meson at production cannot be determined, since the two neutral B mesons produced in a $\Upsilon(4S)$ decay evolve in a coherent P-wave state where they keep opposite flavors at any time. However, as soon as one of them decays, the other follows a time-evolution given by Eqs. (2) or (3), where t is replaced with Δt (which will take negative values half of the time). Hence, the "initial state" tag of a B can be taken as the final-state tag of the other B. Effective tagging efficiencies Q of 30% are achieved by BABAR and Belle [22], using different techniques including $b \to \ell^-$ and $b \to c \to s$ tags. It is worth noting that, in this case, mixing of the other B (*i.e.*, the coherent mixing occurring before the first B decay) does not contribute to the mistag probability.

In the absence of experimental observation of a decay-width difference, oscillation analyses typically neglect $\Delta\Gamma$ in Eq. (4), and describe the data with the physics functions $\Gamma e^{-\Gamma t}(1 \pm \cos(\Delta m t))/2$ (high-energy colliders) or $\Gamma e^{-\Gamma|\Delta t|}(1 \pm \cos(\Delta m \Delta t))/4$ (asymmetric $\Upsilon(4S)$ machines). As can be seen from Eq. (4), a non-zero value of $\Delta\Gamma$ would effectively reduce the oscillation amplitude with a small time-dependent factor that would be very difficult to distinguish from time resolution effects. Measurements of Δm_d are usually extracted from the data using a maximum likelihood fit. To extract information useful for the interpretation of B_s^0 oscillation searches and for the combination of their results, a method [10] is followed in

which a B_s^0 oscillation amplitude $\mathcal{A}$ is measured as a function of a fixed test value of Δm_s, using a maximum likelihood fit based on the functions $\Gamma_s e^{-\Gamma_s t}(1 \pm \mathcal{A}\cos(\Delta m_s t))/2$. To a good approximation, the statistical uncertainty on $\mathcal{A}$ is Gaussian and equal to $1/\mathcal{S}$ from Eq. (13). If Δm_s is equal to its true value, one expects $\mathcal{A} = 1$ within the total uncertainty $\sigma_{\mathcal{A}}$; in case a signal is seen, its observed (or expected) significance will be defined as $\mathcal{A}/\sigma_{\mathcal{A}}$ (or $1/\sigma_{\mathcal{A}}$). However, if Δm_s is (far) below its true value, a measurement consistent with $\mathcal{A} = 0$ is expected. A value of Δm_s can be excluded at 95% CL if $\mathcal{A} + 1.645\,\sigma_{\mathcal{A}} \leq 1$ (since the integral of a normal distribution from $-\infty$ to 1.645 is equal to 0.95). Because of the proper time resolution, the quantity $\sigma_{\mathcal{A}}(\Delta m_s)$ is a steadily increasing function of Δm_s. We define the sensitivity for 95% CL exclusion of Δm_s values (or for a $3\,\sigma$ or $5\,\sigma$ observation of B_s^0 oscillations) as the value of Δm_s for which $1/\sigma_{\mathcal{A}} = 1.645$ (or $1/\sigma_{\mathcal{A}} = 3$ or 5).

B_d^0 mixing studies

Many B_d^0–$\overline{B}_d^0$ oscillations analyses have been published [23] by the ALEPH [24], BABAR [25], Belle [26], CDF [15], DØ [20], DELPHI [14,27], L3 [28], and OPAL [29,30] collaborations. Although a variety of different techniques have been used, the individual Δm_d results obtained at high-energy colliders have remarkably similar precision. Their average is compatible with the recent and more precise measurements from asymmetric B factories. The systematic uncertainties are not negligible; they are often dominated by sample composition, mistag probability, or b-hadron lifetime contributions. Before being combined, the measurements are adjusted on the basis of a common set of input values, including the b-hadron lifetimes and fractions published in this *Review*. Some measurements are statistically correlated. Systematic correlations arise both from common physics sources (fragmentation fractions, lifetimes, branching ratios of b hadrons), and from purely experimental or algorithmic effects (efficiency, resolution, tagging, background description). Combining all published measurements [14,15,20,24–30] and accounting for all identified correlations yields $\Delta m_d = 0.507 \pm 0.003(\text{stat}) \pm 0.003(\text{syst})\ \text{ps}^{-1}$ [31], a result dominated by the B factories.

On the other hand, ARGUS and CLEO have published time-integrated measurements [32–34], which average to $\chi_d = 0.182 \pm 0.015$. Following Ref. 34, the width difference $\Delta\Gamma_d$ could in principle be extracted from the measured value of Γ_d and the above averages for Δm_d and χ_d (see Eq. (5)), provided that $\Delta\Gamma_d$ has a negligible impact on the Δm_d measurements. However, direct time-dependent studies published by DELPHI [14] and BABAR [35] provide stronger constraints, which can be combined to yield $\mathrm{sign}(\mathrm{Re}\lambda_{\mathrm{CP}})\Delta\Gamma_{\mathrm{d}}/\Gamma_{\mathrm{d}} = 0.009 \pm 0.037$ [31].

Assuming $\Delta\Gamma_d = 0$ and no CP violation in mixing, and using the measured B_d^0 lifetime of 1.530 ± 0.009 ps, the Δm_d and χ_d results are combined to yield the world average

$$\Delta m_d = 0.507 \pm 0.005 \ \ \mathrm{ps}^{-1} \tag{14}$$

or, equivalently,

$$\chi_d = 0.1878 \pm 0.0024\,. \tag{15}$$

Evidence for CP violation in B_d^0 mixing has been searched for, both with flavor-specific and inclusive B_d^0 decays, in samples where the initial flavor state is tagged. In the case of semileptonic (or other flavor-specific) decays, where the final-state tag is also available, the following asymmetry [2]

$$\mathcal{A}_{\mathrm{SL}}^d = \frac{N(\overline{B}_d^0(t) \to \ell^+\nu_\ell X) - N(B_d^0(t) \to \ell^-\overline{\nu}_\ell X)}{N(\overline{B}_d^0(t) \to \ell^+\nu_\ell X) + N(B_d^0(t) \to \ell^-\overline{\nu}_\ell X)} \simeq 1 - |q/p|_d^2 \tag{16}$$

has been measured, either in time-integrated analyses at CLEO [34,36], CDF [37,38] and DØ [39], or in time-dependent analyses at LEP [30,40,41], BABAR [35,42,43] and Belle [44]. In the inclusive case, also investigated at LEP [40,41,45], no final-state tag is used, and the asymmetry [46]

$$\frac{N(B_d^0(t) \to \mathrm{all}) - N(\overline{B}_d^0(t) \to \mathrm{all})}{N(B_d^0(t) \to \mathrm{all}) + N(\overline{B}_d^0(t) \to \mathrm{all})} \simeq \mathcal{A}_{\mathrm{SL}}^d \left[\frac{x_d}{2}\sin(\Delta m_d\, t) - \sin^2\left(\frac{\Delta m_d\, t}{2}\right)\right] \tag{17}$$

must be measured as a function of the proper time to extract information on CP violation. In all cases, asymmetries compatible with zero have been found, with a precision limited by the available statistics. A simple average of all

published results for the B_d^0 meson [30,34–36,39,41,42,44,45] yields $\mathcal{A}_{\rm SL}^d = -0.0049 \pm 0.0038$, under the assumption of no CP violation in B_s^0 mixing. Published results at B factories only [34–36,42,44], where no B_s^0 is produced, average to

$$\mathcal{A}_{\rm SL}^d = -0.0005 \pm 0.0056\,, \text{ or } |q/p|_d = 1.0002 \pm 0.0028\,, \quad (18)$$

a result which does not yet constrain the Standard Model.

The Δm_d result of Eq. (14) provides an estimate of $2|M_{12}|$, and can be used, together with Eq. (6), to extract the magnitude of the CKM matrix element V_{td} within the Standard Model [47]. The main experimental uncertainties on the resulting estimate of $|V_{td}|$ come from m_t and Δm_d; however, the extraction is at present completely dominated by the uncertainty on the hadronic matrix element $f_{B_d}\sqrt{B_{B_d}} = 244 \pm 26$ MeV obtained from lattice QCD calculations [48].

B_s^0 mixing studies

In the decade before the Tevatron Run II results became available, B_s^0–$\overline{B}_s^0$ oscillations have been the subject of many studies from ALEPH [49], CDF [50], DELPHI [14,17,51], OPAL [52] and SLD [13,53,54]. The most sensitive analyses appeared to be the ones based on inclusive lepton samples. Because of their better proper time resolution, the small data samples analyzed inclusively at SLD, as well as the fully reconstructed B_s decays at LEP were also very useful to explore the high Δm_s region. However, all results were limited by the available statistics. All published measurements of the B_s^0 oscillation amplitude [13,14,17,49–53] are averaged [31] to yield the combined amplitudes $\mathcal{A}$ shown in Fig. 2 (bottom) as a function of Δm_s. The individual results have been adjusted to common physics inputs, and all known correlations have been accounted for; the sensitivities of the inclusive analyses, which depend directly through Eq. (13) on the assumed fraction f_s of B_s^0 mesons in an unbiased sample of weakly-decaying b hadrons, have also been rescaled to a common average of $f_s = 0.105 \pm 0.009$. The combined sensitivity for 95% CL exclusion of Δm_s values is found to be 18.3 ps^{-1}. All values of Δm_s below 14.6 ps^{-1} are excluded at 95% CL, while the values between 14.6 and 21.7 ps^{-1} cannot be excluded, because the data is compatible



Figure 2: Combined measurements of the B_s^0 oscillation amplitude as a function of Δm_s. Top: CDF result based on Run II data, published in 2006 [19]. Middle: Average of all preliminary DØ results available at the end of 2007 [21]. Bottom: Average of all other results (mainly from LEP and SLD) published between 1997 and 2004. All measurements are dominated by statistical uncertainties. Neighboring points are statistically correlated. See full-color version on color pages at end of book.

with a signal in this region. However, the largest deviation from $\mathcal{A} = 0$ in this range is a $1.9\,\sigma$ effect only, so no signal can be claimed.

Tevatron Run II results based on 1 fb^{-1} of data became available in 2006. After DØ [55] reported $17 < \Delta m_s < 21$ ps^{-1} (90% CL) and a most probable value of 19 ps^{-1} with an observed (expected) significance of $2.5\,\sigma$ ($0.9\,\sigma$), CDF [19] published the first direct evidence of B_s^0 oscillations shortly followed by a $> 5\sigma$ observation (shown at the top of Fig. 2). The measured value of Δm_s is

$$\Delta m_s = 17.77 \pm 0.10(\text{stat}) \pm 0.07(\text{syst})\ \ \text{ps}^{-1}\,, \qquad (19)$$

based on samples of flavour-tagged hadronic and semileptonic B_s^0 decays, partially or fully reconstructed in flavour-specific final states. More recently, DØ [21] obtained with 2.4 fb^{-1} an independent 2.9σ preliminary evidence for B_s^0 oscillations (middle of Fig. 2) at $\Delta m_s = 18.53 \pm 0.93(\text{stat}) \pm 0.30(\text{syst})$ ps^{-1} [56], consistent with the CDF measurement.

The information on $|V_{ts}|$ obtained in the framework of the Standard Model is hampered by the hadronic uncertainty, as in the B_d^0 case. However, several uncertainties cancel in the frequency ratio

$$\frac{\Delta m_s}{\Delta m_d} = \frac{m_{B_s}}{m_{B_d}}\,\xi^2 \left|\frac{V_{ts}}{V_{td}}\right|^2 , \tag{20}$$

where $\xi = (f_{B_s}\sqrt{B_{B_s}})/(f_{B_d}\sqrt{B_{B_d}}) = 1.210\,^{+0.047}_{-0.035}$ is an SU(3) flavor-symmetry breaking factor obtained from lattice QCD calculations [48]. Using the measurements of Eqs. (14) and (19), one can extract

$$\left|\frac{V_{td}}{V_{ts}}\right| = 0.2060 \pm 0.0012(\text{exp})\,^{+0.0080}_{-0.0060}(\text{lattice})\,, \tag{21}$$

in good agreement with (but much more precise than) the recent results obtained by the Belle [57] and BABAR [58] collaborations based on the observation of the $b \to d\gamma$ transition. The CKM matrix can be constrained using experimental results on observables such as Δm_d, Δm_s, $|V_{ub}/V_{cb}|$, ϵ_K, and $\sin(2\beta)$ together with theoretical inputs and unitarity conditions [47,59,60]. The constraint from our knowledge on the ratio $\Delta m_s/\Delta m_d$ is presently more effective in limiting the position of the apex of the CKM unitarity triangle than the one obtained from the Δm_d measurements alone, due to the reduced hadronic uncertainty in Eq. (20). We also note that the measured value of Δm_s is consistent with the Standard Model prediction obtained from CKM fits where no experimental information on Δm_s is used, *e.g.*, 20.6 ± 2.6 ps^{-1} [59] or $17.7\,^{+6.4}_{-2.1}$ ps^{-1} [60].

Information on $\Delta\Gamma_s$ can be obtained by studying the proper time distribution of untagged B_s^0 samples [61]. In the case of an inclusive B_s^0 selection [62], or a semileptonic (or flavour-specific) B_s^0 decay selection [17,63,64], both the short- and

long-lived components are present, and the proper time distribution is a superposition of two exponentials with decay constants $\Gamma_{\rm L,H} = \Gamma_s \pm \Delta\Gamma_s/2$. In principle, this provides sensitivity to both Γ_s and $(\Delta\Gamma_s/\Gamma_s)^2$. Ignoring $\Delta\Gamma_s$ and fitting for a single exponential leads to an estimate of Γ_s with a relative bias proportional to $(\Delta\Gamma_s/\Gamma_s)^2$. An alternative approach, which is directly sensitive to first order in $\Delta\Gamma_s/\Gamma_s$, is to determine the lifetime of B_s^0 candidates decaying to CP eigenstates; measurements exist for $B_s^0 \to K^+K^-$ [65], $B_s^0 \to J/\psi\phi$ [66,67], and $B_s^0 \to D_s^{(*)+}D_s^{(*)-}$ [68], which are mostly CP-even states [69]. However, in the case of $B_s^0 \to J/\psi\phi$, this technique has now been replaced by more sensitive time-dependent angular analyses that allow the simultaneous extraction of $\Delta\Gamma_s/\Gamma_s$ and the CP-even and CP-odd amplitudes [70,71]. Estimates of $\Delta\Gamma_s/\Gamma_s$ have also been obtained directly from measurements of the $B_s^0 \to D_s^{(*)+}D_s^{(*)-}$ branching ratio [68,72], under the assumption that these decays account for all the CP-even final states (however, no systematic uncertainty due to this assumption is given, so the averages quoted below will not include these estimates).

Applying the combination procedure of Ref. 31 (including the constraint from the flavour-specific lifetime measurements) on the published results [17,63,66,68,70,71] yields

$$\Delta\Gamma_s/\Gamma_s = +0.069\,^{+0.058}_{-0.062} \qquad \text{and} \qquad 1/\Gamma_s = 1.470\,^{+0.026}_{-0.027}\ \text{ps}\,, \tag{22}$$

or equivalently

$$1/\Gamma_{\rm L} = 1.419\,^{+0.039}_{-0.038}\ \text{ps} \qquad \text{and} \qquad 1/\Gamma_{\rm H} = 1.525\,^{+0.062}_{-0.063}\ \text{ps}\,, \tag{23}$$

under the assumption of no CP violation in B_s^0 mixing.

Recent studies also consider CP violation, either in untagged [70,71] or tagged [73,74] $B_s^0 \to J/\psi\phi$ decays, and start to constrain the phase difference $-2\beta_s$ between the B_s^0 mixing diagram and the $b \to c\bar{c}s$ tree decay diagram. In the Standard Model, $\beta_s = \arg(-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*))$ is expected to be about one degree [7]. A proper combination of the current Tevatron constraints on β_s requires extra information not available in

the original publications and is being prepared in collaboration between CDF and DØ.

On the other hand CP violation in B_s^0 mixing has been investigated through the asymmetry between positive and negative same-sign muon pairs from semi-leptonic decays of $b\bar{b}$ pairs [37–39], and directly through the charge asymmetry of $B_s^0 \to D_s\mu\nu X$ decays [75]. Combining all published results [37,39,75] with the knowledge of CP violation in B_d^0 mixing from Eq. (18) leads to

$$\mathcal{A}_{\rm SL}^s = -0.0030 \pm 0.0101\,, \;\; \text{or} \;\; |q/p|_s = 1.0015 \pm 0.0051\,. \quad (24)$$

A large New Physics phase could possibly contribute to both CP violation in $B_s^0 \to J/\psi\phi$, and to the mixing phase difference of Eq. (8) on which $\mathcal{A}_{\rm SL}^s$ depends. Combined fits [76,77] of β_s and $\mathcal{A}_{\rm SL}^s$ measurements already yield interesting constraints on this New Physics phase. A deviation from the Standard Model, with a significance of more than 3σ, has recently been claimed [77], based on a preliminary analysis including the latest Tevatron B_s^0 mixing results [19,38,39,73–75].

Average b-hadron mixing probability and b-hadron production fractions in Z decays and at high energy

Mixing measurements can significantly improve our knowledge on the fractions f_u, f_d, f_s, and $f_{\rm baryon}$, defined as the fractions of B_u, B_d^0, B_s^0 and b-baryon in an unbiased sample of weakly decaying b hadrons produced in high-energy collisions. Indeed, time-integrated mixing analyses performed with lepton pairs from $b\bar{b}$ events at high energy measure the quantity

$$\overline{\chi} = f'_d\,\chi_d + f'_s\,\chi_s\,, \quad (25)$$

where f'_d and f'_s are the fractions of B_d^0 and B_s^0 hadrons in a sample of semileptonic b-hadron decays. Assuming that all b hadrons have the same semileptonic decay width implies $f'_q = f_q/(\Gamma_q\tau_b)$ $(q = s, d)$, where τ_b is the average b-hadron lifetime. Hence $\overline{\chi}$ measurements, together with the χ_d average of Eq. (15) and the very good approximation $\chi_s = 1/2$ (in fact $\chi_s = 0.49927 \pm 0.00003$ from Eqs. (5), (19) and (22)), provide constraints on the fractions f_d and f_s.

The LEP experiments have measured $f_s \times \mathrm{BR}(B_s^0 \to D_s^- \ell^+ \nu_\ell X)$ [78], $\mathrm{BR}(b \to \Lambda_b^0) \times \mathrm{BR}(\Lambda_b^0 \to \Lambda_c^+ \ell^- \overline{\nu}_\ell X)$ [79], and $\mathrm{BR}(b \to \Xi_b^-) \times \mathrm{BR}(\Xi_b^- \to \Xi^- \ell^- \overline{\nu}_\ell X)$ [80] from partially reconstructed final states including a lepton, f_{baryon} from protons identified in b events [81], and the production rate of charged b hadrons [82]. The b-hadron fractions measured at CDF using double semileptonic $K^*\mu\mu$ and $\phi\mu\mu$ final states [83] and electron-charm final states [84] are at slight discrepancy with the ones measured at LEP. Furthermore the averages of the $\overline{\chi}$ values measured at LEP, 0.1259 ± 0.0042 [85], and at Tevatron, 0.147 ± 0.011 [39,86], show a $1.8\,\sigma$ deviation with respect to each other. This may be a hint that the fractions at the Tevatron might be different from the ones in Z decays. Combining [31] all the available information under the constraints $f_u = f_d$ and $f_u + f_d + f_s + f_{\mathrm{baryon}} = 1$ yields the two set of averages shown in Table 1. The second set, obtained using both LEP and Tevatron results, has larger errors than the first set, obtained using LEP results only, because we have applied scale factors as advocated by the PDG for the treatment of marginally consistent data.

Table 1: $\overline{\chi}$ and b-hadron fractions (see text).

	in Z decays	at high energy
$\overline{\chi}$	0.1259 ± 0.0042	0.1284 ± 0.0069
$f_u = f_d$	0.402 ± 0.009	0.399 ± 0.011
f_s	0.104 ± 0.009	0.110 ± 0.012
f_{baryon}	0.091 ± 0.015	0.092 ± 0.019

Summary and prospects

B^0–$\overline{B}^0$ mixing has been and still is a field of intense study. While fairly little experimental progress was recently achieved in the B_d^0 sector, impressive new B_s^0 results are becoming available from Run II of the Tevatron. B_s^0 oscillations are now established and the mass difference in the B_s^0–$\overline{B}_s^0$ system is measured very accurately, with a central value compatible with the Standard Model (SM) expectation and a relative precision

(0.7%) matching that in the B_d^0–$\overline{B}_d^0$ system (0.9%). However, the extraction of $|V_{td}/V_{ts}|$ from these measurements in the SM framework is limited by the hadronic uncertainty, which will be an important challenge to reduce in the future. New time-dependent angular analyses of $B_s^0 \to J/\psi\phi$ decays and measurements of time-integrated B_s^0 asymmetries at CDF and DØ are improving our knowledge of the other B_s^0 mixing parameters: while CP violation in B_s^0–$\overline{B}_s^0$ mixing is consistent with zero, with an uncertainty still large compared to the SM prediction, the relative decay width difference $\Delta\Gamma_s/\Gamma_s$ is now determined to an absolute precision of $\sim 6\%$, smaller than the central value of the SM prediction. The data prefer $\Gamma_{\rm L} > \Gamma_{\rm H}$ as predicted in the SM.

Improved B_s^0 results are still to come, with very promising short-term prospects, both for $\Delta\Gamma_s$ and CP-violating phases induced by mixing such as β_s and $\arg(-M_{12}/\Gamma_{12})$. Although first interesting experimental constraints have been published, very little is known yet about these phases, which are predicted to be very small in the SM. A full search for New Physics effects in these observables will require statistics beyond that of the Tevatron. These will eventually become available at CERN's Large Hadron Collider, scheduled to start operation in 2008, where LHCb expects to be able to measure β_s down to the SM value after many years of operations [87].

B mixing may still reveal a surprize, but much effort is needed for this, both on the experimental and theoretical sides, in particular to further reduce the hadronic uncertainties of lattice QCD calculations. In the long term, a stringent check of the consistency of the B_d^0 and B_s^0 mixing amplitudes (magnitudes and phases) with all other measured flavour-physics observables will be possible within the SM, leading to very tight limits (or otherwise new interesting knowledge!) on New Physics.

References

1. T.D. Lee and C.S. Wu, Ann. Rev. Nucl. Sci. **16**, 511 (1966); I.I. Bigi and A.I. Sanda, *CP Violation*, Cambridge Univ. Press, 2000; G.C. Branco, L. Lavoura, and J.P. Silva, *CP Violation*, Clarendon Press Oxford, 1999.

2. See the review on "CP violation in meson decays," by D. Kirkby and Y. Nir in this publication.
3. A.J. Buras, W. Slominski, and H. Steger, Nucl. Phys. **B245**, 369 (1984).
4. T. Inami and C.S. Lim, Prog. Theor. Phys. **65**, 297 (1981); for the power-like approximation, see A.J. Buras and R. Fleischer, page 91 in *Heavy Flavours II*, eds., A.J. Buras and M. Lindner, Singapore World Scientific, 1998.
5. M. Kobayashi and K. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
6. I.I. Bigi *et al.*, in *CP violation,* ed. C. Jarlskog, Singapore World Scientific, 1989.
7. A. Lenz and U. Nierste, JHEP **0607**, 072 (2007), `arXiv: hep-ph/0612167v3`.
8. C. Albajar *et al.* (UA1 Collab.), Phys. Lett. **B186**, 247 (1987).
9. H. Albrecht *et al.* (Argus Collab.), Phys. Lett. **B192**, 245 (1987).
10. H.-G. Moser and A. Roussarie, Nucl. Instrum. Methods **384**, 491 (1997).
11. SLD Collab., SLAC-PUB-7228, SLAC-PUB-7229, and SLAC-PUB-7230, *28th Int. Conf. on High Energy Physics,* Warsaw, 1996; J. Wittlin, PhD thesis, SLAC-R-582, 2001.
12. Aleph Collab., *Int. Europhysics Conf. on High Energy Physics,* Jerusalem, 1997.
13. K. Abe *et al.* (SLD Collab.), Phys. Rev. **D67**, 012006 (2003).
14. J. Abdallah *et al.* (DELPHI Collab.), Eur. Phys. J. **C28**, 155 (2003).
15. F. Abe *et al.* (CDF Collab.), Phys. Rev. Lett. **80**, 2057 (1998) and Phys. Rev. **D59**, 032001 (1999); Phys. Rev. **D60**, 051101 (1999); Phys. Rev. **D60**, 072003 (1999); T. Affolder *et al.* (CDF Collab.), Phys. Rev. **D60**, 112004 (1999).
16. R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C4**, 367 (1998); Eur. Phys. J. **C7**, 553 (1999).
17. P. Abreu *et al.* (DELPHI Collab.), Eur. Phys. J. **C16**, 555 (2000); Eur. Phys. J. **C18**, 229 (2000).
18. See tagging summary on page 160 of K. Anikeev *et al.*, "B physics at the Tevatron: Run II and beyond," FERMILAB-PUB-01/97, `hep-ph/0201071`, and references therein.

19. A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **97**, 242003 (2006); this result supersedes A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **97**, 062003 (2006).

20. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. **D74**, 112002 (2006).

21. DØCollab., DØ note 5474-CONF, August 2007, and DØ note 5254-CONF, October 2006.

22. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **94**, 161803 (2005);
K.-F. Chen *et al.* (Belle Collab.), Phys. Rev. **D72**, 012004 (2005).

23. Throughout this paper, we omit references of results that have been superseded by new published measurements.

24. D. Buskulic *et al.* (Aleph Collab.), Z. Phys. **C75**, 397 (1997).

25. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **88**, 221802 (2002) and Phys. Rev. **D66**, 032003 (2002); Phys. Rev. Lett. **88**, 221803 (2002); Phys. Rev. **D67**, 072002 (2003); Phys. Rev. **D73**, 012004 (2006).

26. N.C. Hastings *et al.* (Belle Collab.), Phys. Rev. **D67**, 052004 (2003); Y. Zheng *et al.* (Belle Collab.), Phys. Rev. **D67**, 092004 (2003); K. Abe *et al.* (Belle Collab.), Phys. Rev. **D71**, 072003 (2005).

27. P. Abreu *et al.* (DELPHI Collab.), Z. Phys. **C76**, 579 (1997).

28. M. Acciarri *et al.* (L3 Collab.), Eur. Phys. J. **C5**, 195 (1998).

29. G. Alexander *et al.* (OPAL Collab.), Z. Phys. **C72**, 377 (1996); K. Ackerstaff *et al.* (OPAL Collab.), Z. Phys. **C76**, 417 (1997); G. Abbiendi *et al.* (OPAL Collab.), Phys. Lett. **B493**, 266 (2000).

30. K. Ackerstaff *et al.* (OPAL Collab.), Z. Phys. **C76**, 401 (1997).

31. E. Barberio *et al.* (HFAG Collab.), "Averages of b-hadron properties at the end of 2006," `arXiv:0704.3575v1 [hep-ex]`, April 2007; the combined results on b-hadron fractions, lifetimes and mixing parameters published in this *Review* have been obtained by the B oscillations working group of the Heavy Flavour Averaging Group (HFAG), using the methods and procedures described in Chapter 3 of the above paper, after updating the list of inputs; for more information, see
`http://www.slac.stanford.edu/xorg/hfag/osc/`.

32. H. Albrecht *et al.* (ARGUS Collab.), Z. Phys. **C55**, 357 (1992); Phys. Lett. **B324**, 249 (1994).
33. J. Bartelt *et al.* (CLEO Collab.), Phys. Rev. Lett. **71**, 1680 (1993).
34. B.H. Behrens *et al.* (CLEO Collab.), Phys. Lett. **B490**, 36 (2000).
35. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **92**, 181801 (2004) and Phys. Rev. **D70**, 012007 (2004).
36. D.E. Jaffe *et al.* (CLEO Collab.), Phys. Rev. Lett. **86**, 5000 (2001).
37. F. Abe *et al.* (CDF Collab.), Phys. Rev. **D55**, 2546 (1997).
38. CDF Collab., CDF note 9015, October 2007.
39. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. **D74**, 092001 (2006).
40. DELPHI Collab., contrib. 449 to *Int. Europhysics Conf. on High Energy Physics,* Jerusalem, 1997.
41. R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C20**, 431 (2001).
42. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **96**, 251802 (2006).
43. BABAR Collab., `arXiv:hep-ex/0607091v1`, *33rd Int. Conf. on High Energy Physics,* Moscow, 2006.
44. E. Nakano *et al.* (Belle Collab.), Phys. Rev. **D73**, 112002 (2006).
45. G. Abbiendi *et al.* (OPAL Collab.), Eur. Phys. J. **C12**, 609 (2000).
46. M. Beneke, G. Buchalla, and I. Dunietz, Phys. Lett. **B393**, 132 (1997); I. Dunietz, Eur. Phys. J. **C7**, 197 (1999).
47. See the review on the CKM quark-mixing matrix by A. Ceccucci, Z. Ligeti, and Y. Sakai in this publication.
48. M. Okamoto, *XXIIIth Int. Symp. on Lattice Field Theory,* Dublin, July 2005, `hep-lat/0510113`; these estimates are obtained by combining the unquenched lattice QCD calculations from A. Gray *et al.* (HPQCD Collab.), Phys. Rev. Lett. **95**, 212001 (2005) and S. Aoki *et al.* (JLQCD Collab.), Phys. Rev. Lett. **91**, 212001 (2003).
49. A. Heister *et al.* (Aleph Collab.), Eur. Phys. J. **C29**, 143 (2003).
50. F. Abe *et al.* (CDF Collab.), Phys. Rev. Lett. **82**, 3576 (1999).
51. J. Abdallah *et al.* (DELPHI Collab.), Eur. Phys. J. **C35**, 35 (2004).

52. G. Abbiendi *et al.* (OPAL Collab.), Eur. Phys. J. **C11**, 587 (1999); Eur. Phys. J. **C19**, 241 (2001).
53. K. Abe *et al.* (SLD Collab.), Phys. Rev. **D66**, 032009 (2002).
54. SLD Collab., SLAC-PUB-8568, *30th Int. Conf. on High Energy Physics,* Osaka, 2000.
55. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **97**, 021802 (2006).
56. D0 Collab., D0 note 5618-CONF v1.1, March 2008.
57. D. Mohapatra *et al.* (Belle Collab.), Phys. Rev. Lett. **96**, 221601 (2006), updated in M. Nakao, talk at *23rd Int. Symp. on Lepton and Photon Interactions at High Energy,* Daegu, 2007.
58. B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **98**, 151802 (2007).
59. M. Bona *et al.* (UTfit Collab.), `arXiv:hep-ph/0606167v2` and updated results at `http://utfit.roma1.infn.it/`.
60. J. Charles *et al.* (CKMfitter Collab.), Eur. Phys. J. **C41**, 1 (2005) & updated results at `http://ckmfitter.in2p3.fr/`.
61. K. Hartkorn and H.-G. Moser, Eur. Phys. J. **C8**, 381 (1999).
62. M. Acciarri *et al.* (L3 Collab.), Phys. Lett. **B438**, 417 (1998).
63. D. Buskulic *et al.* (Aleph Collab.), Phys. Lett. **B377**, 205 (1996); K. Ackerstaff *et al.* (OPAL Collab.), Phys. Lett. **B426**, 161 (1998); F. Abe *et al.* (CDF Collab.), Phys. Rev. **D59**, 032004 (1999).
64. (CDF Collab.), CDF note 7386, March 2005; CDF note 7757, August 2005; V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **97**, 241801 (2006).
65. D. Tonelli for the (CDF Collab.), `arXiv:hep-ex/0605038v1`, May 2006.
66. F. Abe *et al.* (CDF Collab.), Phys. Rev. **D57**, 5382 (1998).
67. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **94**, 041001 (2005); (CDF Collab.), CDF note 7409, May 2004.
68. R. Barate *et al.* (Aleph Collab.), Phys. Lett. **B486**, 286 (2000).
69. R. Aleksan *et al.*, Phys. Lett. **B316**, 567 (1993).
70. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 121803 (2008).
71. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **98**, 121801 (2007).

72. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **99**, 241801 (2007); A. Abulencia *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 021803 (2008).

73. T. Aaltonen *et al.* (CDF Collab.), Phys. Rev. Lett. **100**, 161802 (2008).

74. V.M. Abazov *et al.* (D0 Collab.), `arXiv:0802.2255v1 [hep-ex]`, submitted to Phys. Rev. Lett.

75. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. Lett. **98**, 151801 (2007).

76. V.M. Abazov *et al.* (D0 Collab.), Phys. Rev. **D76**, 057101 (2007).

77. M. Bona *et al.* (UTfit Collab.), `arXiv:0803.0659v1 [hep-ph]`, March 2008.

78. P. Abreu *et al.* (DELPHI Collab.), Phys. Lett. **B289**, 199 (1992); P.D. Acton *et al.* (OPAL Collab.), Phys. Lett. **B295**, 357 (1992); D. Buskulic *et al.* (Aleph Collab.), Phys. Lett. **B361**, 221 (1995).

79. P. Abreu *et al.* (DELPHI Collab.), Z. Phys. **C68**, 375 (1995); R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C2**, 197 (1998).

80. D. Buskulic *et al.* (Aleph Collab.), Phys. Lett. **B384**, 449 (1996); J. Abdallah *et al.* (DELPHI Collab.), Eur. Phys. J. **C44**, 299 (2005).

81. R. Barate *et al.* (Aleph Collab.), Eur. Phys. J. **C5**, 205 (1998).

82. J. Abdallah *et al.* (DELPHI Collab.), Phys. Lett. **B576**, 29 (2003).

83. F. Abe *et al.* (CDF Collab.), Phys. Rev. **D60**, 092005 (1999).

84. T. Affolder *et al.* (CDF Collab.), Phys. Rev. Lett. **84**, 1663 (2000).

85. Aleph, DELPHI, L3, OPAL, and SLD Collabs., "Precision electroweak measurements on the Z resonance," Physics Reports **427**, 257 (2006); we use the $\overline{\chi}$ average given in Eq. (5.39).

86. D. Acosta *et al.* (CDF Collab.), Phys. Rev. **D69**, 012002 (2004).

87. L. Fernández for the LHCb Collab., LHCb note 2006-047, Acta Phys. Pol. **B38**, 931 (2007).