

Review of particle properties: Supplement to 1974 edition

Particle Data Group

Vladimir Chaloupka and Claude Bricman

CERN, 1211 Genève 23, Switzerland

Angela Barbaro-Galtieri, Denyse M. Chew*, Robert L. Kelly, Thomas A. Lasinski†, Alan Rittenberg, Arthur H. Rosenfeld, Thomas G. Trippe, Fumiyo Uchiyama, and George P. Yost

Lawrence Berkeley Laboratory,‡ University of California, Berkeley, California 94720

Naomi Barash-Schmidt

Brandeis University, Waltham, Massachusetts 02154

Matts Roos

Department of Nuclear Physics, University of Helsinki, Helsinki 17, Finland

This supplement to the 1974 edition of "Review of Particle Properties," Particle Data Group [Phys. Lett. **50B**, No. 1 (1974)], contains an announcement concerning the postponement of the usual Review, a list of Errata, and a tabulation of the experimental results on the newly discovered mesons.

For several reasons, outlined below, the Particle Data Group has decided to postpone the next regular edition of the "Review of Particle Properties" until April 1976, at which time it will appear in *Reviews of Modern Physics*. In this supplement we present a brief list of errata to the 1974 edition,¹ and, because of the importance of the recently discovered mesons, we also give addenda to the Data Card Listings containing entries for these particles only. Two mini-reviews in the Listings discuss various aspects of these mesons.

Our reasons for delaying the regular edition are primarily the following: (1) Apart from the excitement generated by the new mesons, there has been somewhat of a shift in the last 2-3 years from experiments studying particle properties to those studying interaction properties. This dimi-

nution in new particle data, especially data of a dramatic nature (again with the exception noted above), leads us to believe that this time we can wait two years for a complete update. (2) In line with the shift in emphasis to interactions, the Berkeley members of the Group have been developing a data storage and retrieval system into which we plan to put all experimental data which we can extract from physics documents (journal articles, preprints, reports, theses, etc.). The primary usage of this system, at least initially, will be for such interaction properties as cross sections, angular and momentum distributions, polarization measurements, and density matrix elements. We are currently in the process of putting together our initial data base and are aiming to produce, as the first output of our system, an "Index of Particle Physics Data".² This Index will enable the user to locate articles on the basis of beam, target, momentum and, to a limited extent, reaction final state and properties measured. Since this project is presently occupying most of the time of the Berkeley Group, publishing the Review this year would significantly delay the Index. (3) Finally, the recent substantial increases in printing costs also contributed to the decision to postpone publication.

² Particle Data Group, "Index of Particle Physics Data", LBL-90 (to be issued).

* On leave of absence from University of Paris VI, Paris, France.

† Presently at Stanford Linear Accelerator Center, Stanford, California 94305, USA.

‡ The Berkeley Particle Data Center is jointly supported by the U.S. Energy Research and Development Administration, the Office of Standard Reference Data of the National Bureau of Standards, and the National Science Foundation.

¹ Particle Data Group: V. Chaloupka, C. Bricman, A. Barbaro-Galtieri, D. M. Chew, R. L. Kelly, T. A. Lasinski, A. Rittenberg, A. H. Rosenfeld, T. G. Trippe, F. Uchiyama, N. Barash-Schmidt, P. Söding, and M. Roos, *Physics Letters* **50B**, No. 1 (1974).

ERRATA

Following is a list of errata in the 1974 edition of Review of Particle Properties [Phys. Lett. 50B, No. 1 (1974)]. Two page numbers are given for each item; the first is for the Review itself, and the second is for the associated 1974 Particle Properties Data Booklet.

In the Physical and Numerical Constants Table (p. 15/35), the line defining an atmosphere should read:

$$1 \text{ atmosphere} = 1033.2275 \text{ dynes cm}^{-2} = 1.01325 \text{ bar.}$$

In the Relativistic Kinematics Table (p. 21/51), equations 20-1 and 21-1 should read:

$$\gamma = \frac{s + m_b^2 - m_a^2}{2m_b \sqrt{s}} = \frac{m_b + E_a^{\text{lab}}}{\sqrt{s}} \quad (20-1),$$

$$\eta = \frac{\sqrt{\Delta(s, m_a^2, m_b^2)}}{2m_b \sqrt{s}} = \frac{P_{\text{inc}}}{\sqrt{s}} \quad (21-1).$$

In the Particle Detectors, Absorbers, and Ranges section, the Atomic and Nuclear Properties of Materials Table (p. 30/74-75) has several errors; as a result we reproduce the entire table, including updated values where they were available.

In the Electromagnetic Relations section (p. 31/76), ϵ and μ are defined to be the relative permittivity and permeability, respectively, for both CGS and MKSA units. To convert the MKSA formulas to more standard notation, where ϵ and μ are defined to be absolute, substitute

$$\epsilon \text{ for } \epsilon\epsilon_0, \text{ and } \mu \text{ for } \mu\mu_0.$$

Atomic and Nuclear Properties of Materials*

Material	Z	A	Nuclear cross section σ^a [barns]	Nuclear collision length L_{coll}^b		Absorption length λ^b [cm]	dE/dx min ^c		Radiation length L_{rad}^d		Density ^e [g/cm ³] () is for gas [g/l]	Refractive index n^e () is $(n-1) \times 10^6$ for gas
				[g/cm ²]	[cm]		[MeV/g/cm ²]	[MeV/cm]	[g/cm ²]	[cm]		
H ₂	1	1.01	0.039	43.0	607	790	4.12	0.292	63.05	890	{0.0708 {(0.090)	{1.112 {(140)
D ₂	1	2.01	0.074	45.1	273	342	2.07	0.342	126.1	764	0.165	1.128
He	2	4.00	0.134	49.6	397	478	1.94	0.243	94.32	755	{0.125 {(0.178)	{1.024 {(35)
Li	3	6.94	0.215	53.6	100.4	120.6	1.65	0.902	82.76	155	0.534	-
Be	4	9.01	0.270	55.4	30.0	36.7	1.61	2.97	65.19	35.3	1.848	-
C	6	12.01	0.340	58.7	≈37.8	49.9	1.78	≈2.76	42.70	≈27.5	≈1.55 ^f	-
N ₂	7	14.01	0.390	59.7	73.8	99.4	1.82	1.47	37.99	47.0	{0.808 {(1.25)	{1.205 {(300)
Ne	10	20.18	0.520	64.4	53.7	74.9	1.73	2.08	28.94	24.0	{1.207 {(0.90)	{1.092 {(67)
Al	13	26.98	0.650	68.9	25.5	37.2	1.62	4.37	24.01	8.9	2.70	-
A	18	39.95	0.890	74.5	53.2	80.9	1.51	2.11	19.55	14.0	{1.40 {(1.78)	{1.233 {(283)
Fe	26	55.85	1.160	79.9	10.2	17.1	1.48	11.6	13.84	1.76	7.87	-
Cu	29	63.54	1.270	83.1	9.3	14.8	1.44	12.9	12.86	1.43	8.96	-
Sn	50	118.69	2.040	96.6	13.2	22.8	1.28	9.4	8.82	1.21	7.31	-
W	74	183.85	2.810	108.6	5.6	10.3	1.17	22.6	6.76	0.35	19.3	-
Pb	82	207.19	3.080	111.7	9.8	18.5	1.13	12.8	6.37	0.56	11.35	-
U	92	238.03	3.380	116.9	≈6.2	12.0	1.09	≈20.7	6.00	≈0.32	≈18.95	-
Air				60.2	50000 ^g	67500 ^g	1.82	0.0022 ^g	36.66	30050 ^g	{0.001205 ^g {(1.29)	{1.000273 ^g {(293)
H ₂ O				58.3	58.3	78.8	2.03	2.03	36.08	36.1	1.00	1.33
H ₂ (bubble chamber 26°K) ^h				43.0	≈683	887	4.12	≈0.26	63.05	≈1000	≈0.063 ^h	1.112
D ₂ (bubble chamber 31°K) ^h				45.1	≈322	403	2.07	≈0.29	126.1	≈900	≈0.140 ^h	1.110
H-Ne mixture (50 mole percent) ⁱ				62.9	154.5	215	1.84	0.75	29.70	73.0	0.407	1.092
Propane (C ₃ H ₈) ^j				55.0	134	176	2.28	0.98	45.38	111	{0.41 ^j {(2.0)	{1.25 ^j {(1005)
Freon 13B1 (CF ₃ Br) ^j				74.3	≈49.5	73.5	1.52	≈2.3	16.53	≈11	{≈1.50 ^j {(8.71)	{1.238 ^j {(750)
Iford emulsion				79.5	23.6	39.1	1.44	5.49	11.02	2.94	3.815	-
NaI				91.9	25.0	41.3	1.32	4.84	9.49	2.59	3.67	1.775
LiF				61.1	23.1	30.7	1.69	4.46	39.25	14.9	2.64	1.394
Polyethylene (CH ₂)				55.7	≈59.6	78.4	2.09	≈1.95	44.78	≈48	0.92-0.95	-
Mylar (C ₅ H ₄ O ₂)				58.5	42.1	56.1	1.91	2.65	39.95	28.7	1.39	-
Polystyrene, scintillator (CH) ^k				57.0	55.2	68.5	2.03	1.97	43.8	42.9	1.032	1.581
Lucite, Plexiglas (C ₅ H ₈ O ₂)				57.7	≈48.9	65.0	1.95	≈1.65	40.55	≈34.5	1.16-1.20	≈1.49
Spark or proportional chamber ^l				0.05%		0.03%	-	0.073		2.7%	0.046	-
Shielding concrete ^m				64.9	26.0	32.2	1.70	4.25	26.7	10.7	2.5	-
CO ₂ ⁿ				60.4	33800	46000	1.82	0.0033	36.2	20210	(1.79) ⁿ	(410) ⁿ
Freon 12 (CCl ₂ F ₂) ⁿ				68.1	13800	20200	1.64	0.0081	23.7	4810	(4.93) ⁿ	(1080) ⁿ
Freon 13 (CClF ₃) ⁿ				66.0	15000	21400	1.70	0.0072	27.15	6380	(4.26) ⁿ	(720) ⁿ

* Table revised January 1975 by J. Engler and F. Mönig. For details and references, see CERN NP Internal Report 74-1.

a) σ of neutrons ($\approx \sigma$ of protons) at 20 GeV from Landolt-Bornstein, New Series I, Vol. 5. Energy dependence for all nuclei $\approx 1/2$ percent/GeV (from 5-25 GeV).

b) $L_{coll} = A/(N\sigma)$. In the absorption length the elastic scattering is subtracted.

c) From W.H. Barkas and M.J. Berger, Tables of Energy Losses and Ranges of Heavy Charged Particles, NASA-SP-3013 (1964).

d) From Y.S. Tsai, Pair Production and Bremsstrahlung of Charged Leptons, SLAC-PUB-1365 (1974), Table III.6.

e) Values for solids, or the liquid phase at boiling point, except where noted. Values in parentheses for gaseous phase STP (0°C, 1 atm.), except where noted.

f) Density variable.

g) Gas at 20°C.

h) Density may vary about $\pm 3\%$, depending on operating conditions.

i) Values for typical working condition with H₂ target: 50 mole percent, 29°K, 7 atm.

j) Values for typical chamber working conditions: Propane $\sim 57^\circ\text{C}$, 8-10 atm. Freon 13B1 $\sim 28^\circ\text{C}$, 8-10 atm.

k) Typical scintillator; e.g. PILOT B and NE 102A have an atomic ratio H/C = 1.10.

l) Values for typical construction: 2 layers 50 μm Cu/Be wires, 8 mm gap, 60% argon, 40% isobutane or CO₂; 2 layers 50 μm Mylar/Aclar foils.

m) Standard shielding blocks, typical composition O₂ 52%, Si 32.5%, Ca 6%, Na 1.5%, Fe 2%, Al 4% plus reinforcing iron bars. Attenuation length $l = 115 \pm 5$ g/cm², also valid for earth (typical $\rho = 2.15$) from CERN-LRL-RHEL Shielding exp. UCRL 17841 (1968).

n) Used in Čerenkov counters, value at 26°C and 1 atm. Indices of refraction from E.R. Hayes, R.A. Schluter, and A. Tamosaitis, ANL-6916 (1964).

NEW HEAVY MESONS

Data Card Listings

The New Heavy Mesons

In the following, we shall briefly review the experiments bearing on the recently-discovered $J/\psi(3100)$ and $\psi(3700)$ particles, as well as a peak in the e^+e^- total cross section, tentatively labeled $X(4100)$ (whose resonant nature has not yet been established).

As of March 1975, the best known of these particles is the $J/\psi(3100)$. A summary of the experiments relevant to this particle is given in Table I. It has been observed in the e^+e^- mass spectrum produced in p-Be collisions at the AGS (AUBERT 74), and in e^+e^- , charged hadronic, and also $\mu^+\mu^-$ decay modes

formed at SPEAR (AUGUSTIN 74). It has also been observed in e^+e^- collisions at ADONE (BACCI 74), at DORIS (BRAUNSCHWEIG 74,75; CRIEGEE 75), and at SPEAR (FORD 75), in photoproduction at FNAL (KNAPP1 75), in n-Be collisions at FNAL (KNAPP2 75), and in photoproduction using a mixture of real and low- q^2 virtual photons at SLAC (DAKIN 75). A Wisconsin-SLAC collaboration has also observed production of the $J/\psi(3100)$, as well as the $\psi(3700)$, in a bremsstrahlung beam of peak energies ranging from 13.5 to 21.5 GeV (Univ. of Wisconsin-SLAC collaboration, private communication).

TABLE I. List of Experiments Bearing on the $J/\psi(3100)$ Meson.

Accelerator	Authors	Decay				Comments
		e^+e^-	$\mu^+\mu^-$	Hadrons	Others	
<u>Formation in e^+e^- Collisions</u>						
SPEAR	{ AUGUSTIN 74 } { LYNCH 75 } { FORD 75 }	x	x ^a	x		• Co-discovery; see AUBERT 74
		x	x		$\gamma\gamma$ ^b	• $\gamma\gamma$ decay mode forbidden if $J=1$, or if $C=-1$ and C conserved
		(See following 3 papers)				
ADONE	{ BACCI 74 } { ASH 74 } { BALDINI-CELIO 74 } { BARBIELLINI 74 }	x	x	x		
		x	x	x		
		x				
DORIS	{ BRAUNSCHWEIG 74 } { BRAUNSCHWEIG 75 } { CRIEGEE 75 }	x				
					$\gamma\gamma, \pi^0\gamma$ ^b	• $\gamma\gamma$ decay mode forbidden if $J=1$, or if $C=-1$ and C conserved
				x		• Signal in collinear 2-prongs also observed
<u>Production in Hadron Collisions</u>						
BNL-AGS	{ AUBERT 74 } { CHRISTENSON 70 }	x				• From p-Be at 28 GeV/c: Co-discovery; see AUGUSTIN 74
			x ^c			• From p-U at 28 GeV/c: Observed a shoulder above 3 GeV in $\mu^+\mu^-$ mass spectrum
FNAL	KNAPP2 75		x			• From n-Be at 70 - 250 GeV/c
<u>Photoproduction</u>						
FNAL	KNAPP1 75		x			• From γ -Be at 80 - 200 GeV: Observed signal of about 60 events above background.
SLAC	{ DAKIN 75 } { MARTIN 75 }		x			• From γ -Be at 14 - 20.5 GeV: Observed 5 events (where 1 event expected)
			x ^b			• From γ -N at 14 - 20.5 GeV
CORNELL	ANDREWS 75		x ^b			• From γ -Be at 11.1 GeV

^a These are only *probable* $\mu^+\mu^-$ events; they are called "collinear events which are not e^+e^- ".

^b Upper limits only.

^c No conclusion drawn, however, on the existence of a narrow resonance.

Data Card Listings

NEW HEAVY MESONS

Evidence bearing on the hadronic nature of the $J/\psi(3100)$ comes from the observation of a large forward photoproduction cross section. Using vector dominance arguments one derives a total cross section for $J/\psi(3100)$ on nucleons over a large energy range of the order of 1 mb (ANDREWS 75, DAKIN 75, KNAPP 75, MARTIN 75). Although this cross section is an order of magnitude smaller than the corresponding cross sections of the well-known vector mesons, it is large enough to suggest that the $J/\psi(3100)$ is probably a hadron.

A preliminary assignment of $I^G(J^P)C = 0^-(1^-)-$ may be made based on the following arguments. A study of the decay angular distribution into lepton pairs (BRAUNSCHWEIG 74, LYNCH 75), and an observed 3-standard-deviation interference effect with QED in the $\mu^+\mu^-$ decay mode (LYNCH 75) suggest that the $J/\psi(3100)$ has the same JPC quantum numbers as a photon, provided parity is conserved in this decay. Non-observation of 2γ decays (BRAUNSCHWEIG 75, FORD 75) supports this interpretation. The observation of a $\Lambda\bar{\Lambda}$ decay mode (SLAC-LBL collaboration, private communication), even though the branching fraction appears to be small, indicates isospin zero, provided the $J/\psi(3100)$ is a hadron and the decay to $\Lambda\bar{\Lambda}$ is not electromagnetic. If $I=0$ and $C=-1$, G-parity must be negative. Support for this assignment comes from the fact (LYNCH 75) that the relative rates for the decay into an odd vs even number of pions is consistent with the assumption that G is conserved in the former and violated in the latter. A natural interpretation, not yet confirmed, is that the $J/\psi(3100)$ is produced in e^+e^- annihilation via a single time-like photon in the direct channel.

The second particle, the $\psi(3700)$, has been observed in e^+e^- collisions at SPEAR (ABRAMS 74) and at DORIS (CRIGEE 75), with cross sections of the same order of magnitude as the $J/\psi(3100)$. It is therefore likely that its formation also takes place via one-photon annihilation, and its quantum numbers would therefore be those of the photon, $J^{PC} = 1^{--}$. No direct evidence for this assignment exists yet. The decay $\psi(3700) \rightarrow J/\psi(3100) + 2\pi$ (LYNCH 75, ABRAMS 75) suggests that the G-parity is the same as for the $J/\psi(3100)$. If $GC = +1$, then I is even.

The $X(4100)$ has been observed as a broad enhancement in the e^+e^- total cross section at SPEAR (AUGUSTIN 75). Although the interpretation is still unclear, if it is in fact a resonance, then its large width

suggests that it is hadronic. Further, as in the case of the $\psi(3700)$, if one-photon annihilation is responsible for the effect, the quantum numbers would then be those of the photon, $J^{PC} = 1^{--}$.

Note added in proof: after this mini-review was completed, an additional paper, BOYARSKI 75, was received. The results of this paper, which support the above preliminary assignments of the $J/\psi(3100)$ quantum numbers, are included in the following Listings.

Extracting Resonance Widths from e^+e^- Colliding Beam Formation Experiments

In an e^+e^- colliding beam formation experiment, the true shape of an observed resonance is distorted primarily by the effects of (1) soft-photon processes, and (2) beam energy spread due to processes such as quantum fluctuations in emission of synchrotron radiation. The spread in energy due to (2) may usually be approximated by a Gaussian distribution, the effect of which vanishes rapidly at energies sufficiently removed from resonance. The major effect of (1) is a decrease in the effective c.m. energy for some fraction of the collisions, because of the emission of bremsstrahlung by the electron or positron before annihilation. Hence, though the nominal beam energy may be well above the resonance region, a certain fraction of the collisions occur at or near resonance. This gives rise to the well-known high-mass radiative tails of the $J/\psi(3100)$ and $\psi(3700)$ resonances. These radiative processes, taken all together, can decrease the peak cross section for a narrow resonance by 50% or more, depending on the mass and the observed width.

Because of these effects, perhaps the most reliable means for determining resonance widths is to use a method based on the area under the line shape. This method, familiar in nuclear physics, minimizes the sensitivity to the details of the beam energy spread. Corrections for the radiative processes, which depend on the limits of integration of the areas, still need to be made. This discussion assumes the resolution is adequate to allow a reasonable separation of signal from background (which itself is subject to radiative processes).

For formation of a resonance of mass M in e^+e^- collisions, with subsequent decay via channel i ,

$$\sigma_i(W) = \sigma_0 \frac{\Gamma_e \Gamma_i / 4}{(M - W)^2 + \Gamma^2 / 4}$$

NEW HEAVY MESONS

where W is the total center of mass energy; Γ , Γ_e , and Γ_i are the total width and partial widths for coupling to e^+e^- and channel i , respectively; and a Breit-Wigner line shape with energy-independent partial widths is assumed. The quantity σ_0 is given by

$$\sigma_0(W) = \frac{4\pi(2J+1)}{W^2}$$

where J is the spin of the resonance. For a narrow resonance, the area under the resonant line is given by

$$A_i = \frac{\pi}{2} \frac{\Gamma_e \Gamma_i}{\Gamma} \sigma_0(M)$$

independent of the energy resolution of the apparatus (but after correction for reduction in peak height and radiative tail caused by soft-photon processes). This area is tabulated in the Data Card Listings in the section labeled Integrated Channel Cross Sections. Substituting for σ_0 , we have

$$\frac{\Gamma_e \Gamma_i}{\Gamma} = \frac{M^2}{2(2J+1)\pi^2} A_i$$

Determination of the area for different decay channels thus makes it possible to determine the various partial widths.

This mini-review is based largely on "Notes from the SLAC Theory Workshop on the ψ ", ed. R. Pearson, SLAC-PUB-1515 (1974). We also wish to thank J. D. Jackson of LBL, Y.-S. Tsai of SLAC, and M. Chanowitz of LBL for fruitful discussions.

Note on the Data Card Listings

The closing date for data for the following Listings was March 28, 1975. We recognize that, because of the intense interest in, and activity relating to, the new mesons, some of the values tabulated may already be superseded by the time of publication of this supplement. We hope, nevertheless, that these Listings will at least serve as a starting point for those studying the new particles. As always we urge the reader to refer to the original articles for all details.

The meaning of the columns and the various abbreviations appearing below can be found in the Illustrative Key in the 1974 edition of the Review. Several new abbreviations for measurement techniques (or detectors) have been introduced this year; they are

Data Card Listings

- ELEC Electronic detector - a combination of chambers, counters, etc.
- DASP DESY double-arm spectrometer
- PLUT DESY PLUTO detector
- SMAG SPEAR magnetic detector
- SPEC Spectrometer

J/ ψ (3100)

70 J/PSI (3100, JPCI CONSISTENT WITH I--0)

70 J/PSI(3100) MASS (MEV)

M	FL	(3100.)	APPROX.	AUBERT	74	SPEC	0	28.	PP(E+E-)	2/75*
M	P	(3105.)	(3.)	AUGUSTIN	74	SMAG	0	3.10	E+E-(+ALL)	2/75*
M	P	(3101.)	(2.)	BALDINI-	74	ELEC	0	3.1	E+E-(+ALL)	2/75*
M	O	(3090.)	4.	BRAUNSCH	74	DASP	0	3.10	E+E-(+E+E-)	2/75*
M	S	(3089.5)		BOYARSKI	75	SMAG	0	3.1	E+E-(+ALL)	3/75*
M	F	(3105.)		CRIGEEE	75	PLUT	0	3.10	E+E-(+ALL)	2/75*
M	F	(3100.)	APPROX.	FORD	75	SPEC	0	3.1	E+E-(2LEPT.)	2/75*
M	F	(3100.)	APPROX.	KNAPPL	75	SPEC	0		GAMMA-BE(MU+MU-)	2/75*
M	F	(3100.)	APPROX.	KNAPP2	75	SPEC	0		N-BE(MU+MU-)	2/75*
M	LP	(3095.)	(5.)	LYNCH	75	SMAG	0	3.10	E+E-(+ALL)	2/75*
M	F	AUGUSTIN 74 AND FORD 75 VALUES ARE NOT INDEPENDENT MASS MEASUREMENTS. THEY RESULT FROM THE SAME BEAM ENERGY CALIBRATION, AND ARE THUS BOTH SUBJECT TO THE RECALIBRATION REFERRED TO IN NOTE L.								
M	L	LYNCH 75 IS A REEVALUATION OF AUGUSTIN 74 BASED ON A RECALIBRATION OF THE SPEAR BEAM ENERGY. SUPERSEDED BY BOYARSKI 75.								
M	P	PRELIMINARY RESULTS.								
M	O	MASS, WIDTH, PARTIAL WIDTHS, AND BRANCHING RATIOS ALL OBTAINED FROM ONE OVERALL FIT TO DATA OF THIS EXPERIMENT.								
M	S	ERROR OF ABOUT 1 PER CENT FROM THE UNCERTAINTY IN CALIBRATION OF THE BEAM ENERGY.								

70 J/PSI(3100) WIDTH (KEV)

THESE VALUES (AS WELL AS THE PARTIAL WIDTHS GIVEN BELOW) ARE BASED ON INTEGRATED CHANNEL CROSS SECTIONS WITH THE ASSUMPTIONS THAT J ψ 1- AND THAT THE TOTAL WIDTH IS THE SUM OF THE HADRON, (E+ E-), AND (MU+ MU-) WIDTHS. UNLESS OTHERWISE NOTED, RADIATIVE CORRECTIONS ARE ASSUMED INCLUDED (SEE JACKSON 74 AND YENNIE 75).

W	BP	(32.1	(12.)	BALDINI-	74	ELEC	0	3.10	E+E-(+ALL)	2/75*
W	O	69.	15.	BOYARSKI	75	SMAG	0	3.1	E+E-(+ALL)	3/75*
W	LP	(77.)	(19.)	LYNCH	75	SMAG	0	3.10	E+E-(+ALL)	2/75*
W	B	CALCULATED BY US ASSUMING THE TOTAL WIDTH TO BE THE SUM OF THEIR HADRON, (E+ E-), AND (MU+ MU-) PARTIAL WIDTHS.								
W	P	PRELIMINARY RESULTS.								
W	O	FROM OVERALL FIT. SEE NOTE ON BOYARSKI 75 MASS.								
W	L	SUPERSEDED BY BOYARSKI 75.								

70 J/PSI(3100) PARTIAL DECAY MODES

P1	J/PSI(3100)	INTO E+ E-	DECAY MASSES
P2	J/PSI(3100)	INTO MU+ MU-	+5+ .5
P3	J/PSI(3100)	INTO HADRONS	105+ 105
P4	J/PSI(3100)	INTO 2 GAMMA	0+ 0
ABOVE MODE FORBIDDEN IF J=1, OR IF C=-1 (AND C IS CONSERVED). SEE BRAUNSCHWEIG 75.			
P5	J/PSI(3100)	INTO PI0 GAMMA	134+ 0
P6	J/PSI(3100)	INTO K+ K-	493+ 493
P7	J/PSI(3100)	INTO PROTON ANTI-PROTON	938+ 938
P8	J/PSI(3100)	INTO K+ K- PI+ PI-	493+ 493+ 139+ 139
P9	J/PSI(3100)	INTO 3PI	139+ 139+ 134
P10	J/PSI(3100)	INTO RHO PI	770+ 139
P11	J/PSI(3100)	INTO 5PI	
P12	J/PSI(3100)	INTO OMEGA PI+ PI-	782+ 139+ 139
P13	J/PSI(3100)	INTO 7PI	
P14	J/PSI(3100)	INTO LAMBDA ANTI-LAMBDA	1115+1115

70 J/PSI(3100) PARTIAL WIDTHS (KEV)

SEE NOTE ABOVE ON TOTAL WIDTH.

W1	J/PSI(3100)	INTO E+ E-	(G1)						
W1 P	(3.4)	(1.2)	BALDINI-	74	ELEC	0	3.1	E+E-(+E+E-)	2/75*
W1 O	4.8	0.6	BOYARSKI	75	SMAG	0	3.1	E+E-(+E+E-)	3/75*
W1 L	(5.2)	(1.3)	LYNCH	75	SMAG	0	3.1	E+E-(+E+E-)	2/75*
W1 P	PRELIMINARY RESULTS NOT INDEPENDENT OF THEIR IC1, IC2, AND IC3 CROSS SECTIONS BELOW.								
W1 O	FROM OVERALL FIT. SEE NOTE ON BOYARSKI 75 MASS.								
W1 L	PRELIMINARY RESULT. SUPERSEDED BY BOYARSKI 75.								
W2	J/PSI(3100)	INTO MU+ MU-	(G2)						
W2 P	(2.4)	(1.1)	BALDINI-	74	ELEC	0	3.1E+E-(MU+MU-)	2/75*	
W2 O	4.8	0.6	BOYARSKI	75	SMAG	0	3.1	E+E-(+E+E-)	3/75*
W2 P	PRELIMINARY RESULTS NOT INDEPENDENT OF THEIR IC1, IC2, AND IC3 CROSS SECTIONS BELOW.								
W2 O	FROM OVERALL FIT. SEE NOTE ON BOYARSKI 75 MASS.								
W3	J/PSI(3100)	INTO HADRONS	(G3)						
W3 P	(25.1)	(11.)	BALDINI-	74	ELEC	0	3.1	E+E-(+HAD.)	2/75*
W3 O	59.	14.	BOYARSKI	75	SMAG	0	3.1	E+E-(+HAD.)	3/75*
W3 P	PRELIMINARY RESULTS NOT INDEPENDENT OF THEIR IC1, IC2, AND IC3 CROSS SECTIONS BELOW.								
W3 O	FROM OVERALL FIT. SEE NOTE ON BOYARSKI 75 MASS.								

Data Card Listings

NEW HEAVY MESONS

Table with columns for particle ID, name, and properties. Section: 70 J/PSI(3100) BRANCHING RATIOS. Includes rows for R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, R12.

Table with columns for particle ID, name, and properties. Section: 70 J/PSI(3100) INTEGRATED CHANNEL CROSS SECTIONS. Includes text for a discussion of this quantity and rows for IC1, IC2, IC3, IC4, IC5.

Table with columns for particle ID, name, and properties. Section: REFERENCES FOR J/PSI(3100). Includes names like CHRISTEN, ABRAMS, ASH, AUBERT, AUGUSTIN, BACCI, etc.

Table with columns for particle ID, name, and properties. Section: 71 PSI(3700) MASS (MEV). Includes rows for M L, M S, M LP, M L, M S, M P.

Table with columns for particle ID, name, and properties. Section: 71 PSI(3700) PARTIAL DECAY MODES. Includes rows for P1, P2, P3, P4, P5, P6.

Table with columns for particle ID, name, and properties. Section: 71 PSI(3700) PARTIAL WIDTHS (KEV). Includes rows for W1, W1 P, W1 P.

Table with columns for particle ID, name, and properties. Section: 71 PSI(3700) BRANCHING RATIOS. Includes rows for R1, R1, R2, R2, R3, R3, R3.

Table with columns for particle ID, name, and properties. Section: REFERENCES FOR PSI(3700). Includes names like ABRAMS, JACKSON, AUBERT, CRIGEE, LYNCH.

Table with columns for particle ID, name, and properties. Section: 72 X(4100) MASS (MEV). Includes row for M (415.0).

Table with columns for particle ID, name, and properties. Section: REFERENCES FOR X(4100). Includes name AUGUSTIN.

