

$\chi_{c1}(3872)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to $< 4\%$ at 95% CL.

See the review on "Spectroscopy of Mesons Containing Two Heavy Quarks."

 $\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

| VALUE (MeV) | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|------------|-----------------------------|-----------|---|
| 3871.69 ± 0.17 OUR AVERAGE | | | | |
| $3871.9 \pm 0.7 \pm 0.2$ | 20 ± 5 | ABLIKIM | 14 BES3 | $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$ |
| $3871.95 \pm 0.48 \pm 0.12$ | 0.6k | AAIJ | 12H LHCb | $pp \rightarrow J/\psi\pi^+\pi^-X$ |
| $3871.85 \pm 0.27 \pm 0.19$ | ~ 170 | ¹ CHOI | 11 BELL | $B \rightarrow K\pi^+\pi^- J/\psi$ |
| $3873 \begin{smallmatrix} +1.8 \\ -1.6 \end{smallmatrix} \pm 1.3$ | 27 ± 8 | ² DEL-AMO-SA.10B | BABR | $B \rightarrow \omega J/\psi K$ |
| $3871.61 \pm 0.16 \pm 0.19$ | 6k | ^{2,3} AALTONEN | 09AU CDF2 | $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$ |
| $3871.4 \pm 0.6 \pm 0.1$ | 93.4 | AUBERT | 08Y BABR | $B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$ |
| $3868.7 \pm 1.5 \pm 0.4$ | 9.4 | AUBERT | 08Y BABR | $B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$ |
| $3871.8 \pm 3.1 \pm 3.0$ | 522 | ^{2,4} ABAZOV | 04F D0 | $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $3873.3 \pm 1.1 \pm 1.0$ | 45 | ⁵ ABLIKIM | 19V BES | $e^+e^- \rightarrow \gamma\omega J/\psi$ |
| 3860.0 ± 10.4 | 13.6 | ^{2,6} AGHASYAN | 18A COMP | $\gamma^* N \rightarrow X\pi^\pm N'$ |
| $3868.6 \pm 1.2 \pm 0.2$ | 8 | ⁷ AUBERT | 06 BABR | $B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$ |
| $3871.3 \pm 0.6 \pm 0.1$ | 61 | ⁷ AUBERT | 06 BABR | $B^- \rightarrow K^- J/\psi\pi^+\pi^-$ |
| 3873.4 ± 1.4 | 25 | ⁸ AUBERT | 05R BABR | $B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$ |
| $3871.3 \pm 0.7 \pm 0.4$ | 730 | ^{2,9} ACOSTA | 04 CDF2 | $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$ |
| $3872.0 \pm 0.6 \pm 0.5$ | 36 | ¹⁰ CHOI | 03 BELL | $B \rightarrow K\pi^+\pi^- J/\psi$ |
| 3836 ± 13 | 58 | ^{2,11} ANTONIAZZI | 94 E705 | $300 \pi^\pm \text{Li} \rightarrow J/\psi\pi^+\pi^-X$ |

¹The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

²Width consistent with detector resolution.

- ³ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/c² is excluded at 95% CL.
⁴ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.
⁵ Fit with fixed width and including two resonances, X(3915) and X(3960).
⁶ Could be a different state.
⁷ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.
⁸ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.
⁹ Superseded by AALTONEN 09AU.
¹⁰ Superseded by CHOI 11.
¹¹ A lower mass value can be due to an incorrect momentum scale for soft pions.

$\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|------------------------|----------|---------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $3872.9^{+0.6+0.4}_{-0.4-0.5}$ | 50 | 1,2 AUSHEV | 10 BELL | $B \rightarrow \bar{D}^{*0} D^0 K$ |
| $3875.1^{+0.7}_{-0.5} \pm 0.5$ | 33 ± 6 | ² AUBERT | 08B BABR | $B \rightarrow \bar{D}^{*0} D^0 K$ |
| $3875.2 \pm 0.7^{+0.9}_{-1.8}$ | 24 ± 6 | ^{2,3} GOKHROO | 06 BELL | $B \rightarrow D^0 \bar{D}^0 \pi^0 K$ |

- ¹ Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.
² Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.
³ Superseded by AUSHEV 10.

$m_{\chi_{c1}(3872)} - m_{J/\psi}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|--------|---|
| $774.9 \pm 3.1 \pm 3.0$ | 522 | ABAZOV | 04F D0 | $p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$ |

$m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|----------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 187.4 ± 1.4 | 25 | ¹ AUBERT | 05R BABR | $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$ |

- ¹ Superseded by AUBERT 06.

$\chi_{c1}(3872)$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------------|----------|---|
| <1.2 | 90 | | CHOI | 11 BELL | $B \rightarrow K \pi^+ \pi^- J/\psi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <2.4 | 90 | | ABLIKIM | 14 BES3 | $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$ |
| <3.3 | 90 | | AUBERT | 08Y BABR | $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$ |
| <4.1 | 90 | 69 | AUBERT | 06 BABR | $B \rightarrow K \pi^+ \pi^- J/\psi$ |
| <2.3 | 90 | 36 | ¹ CHOI | 03 BELL | $B \rightarrow K \pi^+ \pi^- J/\psi$ |

- ¹ Superseded by CHOI 11.

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------------|---------------------|------|---|
| $3.9^{+2.8+0.2}_{-1.4-1.1}$ | 50 | ¹ AUSHEV | 10 | BELL $B \rightarrow \bar{D}^{*0} D^0 K$ |
| $3.0^{+1.9}_{-1.4} \pm 0.9$ | 33 ± 6 | AUBERT | 08B | BABR $B \rightarrow \bar{D}^{*0} D^0 K$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

 $\chi_{c1}(3872)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| Γ_1 $e^+ e^-$ | |
| Γ_2 $\pi^+ \pi^- J/\psi(1S)$ | > 3.2 % |
| Γ_3 $\rho^0 J/\psi(1S)$ | |
| Γ_4 $\omega J/\psi(1S)$ | > 2.3 % |
| Γ_5 $D^0 \bar{D}^0 \pi^0$ | >40 % |
| Γ_6 $\bar{D}^{*0} D^0$ | >30 % |
| Γ_7 $\gamma\gamma$ | |
| Γ_8 $D^0 \bar{D}^0$ | |
| Γ_9 $D^+ D^-$ | |
| Γ_{10} $\gamma \chi_{c1}$ | |
| Γ_{11} $\gamma \chi_{c2}$ | |
| Γ_{12} $\pi^0 \chi_{c2}$ | |
| Γ_{13} $\pi^0 \chi_{c1}$ | > 2.8 % |
| Γ_{14} $\pi^0 \chi_{c0}$ | |
| Γ_{15} $\gamma J/\psi$ | > 7×10^{-3} |
| Γ_{16} $\gamma \psi(2S)$ | > 4 % |
| Γ_{17} $\pi^+ \pi^- \eta_c(1S)$ | not seen |
| Γ_{18} $\pi^+ \pi^- \chi_{c1}$ | not seen |
| Γ_{19} $\rho \bar{\rho}$ | not seen |

C-violating decays

Γ_{20} $\eta J/\psi$

 $\chi_{c1}(3872)$ PARTIAL WIDTHS

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_1 |
|------------|-----|-------------|------|---------|------------|
|------------|-----|-------------|------|---------|------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|----------------------|-----|------|--|
| < 4.3 | 90 | ¹ ABLIKIM | 15v | BES3 | $4.0-4.4 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ |
| <280 | 90 | ² YUAN | 04 | RVUE | $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ |

¹ABLIKIM 15v reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-) / \Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) / \Gamma = 3\%$.

² Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. Assuming that $\Gamma(\pi^+\pi^-J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).

$\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^-J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_1/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|----------|--|
| < 0.13 | 90 | ABLIKIM | 15V BES3 | 4.0–4.4 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 6.2 | 90 | ^{1,2} AUBERT | 05D BABR | 10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$ |
| < 8.3 | 90 | ² DOBBS | 05 CLE3 | $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ |
| < 10 | 90 | ³ YUAN | 04 RVUE | $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ |

¹ Using $B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.

² Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$.

³ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.

$\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^-J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_7/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------------|---------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 12.9 | 90 | ¹ DOBBS | 05 CLE3 | $e^+e^- \rightarrow \pi^+\pi^-J/\psi\gamma$ |

¹ Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.

$\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_7/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------|-----------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| < 1.7 | 90 | ¹ LEES | 12AD BABR | $e^+e^- \rightarrow e^+e^-\omega J/\psi$ |

¹ Assuming $\chi_{c1}(3872)$ has spin 2.

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_7/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------|-----|-------------|-----------|---|
| < 11.1 | 90 | LEES | 12AE BABR | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$ |

$\chi_{c1}(3872)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}$ Γ_2/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------|---------|---------------------|----------|----------------------------------|
| > 0.032 | 93 ± 17 | ¹ AUBERT | 08Y BABR | $B \rightarrow \chi_{c1}(3872)K$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|------------|---------------------|-----|------|--|
| seen | 151 | ² BALA | 15 | BELL | $B \rightarrow \chi_{c1}(3872) K \pi$ |
| >0.05 | 30 | ³ AUBERT | 05R | BABR | $B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$ |
| >0.05 | 36 ± 7 | ⁴ CHOI | 03 | BELL | $B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$ |

¹ AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

² BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872) K^+ \pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872) K^0 \pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

³ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

⁴ CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] / [B(B^+ \rightarrow \psi(2S) K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S) K^+) = (6.19 \pm 0.22) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$.

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|------------------------------|-------------|-------------------------------------|
| >0.023 | 21 ± 7 | ¹ DEL-AMO-SA..10B | BABR | $B^+ \rightarrow \omega J/\psi K^+$ |

¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872) K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi \omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.

$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_4/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|--|
| 1.1 ± 0.4 OUR AVERAGE | Error includes scale factor of 1.7. | | |
| $1.6^{+0.4}_{-0.3} \pm 0.2$ | ¹ ABLIKIM | 19V | BES $e^+ e^- \rightarrow \gamma \omega J/\psi$ |
| 0.8 ± 0.3 | ² DEL-AMO-SA..10B | BABR | $B \rightarrow \omega J/\psi K$ |

¹ Fit with fixed width and including two resonances, X(3915) and X(3960).

² Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|-------------|----------------------|-------------|--|
| >0.4 | 17 ± 5 | ¹ GOKHROO | 06 | BELL $B^+ \rightarrow D^0 \bar{D}^0 \pi^0 K^+$ |

¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

$\Gamma(D^0 \bar{D}^0 \pi^0) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_5 / Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|------|-------------------------|------|---------------------------------------|
| seen | ¹ GOKHROO 06 | BELL | $B \rightarrow D^0 \bar{D}^0 \pi^0 K$ |
|------|-------------------------|------|---------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|------|-----------|------|---------------------------------------|
| seen | AUSHEV 10 | BELL | $B \rightarrow D^0 \bar{D}^0 \pi^0 K$ |
|------|-----------|------|---------------------------------------|

¹ May not necessarily be the same state as that observed in the $J/\psi \pi^+ \pi^-$ mode. Supersedes CHISTOV 04.

 $\Gamma(\bar{D}^{*0} D^0) / \Gamma_{\text{total}}$ Γ_6 / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

| | | | | |
|-------|----------------|------------------------|------|--|
| >0.30 | 41^{+9}_{-8} | ¹ AUSHEV 10 | BELL | $B^+ \rightarrow D^{*0} \bar{D}^0 K^+$ |
|-------|----------------|------------------------|------|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|------------|-------------------------|------|--|
| >0.6 | 27 ± 6 | ² AUBERT 08B | BABR | $B^+ \rightarrow \bar{D}^{*0} D^0 K^+$ |
|------|------------|-------------------------|------|--|

¹ AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

² AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

 $\Gamma(D^0 \bar{D}^0) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_8 / Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|----------|------------|------|---------------------------------|
| not seen | CHISTOV 04 | BELL | $B \rightarrow K D^0 \bar{D}^0$ |
|----------|------------|------|---------------------------------|

 $\Gamma(D^+ D^-) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_9 / Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|----------|------------|------|---------------------------|
| not seen | CHISTOV 04 | BELL | $B \rightarrow K D^+ D^-$ |
|----------|------------|------|---------------------------|

 $\Gamma(\gamma \chi_{c1}) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{10} / Γ_2

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|----------|--|--------------------------|------|--|
| not seen | | ¹ BHARDWAJ 13 | BELL | $B^+ \rightarrow \chi_{c1} \gamma K^+$ |
|----------|--|--------------------------|------|--|

| | | | | |
|-------|----|---------|------|--------------------------------------|
| <0.89 | 90 | CHOI 03 | BELL | $B \rightarrow K \pi^+ \pi^- J/\psi$ |
|-------|----|---------|------|--------------------------------------|

¹ Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1}) < 1.9 \times 10^{-6}$ at 90% CL.

 $\Gamma(\gamma \chi_{c2}) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{11} / Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|----------|--------------------------|------|--|
| not seen | ¹ BHARDWAJ 13 | BELL | $B^\pm \rightarrow \chi_{c2} \gamma K^\pm$ |
|----------|--------------------------|------|--|

¹ Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2}) < 6.7 \times 10^{-6}$ at 90% CL.

$\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

>7 × 10⁻³ ¹ BHARDWAJ 11 BELL $B^\pm \rightarrow \gamma J/\psi K^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.011 20 ² AUBERT 09B BABR $B^+ \rightarrow \gamma J/\psi K^+$

>0.013 19 ³ AUBERT,BE 06M BABR $B^+ \rightarrow \gamma J/\psi K^+$

¹ BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

³ Superseded by AUBERT 09B. AUBERT,BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

$\Gamma(\gamma\psi(2S))/\Gamma_{\text{total}}$ Γ_{16}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

seen 36 ± 9 ¹ AAIJ 14AH LHCB $B^+ \rightarrow \gamma\psi(2S) K^+$

>0.04 25 ± 7 ² AUBERT 09B BABR $B^+ \rightarrow \gamma\psi(2S) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ³ BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S) K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ.

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

³ BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma\psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$ Γ_{16}/Γ_{15}

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|-----|------|-------------|------|---------|
|-------|-----|------|-------------|------|---------|

2.6 ± 0.6 OUR AVERAGE

2.46 ± 0.64 ± 0.29 36 ± 9 ¹ AAIJ 14AH LHCB $B^+ \rightarrow \gamma\psi(2S) K^+$

3.4 ± 1.4 AUBERT 09B BABR $B^+ \rightarrow \gamma c\bar{c} K'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1 90 BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S) K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ.

$\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$ Γ_{18}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

not seen ¹ BHARDWAJ 16 BELL $B^+ \rightarrow \pi^+ \pi^- \chi_{c1} K^+$

¹ BHARDWAJ 16 quotes $B(B^+ \rightarrow \chi_{c1}(3872) K^+) \cdot B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \chi_{c1}) < 1.5 \times 10^{-6}$ at 90% CL.

| $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{19}/Γ |
|--|-----|-------------------|-----------|--|----------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| not seen | | ¹ AAIJ | 17AD LHCb | $pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$ | |
| ¹ AAIJ 17AD reports $B(B^+ \rightarrow \chi_{c1}(3872)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) < 2.0 (2.5) \times 10^{-3}$ at 90% (95%) CL. | | | | | |

| $\Gamma(p\bar{p})/\Gamma(\pi^+\pi^- J/\psi(1S))$ | | | | | Γ_{19}/Γ_2 |
|---|-----|-------------------|----------|-------------------------------|------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <2.0 × 10⁻³ | 95 | ¹ AAIJ | 13S LHCb | $B^+ \rightarrow p\bar{p}K^+$ | |
| ¹ AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$. | | | | | |

| $\Gamma(\pi^0\chi_{c0})/\Gamma(\pi^+\pi^- J/\psi(1S))$ | | | | | Γ_{14}/Γ_2 |
|--|-----|-------------|----------|--|------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <19 | 90 | ABLIKIM | 19U BES3 | $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$ | |

| $\Gamma(\pi^0\chi_{c1})/\Gamma(\pi^+\pi^- J/\psi(1S))$ | | | | | Γ_{13}/Γ_2 |
|--|------|------|-------------|----------|--|
| VALUE (units 10 ⁻²) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 88⁺³³₋₂₇ ± 10 | 10.8 | | ABLIKIM | 19U BES3 | $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---|----|-----------------------|---------|--|--|
| <97 | 90 | ¹ BHARDWAJ | 19 BELL | $B^\pm \rightarrow \chi_{c1}\pi^0 K^\pm$ | |
| ¹ BHARDWAJ 19 reports $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \pi^0\chi_{c1}) < 8.1 \times 10^{-6}$ at 90% CL which was divided by $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) = (8.63 \pm 0.97) \times 10^{-6}$ from CHOI 11. | | | | | |

| $\Gamma(\pi^0\chi_{c2})/\Gamma(\pi^+\pi^- J/\psi(1S))$ | | | | | Γ_{12}/Γ_2 |
|--|-----|-------------|----------|--|------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <1.1 | 90 | ABLIKIM | 19U BES3 | $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$ | |

————— C-violating decays —————

| $\Gamma(\eta J/\psi)/\Gamma(\pi^+\pi^- J/\psi(1S))$ | | | | | Γ_{20}/Γ_2 |
|---|-----|-------------------------|----------|------------------------------|------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.4 | 90 | ^{1,2} IWASHITA | 14 BELL | $B \rightarrow K\eta J/\psi$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.6 | 90 | AUBERT | 04Y BABR | $B \rightarrow K\eta J/\psi$ | |
| ¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$. | | | | | |
| ² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals. | | | | | |

$\chi_{c1}(3872)$ REFERENCES

| | | | | |
|---------------|------|------------------|----------------------------------|-------------------|
| ABLIKIM | 19U | PRL 122 202001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19V | PRL 122 232002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| BHARDWAJ | 19 | PR D99 111101 | V. Bhardwaj <i>et al.</i> | (BELLE Collab.) |
| AGHASYAN | 18A | PL B783 334 | M. Aghasyan <i>et al.</i> | (COMPASS Collab.) |
| AAIJ | 17AD | PL B769 305 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| BHARDWAJ | 16 | PR D93 052016 | V. Bhardwaj <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 15AO | PR D92 011102 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 15V | PL B749 414 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| BALA | 15 | PR D91 051101 | A. Bala <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 14AH | NP B886 665 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 14 | PRL 112 092001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| IWASHITA | 14 | PTEP 2014 043C01 | T. Iwashita <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 13Q | PRL 110 222001 | R. Aaij <i>et al.</i> | (LHCb Collab.) JP |
| AAIJ | 13S | EPJ C73 2462 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| BHARDWAJ | 13 | PRL 111 032001 | V. Bhardwaj <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 12H | EPJ C72 1972 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| LEES | 12AD | PR D86 072002 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 12AE | PR D86 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| BHARDWAJ | 11 | PRL 107 091803 | V. Bhardwaj <i>et al.</i> | (BELLE Collab.) |
| CHOI | 11 | PR D84 052004 | S.-K. Choi <i>et al.</i> | (BELLE Collab.) |
| AUSHEV | 10 | PR D81 031103 | T. Aushev <i>et al.</i> | (BELLE Collab.) |
| DEL-AMO-SA... | 10B | PR D82 011101 | P. del Amo Sanchez <i>et al.</i> | (BABAR Collab.) |
| AALTONEN | 09AU | PRL 103 152001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AUBERT | 09B | PRL 102 132001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 08B | PR D77 011102 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 08Y | PR D77 111101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06 | PR D73 011101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06M | PR D74 071101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| GOKHROO | 06 | PRL 97 162002 | G. Gokhroo <i>et al.</i> | (BELLE Collab.) |
| AUBERT | 05B | PR D71 031501 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05D | PR D71 052001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05R | PR D71 071103 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| DOBBS | 05 | PRL 94 032004 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABAZOV | 04F | PRL 93 162002 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ACOSTA | 04 | PRL 93 072001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| AUBERT | 04Y | PRL 93 041801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| CHISTOV | 04 | PRL 93 051803 | R. Chistov <i>et al.</i> | (BELLE Collab.) |
| PDG | 04 | PL B592 1 | S. Eidelman <i>et al.</i> | (PDG Collab.) |
| YUAN | 04 | PL B579 74 | C.Z. Yuan <i>et al.</i> | |
| CHOI | 03 | PRL 91 262001 | S.-K. Choi <i>et al.</i> | (BELLE Collab.) |
| BAI | 98E | PR D57 3854 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ANTONIAZZI | 94 | PR D50 4258 | L. Antoniazzi <i>et al.</i> | (E705 Collab.) |