

$\eta_c(2S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.

### $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3637.5±1.1 OUR AVERAGE</b>				Error includes scale factor of 1.2.
3635.1±3.7±2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6±1.7±0.6	106	<sup>1</sup> AAIJ	17ADLHCb	$p p \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4±4.1±0.7	365	<sup>2</sup> AAIJ	17BBLHCb	$p p \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-)X$
3637.0±5.7±3.4	178	<sup>3,4</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^0$
3635.1±5.8±2.1	47	<sup>3,5</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta$
3646.9±1.6±3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6±2.9±1.6	127 ± 18	<sup>6</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, K K\pi^0$
3638.5±1.5±0.8	624	<sup>3</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5±3.2±2.5	1201	<sup>3</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 <sup>+3.9</sup> <sub>-4.2</sub> <sup>+0.7</sup> <sub>-2.0</sub>	128	<sup>7</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	<sup>8</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0±5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9±3.1±1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3639 ± 7	98 ± 52	<sup>9</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$
3630.8±3.4±1.0	112 ± 24	<sup>10</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ± 6 ± 8	39 ± 11	<sup>11</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		<sup>12</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup>AAIJ 17AD report  $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$  MeV. We use the current value  $m_{\psi(2S)} = 3686.097 \pm 0.025$  MeV to obtain the quoted mass.

<sup>2</sup>From a fit of the  $\phi\phi$  invariant mass with the width of  $\eta_c(2S)$  fixed to the PDG 16 value.

<sup>3</sup>Ignoring possible interference with continuum.

<sup>4</sup>With a width fixed to 11.3 MeV.

<sup>5</sup>With a width fixed to 11.3 MeV. Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

<sup>6</sup>From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>7</sup>Accounts for interference with non-resonant continuum.

<sup>8</sup>From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>9</sup>From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>10</sup>Superseded by DEL-AMO-SANCHEZ 11M.

<sup>11</sup>Superseded by VINOKUROVA 11.

<sup>12</sup>Assuming mass of  $\psi(2S) = 3686$  MeV.

## $\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.3<sup>+ 3.2</sup><sub>- 2.9</sub> OUR AVERAGE</b>					
9.9 ± 4.8 ± 2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8		127 ± 18	<sup>13</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi,$ $K K\pi^0$
13.4 ± 4.6 ± 3.2		624	<sup>14</sup> DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
6.6 <sup>+ 8.4 + 2.6</sup> <sub>- 5.1 - 0.9</sub>		128	<sup>15</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 23	90	98 ± 52	<sup>16</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
17.0 ± 8.3 ± 2.5		112 ± 24	<sup>17</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
< 55	90	39 ± 11	<sup>18</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95		<sup>19</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$
<sup>13</sup> From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.					
<sup>14</sup> Ignoring possible interference with continuum.					
<sup>15</sup> Accounts for interference with non-resonant continuum.					
<sup>16</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.					
<sup>17</sup> Superseded by DEL-AMO-SANCHEZ 11M.					
<sup>18</sup> For a mass value of $3654 \pm 6$ MeV. Superseded by VINOKUROVA 11.					
<sup>19</sup> For a mass value of $3594 \pm 5$ MeV					

## $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons		not seen
$\Gamma_2$ $K\bar{K}\pi$	( 1.9 ± 1.2 ) %	
$\Gamma_3$ $K\bar{K}\eta$	( 5 ± 4 ) × 10 <sup>-3</sup>	
$\Gamma_4$ $2\pi^+ 2\pi^-$	not seen	
$\Gamma_5$ $\rho^0 \rho^0$	not seen	
$\Gamma_6$ $3\pi^+ 3\pi^-$	not seen	
$\Gamma_7$ $K^+ K^- \pi^+ \pi^-$	not seen	
$\Gamma_8$ $K^{*0} \bar{K}^{*0}$	not seen	
$\Gamma_9$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.4 ± 1.0 ) %	
$\Gamma_{10}$ $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
$\Gamma_{11}$ $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	seen	
$\Gamma_{12}$ $2K^+ 2K^-$	not seen	
$\Gamma_{13}$ $\phi\phi$	not seen	

$\Gamma_{14}$	$p\bar{p}$	seen		
$\Gamma_{15}$	$p\bar{p}\pi^+\pi^-$	seen		
$\Gamma_{16}$	$\gamma\gamma$	$(1.9 \pm 1.3) \times 10^{-4}$		
$\Gamma_{17}$	$\gamma J/\psi(1S)$	$< 1.4 \quad \%$	90%	
$\Gamma_{18}$	$\pi^+\pi^-\eta$	not seen		
$\Gamma_{19}$	$\pi^+\pi^-\eta'$	not seen		
$\Gamma_{20}$	$\pi^+\pi^-\eta_c(1S)$	$< 25 \quad \%$	90%	

 **$\eta_c(2S)$  PARTIAL WIDTHS** **$\Gamma(\gamma\gamma)$**  **$\Gamma_{16}$** 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.44 \pm 0.14$	106	20 XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$1.3 \pm 0.6$		21 ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

20 Assuming that the branching fraction into  $\eta'\pi^+\pi^-$  is the same as for  $\eta_c(1S)$ .21 They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV. **$\Gamma(\gamma\gamma) \times \Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$**  **$\Gamma_{16}\Gamma_{19}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5.6^{+1.2}_{-1.1} \pm 1.1$	106	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

 **$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**  **$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_4\Gamma_{16}/\Gamma$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.5$	90	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

 **$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_2\Gamma_{16}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$41^{+4}_{-6} \pm 6$	624	22 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

22 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 **$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_7\Gamma_{16}/\Gamma$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.0$	90	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

 **$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_9\Gamma_{16}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$30^{+6}_{-5} \pm 5$	1201	23 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

23 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{12}\Gamma_{16}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.9	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$
$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{20}\Gamma_{16}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<133	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$	

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$
VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
< 5.6	90	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 8.0	90	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
<12.0	90	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
24 Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.					
25 For a total width $\Gamma=5$ MeV.					
26 For the resonance mass region $3589$ – $3599$ MeV/ $c^2$ .					
27 For the resonance mass region $3575$ – $3660$ MeV/ $c^2$ .					

### $\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABREU	980 DLPH	$e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
seen	28 EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$		
28 For a mass value of $3594 \pm 5$ MeV					

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.9±0.4±1.1</b>	$59 \pm 12$	29 AUBERT	08AB BABR	$B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
seen	$127 \pm 18$	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$	
seen	$39 \pm 11$	30 CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$	
29 Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S) K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S) K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .					
30 For a mass value of $3654 \pm 6$ MeV					

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$					$\Gamma_3/\Gamma_2$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>27.3±7.0±9.0</b>	225	31 LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$	
31 LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$ , which we divide by 3 to account for isospin symmetry.					

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

 $\Gamma_4/\Gamma$  $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

 $\Gamma_5/\Gamma$  $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

 $\Gamma_7/\Gamma$  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.73 ± 0.17 ± 0.17</b>	1201	32 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>32</sup> We have multiplied the value of  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 $\Gamma_9/\Gamma_2$  $\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

 $\Gamma_8/\Gamma$  $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	57 ± 17	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

 $\Gamma_{11}/\Gamma$  $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

 $\Gamma_{12}/\Gamma$  $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

 $\Gamma_{13}/\Gamma$  $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	106	33 AAIJ	17AD LHCb	$p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$

 $\Gamma_{14}/\Gamma$ 

<sup>33</sup> AAIJ 17AD report a 6.4 standard deviation signal, with  $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$ .

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	110	34 CHILIKIN	19	BELL $e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma_{15}/\Gamma$ 

<sup>34</sup> CHILIKIN 19 reports signals in  $B^+ \rightarrow \eta_c(2S) K^+$  and  $B^0 \rightarrow \eta_c(2S) K_S^0$  with 12.3 and 5.9 standard deviations, respectively.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<4 \times 10^{-4}$	90	35 WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma\gamma$	
not seen		AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
$<0.01$	90	LEE	85	CBAL $\psi' \rightarrow \text{photons}$	
35 WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$ .					

$\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$					$\Gamma_{20}/\Gamma_2$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;3.33</b>	90	36 LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$	

36 We divided the reported limit by 3 to take into account isospin relations.

### $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<11.8 \times 10^{-6}$	90	37 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$	

37 CRONIN-HENNESSY 10 reports a limit of  $< 5.9 \times 10^{-6}$  for the decay  $\eta_c(2S) \rightarrow K^+ K^- \eta$  which we multiply by 2 account for isospin symmetry. It assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. It also gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;14.6 x 10^-6</b>	90	38 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	

38 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<12.7 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	

$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_6/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<13.2 \times 10^{-6}$	90	39 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$	

39 Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}^{\psi(2S)}$$

$$\Gamma_7/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	40 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>40</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_8/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}^{\psi(2S)}$$

$$\Gamma_9/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	41 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>41</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	42 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>42</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{11}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60	ABLIKIM	13K BES3		$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15.2$	90	43 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

<sup>43</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{13}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					
$\Gamma_{14}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					
$\Gamma_{17}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$					
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33	44	ABLIKIM	17N BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$

<sup>44</sup> Uses  $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$ .

$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					
$\Gamma_{18}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.3 \times 10^{-6}$	90	45	CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

<sup>45</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					
$\Gamma_{19}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<14.2 \times 10^{-6}$	90	46	CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>46</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					
$\Gamma_{20}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-4}$	90	47	CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$

<sup>47</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

## $\eta_c(2S)$ REFERENCES

CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA... VINOKUROVA	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN... AUBERT	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)

UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)