

$\omega(1650)$ $I^G(J^{PC}) = 0^-(1^{--})$ See also the $\omega(1420)$ particle listing. **$\omega(1650)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670 ± 30 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1651 \pm 3 $^{+16}_{-6}$	183k	1 ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
1673 $^{+6}_{-7}$		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
1671 \pm 6 \pm 10	824	2 AKHMETSHIN	17A CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
1660 \pm 10	898	3 ACHASOV	16B SND	1.34–2.00 $e^+ e^- \rightarrow \omega \eta$
1680 \pm 10	13.1k	4 AULCHENKO	15A SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1667 \pm 13 \pm 6		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
1645 \pm 8	13	AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \omega \eta \gamma$
1660 \pm 10 \pm 2		AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1770 \pm 50 \pm 60	1.2M	5 ACHASOV	03D RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1619 \pm 5		6 HENNER	02 RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
1700 \pm 20		EUGENIO	01 SPEC	18 $\pi^- p \rightarrow \omega \eta n$
1705 \pm 26	612	7 AKHMETSHIN	00D CMD2	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1820 $^{+190}_{-150}$		8 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1840 $^{+100}_{-70}$		9 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1780 $^{+170}_{-300}$		10 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K^+ K^-$
\sim 2100		11 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606 \pm 9		12 CLEGG	94 RVUE	
1662 \pm 13	750	13 ANTONELLI	92 DM2	1.34–2.4 $e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
1670 \pm 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657 \pm 13		CORDIER	81 DM1	$e^+ e^- \rightarrow \omega 2\pi$
1679 \pm 34	21	ESPOSITO	80 FRAM	$e^+ e^- \rightarrow 3\pi$
1652 \pm 17		COSME	79 OSPK	$e^+ e^- \rightarrow 3\pi$

¹ Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X \pi^0 \rightarrow K^+ K^- \pi^0 = (5.3 \pm 0.3)^{+0.6}_{-0.5} \times 10^{-5}$.

² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

³ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

⁴ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

⁵ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁶ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

- ⁷ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
⁸ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.
⁹ Using the data from ANTONELLI 92.
¹⁰ Using the data from IVANOV 81 and BISELLO 88B.
¹¹ Using the data from BISELLO 91C.
¹² From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
¹³ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

$\omega(1650)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315 ± 35 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
194 \pm 8 \pm 15 7	183k	¹ ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
95 \pm 11		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
113 \pm 9 \pm 10	824	² AKHMETSHIN	17A CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
110 \pm 20	898	³ ACHASOV	16B SND	1.34–2.00 $e^+ e^- \rightarrow \omega \eta$
310 \pm 30	13.1k	⁴ AULCHENKO	15A SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
222 \pm 25 \pm 20		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
114 \pm 14	13	AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \omega \eta \gamma$
230 \pm 30 \pm 20		AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
490 $^{+200}_{-150}$ \pm 130	1.2M	⁵ ACHASOV	03D RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
250 \pm 14		⁶ HENNER	02 RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
250 \pm 50		EUGENIO	01 SPEC	18 $\pi^- p \rightarrow \omega \eta n$
370 \pm 25	612	⁷ AKHMETSHIN	00D CMD2	1.4–2.0 $e^+ e^- \rightarrow \omega \pi^+ \pi^-$
113 \pm 20		⁸ CLEGG	94 RVUE	
280 \pm 24	750	⁹ ANTONELLI	92 DM2	1.34–2.4 $e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
160 \pm 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
136 \pm 46		CORDIER	81 DM1	$e^+ e^- \rightarrow \omega 2\pi$
99 \pm 49	21	ESPOSITO	80 FRAM	$e^+ e^- \rightarrow 3\pi$
42 \pm 17		COSME	79 OSPK	$e^+ e^- \rightarrow 3\pi$

¹ Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X \pi^0 \rightarrow K^+ K^- \pi^0 = (5.3 \pm 0.3)^{+0.6}_{-0.5) \times 10^{-5}}$.

² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

³ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

⁴ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

⁵ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁶ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁷ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁸ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁹ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

$\omega(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \omega\pi\pi$	seen
$\Gamma_3 \omega\eta$	seen
$\Gamma_4 e^+e^-$	seen
$\Gamma_5 \pi^0\gamma$	not seen

$$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$$

$$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_4/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
1.56 ± 0.23	13.1k	¹ AULCHENKO	15A SND	$1.05-1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1.3 ± 0.1	± 0.1	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1.2 ± 0.4	± 0.8	1.2M ^{2,3} ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921 ± 0.230		4,5 CLEGG	94 RVUE	
0.479 ± 0.050	750	^{6,7} ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

² Calculated by us from the cross section at the peak.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁵ From the partial and leptonic width given by the authors.

⁶ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

⁷ From the product of the leptonic width and partial branching ratio given by the authors.

$$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma \times \Gamma_4/\Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
7.0 ± 0.5		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 ± 0.9	± 1.3	1.2M ^{1,2} ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		³ AKHMETSHIN 00D	CMD2	$1.2-1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		4,5 CLEGG	94 RVUE	
6.07 ± 0.61	750	^{6,7} ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ Calculated by us from the cross section at the peak.

² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

³ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

- ⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
⁵ From the partial and leptonic width given by the authors.
⁶ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.
⁷ From the product of the leptonic width and partial branching ratio given by the authors.

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_4/\Gamma$				
<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.62 ^{+0.45} _{-0.42}		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$	
4.5 ± 0.3 ± 0.3	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$	
4.4 ± 0.5	898	² ACHASOV	16B	SND 1.34–2.00 $e^+e^- \rightarrow \omega\eta$	
5.7 ± 0.6	13	AUBERT	06D	BABR 10.6 $e^+e^- \rightarrow \omega\eta\gamma$	
< 60 at 90% CL		³ AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$	
¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1650) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1650) \rightarrow e^+e^-) = 51 \pm 3$ eV.					
² From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.					
³ $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.					

$\omega(1650)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	Γ_1/Γ				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 0.65	1.2M	¹ ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
0.380 ± 0.014		² HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$	
¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.					
² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.					

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	Γ_2/Γ				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 0.35	1.2M	¹ ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
0.620 ± 0.014		² HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$	
¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.					
² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.					

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 18	1.2M	1,2 ACHASOV	03D RVUE	$0.44\text{--}2.00 \text{ } e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
32 ± 1		2 HENNER	02 RVUE	$1.2\text{--}2.0 \text{ } e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ Calculated by us from the cross section at the peak.² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only. $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	1 ACHASOV	10D SND	$1.075\text{--}2.0 \text{ } e^+ e^- \rightarrow \pi^0\gamma$
1 From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. The width of the highest mass effective resonance is fixed at 315 MeV.			

 $\omega(1650)$ REFERENCES

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
		Translated from ZETF 148 34.		
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)