


**$\pi_2(1670)$**  $I^G(J^{PC}) = 1^-(2^-+)$  **$\pi_2(1670)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1670.6<math>^{+2.9}_{-1.2}</math> OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.			
1642 $\pm 12$	46M	1 AGHASYAN	18B COMP		$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1749 $\pm 10$	$\pm 100$	145k LU	05 B852		$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
1676 $\pm 3$	$\pm 8$	2 CHUNG	02 B852		$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1685 $\pm 10$	$\pm 30$	BARBERIS	01		$450 pp \rightarrow p_f 3\pi^0 p_s$
1687 $\pm 9$	$\pm 15$	AMELIN	99 VES		$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1669 $\pm 4$		BARBERIS	98B		$450 pp \rightarrow p_f \rho \pi p_s$
1670 $\pm 4$		BARBERIS	98B		$450 pp \rightarrow p_f f_2(1270) \pi p_s$
1690 $\pm 14$		3 BERDNIKOV	94 VES		$37 \pi^- A \rightarrow K^+ K^- \pi^- A$
1710 $\pm 20$	700	ANTIPOV	87 SIGM	-	$50 \pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
1676 $\pm 6$		3 EVANGELIS...	81 OMEG	-	$12 \pi^- p \rightarrow 3\pi p$
1657 $\pm 14$		3,4 DAUM	80D SPEC	-	$63-94 \pi p \rightarrow 3\pi X$
1662 $\pm 10$	2000	3 BALTAY	77 HBC	+	$15 \pi^+ p \rightarrow p 3\pi$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
1658 $\pm 3$	$\pm 24$	420k 5 ALEKSEEV	10 COMP		$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1730 $\pm 20$		6 AMELIN	95B VES		$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1742 $\pm 31$	$\pm 49$	ANTREASYAN	90 CBAL		$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
1624 $\pm 21$		2 BELLINI	85 SPEC		$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1622 $\pm 35$		7 BELLINI	85 SPEC		$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1693 $\pm 28$		8 BELLINI	85 SPEC		$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1710 $\pm 20$		9 DAUM	81B SPEC	-	$63,94 \pi^- p$
1660 $\pm 10$		3 ASCOLI	73 HBC	-	$5-25 \pi^- p \rightarrow p \pi_2$

<sup>1</sup> Statistical error negligible.<sup>2</sup> From  $f_2(1270)\pi$  decay.<sup>3</sup> From a fit to  $J^P = 2^-$  S-wave  $f_2(1270)\pi$  partial wave.<sup>4</sup> Clear phase rotation seen in  $2^- S$ ,  $2^- P$ ,  $2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.<sup>5</sup> Superseded by AGHASYAN 2018B.<sup>6</sup>  $J^{PC}$  ambiguous.<sup>7</sup> From  $\rho\pi$  decay.

<sup>8</sup> From  $\sigma\pi$  decay.

<sup>9</sup> From a two-resonance fit to four  $2^-0^+$  waves. This should not be averaged with all the single resonance fits.



### $\pi_2(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>258^{+8}_{-9}</math> OUR AVERAGE</b>					Error includes scale factor of 1.2.
$311^{+12}_{-23}$	46M	10 AGHASYAN	18B	COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$408 \pm 60 \pm 250$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$254 \pm 3 \pm 31$		11 CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$265 \pm 30 \pm 40$		BARBERIS	01		$450 pp \rightarrow p_f 3\pi^0 p_s$
$168 \pm 43 \pm 53$		AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
$268 \pm 15$		BARBERIS	98B		$450 pp \rightarrow p_f \rho \pi p_s$
$256 \pm 15$		BARBERIS	98B		$450 pp \rightarrow p_f f_2(1270) \pi p_s$
$190 \pm 50$		12 BERDNIKOV	94	VES	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$
$170 \pm 80$	700	ANTIPOV	87	SIGM	$50 \pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
$260 \pm 20$		12 EVANGELIS...	81	OMEG	$12 \pi^- p \rightarrow 3\pi p$
$219 \pm 20$		12,13 DAUM	80D	SPEC	$63-94 \pi p \rightarrow 3\pi X$
$285 \pm 60$	2000	12 BALTAY	77	HBC	$15 \pi^+ p \rightarrow p 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$271 \pm 9^{+22}_{-24}$	420k	<sup>14</sup> ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$310 \pm 20$		<sup>15</sup> AMELIN	95B	VES	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
$236 \pm 49 \pm 36$		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
$304 \pm 22$		<sup>11</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$404 \pm 108$		<sup>16</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$330 \pm 90$		<sup>17</sup> BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
$312 \pm 50$		<sup>18</sup> DAUM	81B	SPEC	$63,94 \pi^- p$
$270 \pm 60$		<sup>12</sup> ASCOLI	73	HBC	$5-25 \pi^- p \rightarrow p \pi_2^-$

<sup>10</sup> Statistical error negligible.

<sup>11</sup> From  $f_2(1270)\pi$  decay.

<sup>12</sup> From a fit to  $J^P = 2^- f_2(1270)\pi$  partial wave.

<sup>13</sup> Clear phase rotation seen in  $2^- S$ ,  $2^- P$ ,  $2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.

<sup>14</sup> Superseded by AGHASYAN 2018B.

<sup>15</sup>  $JPC$  ambiguous.

<sup>16</sup> From  $\rho\pi$  decay.

<sup>17</sup> From  $\sigma\pi$  decay.

<sup>18</sup> From a two-resonance fit to four  $2^- 0^+$  waves. This should not be averaged with all the single resonance fits.

## $\pi_2(1670)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 3\pi$	(95.8±1.4) %	
$\Gamma_2 \pi^+ \pi^- \pi^0$		
$\Gamma_3 \pi^0 \pi^0 \pi^0$		
$\Gamma_4 f_2(1270)\pi$	(56.3±3.2) %	
$\Gamma_5 \rho\pi$	(31 ± 4) %	
$\Gamma_6 \sigma\pi$	(10 ± 4) %	
$\Gamma_7 \pi(\pi\pi)_{S\text{-wave}}$	( 8.7±3.4) %	
$\Gamma_8 \pi^\pm \pi^+ \pi^-$	(53 ± 4) %	
$\Gamma_9 K\bar{K}^*(892) + \text{c.c.}$	( 4.2±1.4) %	
$\Gamma_{10} \omega\rho$	( 2.7±1.1) %	
$\Gamma_{11} \pi^\pm \gamma$	( 7.0±1.2) × 10 <sup>-4</sup>	
$\Gamma_{12} \gamma\gamma$	< 2.8 × 10 <sup>-7</sup>	90%
$\Gamma_{13} \eta\pi$	< 5 %	
$\Gamma_{14} \pi^\pm 2\pi^+ 2\pi^-$	< 5 %	
$\Gamma_{15} \rho(1450)\pi$	< 3.6 × 10 <sup>-3</sup>	97.7%
$\Gamma_{16} b_1(1235)\pi$	< 1.9 × 10 <sup>-3</sup>	97.7%
$\Gamma_{17} \eta 3\pi$	possibly seen	
$\Gamma_{18} f_1(1285)\pi$	not seen	
$\Gamma_{19} a_2(1320)\pi$		

## CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 1.9$  for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_5$	-53	
$x_7$	-29	-59
$x_9$	-8	-21
	$x_4$	$x_5$
		$x_7$

## $\pi_2(1670)$ PARTIAL WIDTHS

### $\Gamma(\pi^\pm \gamma)$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_{11}$
<b>181±11±27</b>	19 ADOLPH	14 COMP	-	$190 \pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$	

19 Primakoff reaction. Assumes incoherent  $f_2(1270)\pi$  contribution to  $3\pi$  final state and uses  $B(\pi_2(1670) \rightarrow f_2\pi) = 56\%$ .

### $\Gamma(\gamma\gamma)$

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_{12}$
<b>&lt;0.072</b>	90	20 ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>						
<0.19	90	20 ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
1.41 $\pm 0.23 \pm 0.28$		ANTREASYAN	90	CBAL	0 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$	
0.8 $\pm 0.3 \pm 0.12$		21 BEHREND	90C	CELL	0 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
1.3 $\pm 0.3 \pm 0.2$		22 BEHREND	90C	CELL	0 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	

20 Decaying into  $f_2(1270)\pi$  and  $\rho\pi$ .

21 Constructive interference between  $f_2(1270)\pi, \rho\pi$  and background.

22 Incoherent Ansatz.

## $\pi_2(1670) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

### $\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2 \Gamma_{12} / \Gamma$
<b>&lt;0.1</b>	95	23 SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	

23 From analysis of L3 data at 183–209 GeV.

**$\pi_2(1670)$  BRANCHING RATIOS** $\Gamma(3\pi)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID
<b>0.958±0.014 OUR FIT</b>	

 $\Gamma_1/\Gamma = (\Gamma_4 + \Gamma_5 + \Gamma_7)/\Gamma$  $\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ 

VALUE	DOCUMENT ID	COMMENT
<b>0.29±0.03±0.05</b>	BARBERIS 01	$450 \text{ pp} \rightarrow p_f 3\pi^0 p_s$

 $\Gamma_3/\Gamma_2$  $\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi)$ (With  $f_2(1270) \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.97±0.09 OUR AVERAGE</b>			Error includes scale factor of 1.9.
0.76±0.07±0.10	CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1.01±0.05	BARBERIS 98B		$450 \text{ pp} \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$

 $\Gamma_5/0.565\Gamma_4$  $\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi)$  $\Gamma_6/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.17±0.02±0.07</b>	CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.24±0.10	BAKER 24,25	SPEC 99	$1.94 \bar{p}p \rightarrow 4\pi^0$

 $\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$  $\frac{1}{2}\Gamma_5/\Gamma_8 = \frac{1}{2}\Gamma_5/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.29±0.04 OUR FIT</b>				
<b>0.29±0.05</b>	26 DAUM	81B SPEC		$63,94 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.3	BARTSCH 68	HBC +		$8 \pi^+ p \rightarrow 3\pi p$

 $0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$  $0.565\Gamma_4/\Gamma_8 = 0.565\Gamma_4/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ (With  $f_2(1270) \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.604±0.035 OUR FIT</b>				
<b>0.60 ±0.05 OUR AVERAGE</b>				Error includes scale factor of 1.3.
0.61 ± 0.04	26 DAUM	81B SPEC		$63,94 \pi^- p$
0.76 +0.24 -0.34	ARMENISE 69	DBC +		$5.1 \pi^+ d \rightarrow d 3\pi$
0.35 ± 0.20	BALTAY 68	HBC +		$7-8.5 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.59	BARTSCH 68	HBC +		$8 \pi^+ p \rightarrow 3\pi p$

 $0.624\Gamma(\pi(\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm\pi^+\pi^-)$  $0.624\Gamma_7/\Gamma_8 = 0.624\Gamma_7/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ (With  $(\pi\pi)_{S\text{-wave}} \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.10±0.04 OUR FIT</b>			
<b>0.10±0.05</b>	26 DAUM	81B SPEC	$63,94 \pi^- p$

$\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(f_2(1270)\pi)$   $\Gamma_9/\Gamma_4$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>0.075±0.025 OUR FIT</b>				
<b>0.075±0.025</b>	27 ARMSTRONG 82B OMEG	—	—	$16 \pi^- p \rightarrow K^+ K^- \pi^- p$

 $\Gamma(\omega\rho)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.027±0.004±0.010</b>	28 AMELIN 99 VES	37	$\pi^- A \xrightarrow{\omega\pi^- \pi^0} A^*$

 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$   $\Gamma_{13}/\Gamma_8 = \Gamma_{13}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.09	BALTAY 68 HBC	+	—	$7-8.5 \pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10	CRENNELL 70 HBC	—	—	$6 \pi^- p \rightarrow f_2 \pi^- N$
-------	-----------------	---	---	-------------------------------------

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$   $\Gamma_{14}/\Gamma_8 = \Gamma_{14}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.10	CRENNELL 70 HBC	—	—	$6 \pi^- p \rightarrow f_2 \pi^- N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	BALTAY 68 HBC	+	—	$7,8.5 \pi^+ p$
------	---------------	---	---	-----------------

 $\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0036	97.7	AMELIN 99 VES	37	$\pi^- A \xrightarrow{\omega\pi^- \pi^0} A^*$

 $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0019	97.7	AMELIN 99 VES	37	$\pi^- A \xrightarrow{\omega\pi^- \pi^0} A^*$

 $\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
possibly seen	69k	KUHN 04 B852	18	$\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	69k	KUHN 04 B852	18	$\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

**D-wave/S-wave RATIO FOR  $\pi_2(1670) \rightarrow f_2(1270)\pi$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.18±0.06	24 BAKER 99 SPEC	1.94	$\bar{p}p \rightarrow 4\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22±0.10	26 DAUM 81B SPEC	63,94	$\pi^- p$
-----------	------------------	-------	-----------

**F-wave/P-wave RATIO FOR  $\pi_2(1670) \rightarrow \rho\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.72±0.07±0.14</b>	CHUNG	02	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

24 Using preliminary CBAR data.

25 With the  $\sigma\pi$  in  $L=2$  and the  $f_2(1270)\pi$  in  $L=0$ .26 From a two-resonance fit to four  $2^-0^+$  waves.27 From a partial-wave analysis of  $K^+ K^- \pi^-$  system.28 Normalized to the  $B(\pi_2(1670) \rightarrow f_2\pi)$ . **$\pi_2(1670)$  REFERENCES**

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 62 487.		
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
		Translated from YAF 41 1223.		
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
Also		NP B186 594	C. Evangelista	
DAUM	80D	PL 89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TNTO, GENO, HAMB, MILA+) JP
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)
ARMENISE	69	LNC 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP