

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------------|------|---------------------------------------|
| 3686.10 ± 0.06 OUR FIT | | | | Error includes scale factor of 5.9. |
| 3686.097 ± 0.010 OUR AVERAGE | | | | |
| 3686.099 ± 0.004 ± 0.009 | | ¹ ANASHIN 15 | KEDR | $e^+e^- \rightarrow \text{hadrons}$ |
| 3686.12 ± 0.06 ± 0.10 | 4k | AAIJ 12H | LHCB | $pp \rightarrow J/\psi \pi^+ \pi^- X$ |
| 3685.95 ± 0.10 | 413 | ² ARTAMONOV 00 | OLYA | $e^+e^- \rightarrow \text{hadrons}$ |
| 3685.98 ± 0.09 ± 0.04 | | ³ ARMSTRONG 93B | E760 | $\bar{p}p \rightarrow e^+e^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 3686.114 ± 0.007 ^{+0.011} _{-0.016} | | ⁴ ANASHIN 12 | KEDR | $e^+e^- \rightarrow \text{hadrons}$ |
| 3686.111 ± 0.025 ± 0.009 | | AULCHENKO 03 | KEDR | $e^+e^- \rightarrow \text{hadrons}$ |
| 3686.00 ± 0.10 | 413 | ⁵ ZHOLENTZ 80 | OLYA | e^+e^- |

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁵ Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|--|
| 589.188 ± 0.028 OUR AVERAGE | | | |
| 589.194 ± 0.027 ± 0.011 | ¹ AULCHENKO 03 | KEDR | $e^+e^- \rightarrow \text{hadrons}$ |
| 589.7 ± 1.2 | LEMOIGNE 82 | GOLI | $185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$ |
| 589.07 ± 0.13 | ¹ ZHOLENTZ 80 | OLYA | e^+e^- |
| 588.7 ± 0.8 | LUTH 75 | MRK1 | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 588 ± 1 | ² BAI 98E | BES | e^+e^- |

¹ Redundant with data in mass above.

² Systematic errors not evaluated.

$\psi(2S)$ WIDTH

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|------------------------|----------|---|
| 294 ± 8 OUR FIT | | | | |
| 286 ± 16 OUR AVERAGE | | | | |
| 358 ± 88 ± 4 | | ABLIKIM | 08B BES2 | $e^+e^- \rightarrow \text{hadrons}$ |
| 290 ± 25 ± 4 | 2.7k | ANDREOTTI | 07 E835 | $\rho\bar{\rho} \rightarrow e^+e^-, J/\psi X$ |
| 331 ± 58 ± 2 | | ABLIKIM | 06L BES2 | $e^+e^- \rightarrow \text{hadrons}$ |
| 264 ± 27 | | ¹ BAI | 02B BES2 | e^+e^- |
| 287 ± 37 ± 16 | | ² ARMSTRONG | 93B E760 | $\bar{p}p \rightarrow e^+e^-$ |

¹From a simultaneous fit to the hadronic and $\mu^+\mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

²The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

 $\psi(2S)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|------------------------------------|-----------------------------------|
| Γ_1 hadrons | (97.85 ± 0.13) % | |
| Γ_2 virtual $\gamma \rightarrow$ hadrons | (1.73 ± 0.14) % | S=1.5 |
| Γ_3 ggg | (10.6 ± 1.6) % | |
| Γ_4 γgg | (1.03 ± 0.29) % | |
| Γ_5 light hadrons | (15.4 ± 1.5) % | |
| Γ_6 e^+e^- | (7.93 ± 0.17) × 10 ⁻³ | |
| Γ_7 $\mu^+\mu^-$ | (8.0 ± 0.6) × 10 ⁻³ | |
| Γ_8 $\tau^+\tau^-$ | (3.1 ± 0.4) × 10 ⁻³ | |

Decays into $J/\psi(1S)$ and anything

| | |
|--------------------------------------|--------------------------------------|
| Γ_9 $J/\psi(1S)$ anything | (61.4 ± 0.6) % |
| Γ_{10} $J/\psi(1S)$ neutrals | (25.38 ± 0.32) % |
| Γ_{11} $J/\psi(1S)\pi^+\pi^-$ | (34.68 ± 0.30) % |
| Γ_{12} $J/\psi(1S)\pi^0\pi^0$ | (18.24 ± 0.31) % |
| Γ_{13} $J/\psi(1S)\eta$ | (3.37 ± 0.05) % |
| Γ_{14} $J/\psi(1S)\pi^0$ | (1.268 ± 0.032) × 10 ⁻³ |

Hadronic decays

| | | |
|--|------------------------------------|--------|
| Γ_{15} $\pi^0 h_c(1P)$ | (8.6 ± 1.3) × 10 ⁻⁴ | |
| Γ_{16} $3(\pi^+\pi^-\pi^0)$ | (3.5 ± 1.6) × 10 ⁻³ | |
| Γ_{17} $2(\pi^+\pi^-\pi^0)$ | (2.9 ± 1.0) × 10 ⁻³ | S=4.7 |
| Γ_{18} $\rho a_2(1320)$ | (2.6 ± 0.9) × 10 ⁻⁴ | |
| Γ_{19} $\pi^+\pi^-\pi^0\pi^0\pi^0$ | (5.3 ± 0.9) × 10 ⁻³ | |
| Γ_{20} $\rho^\pm\pi^\mp\pi^0\pi^0$ | < 2.7 × 10 ⁻³ | CL=90% |
| Γ_{21} $\rho\bar{\rho}$ | (2.94 ± 0.08) × 10 ⁻⁴ | |
| Γ_{22} $n\bar{n}$ | (3.06 ± 0.15) × 10 ⁻⁴ | |
| Γ_{23} $\Delta^{++}\bar{\Delta}^{--}$ | (1.28 ± 0.35) × 10 ⁻⁴ | |
| Γ_{24} $\Lambda\bar{\Lambda}\pi^0$ | < 2.9 × 10 ⁻⁶ | CL=90% |
| Γ_{25} $\Lambda\bar{\Lambda}\eta$ | (2.5 ± 0.4) × 10 ⁻⁵ | |

| | | | |
|---------------|---|--|-------|
| Γ_{26} | $\Lambda \bar{p} K^+$ | $(1.00 \pm 0.14) \times 10^{-4}$ | |
| Γ_{27} | $K^*(892)^+ \bar{p} \Lambda + \text{c.c.}$ | $(6.3 \pm 0.7) \times 10^{-5}$ | |
| Γ_{28} | $\Lambda \bar{p} K^+ \pi^+ \pi^-$ | $(1.8 \pm 0.4) \times 10^{-4}$ | |
| Γ_{29} | $\Lambda \bar{\Lambda} \pi^+ \pi^-$ | $(2.8 \pm 0.6) \times 10^{-4}$ | |
| Γ_{30} | $\Lambda \bar{\Lambda}$ | $(3.81 \pm 0.13) \times 10^{-4}$ | S=1.4 |
| Γ_{31} | $\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.}$ | $(1.40 \pm 0.13) \times 10^{-4}$ | |
| Γ_{32} | $\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.}$ | $(1.54 \pm 0.14) \times 10^{-4}$ | |
| Γ_{33} | $\Lambda \bar{\Sigma}^0$ | $(1.23 \pm 0.24) \times 10^{-5}$ | |
| Γ_{34} | $\Sigma^0 \bar{p} K^+ + \text{c.c.}$ | $(1.67 \pm 0.18) \times 10^{-5}$ | |
| Γ_{35} | $\Sigma^+ \bar{\Sigma}^-$ | $(2.32 \pm 0.12) \times 10^{-4}$ | |
| Γ_{36} | $\Sigma^0 \bar{\Sigma}^0$ | $(2.35 \pm 0.09) \times 10^{-4}$ | S=1.1 |
| Γ_{37} | $\Sigma(1385)^+ \bar{\Sigma}(1385)^-$ | $(8.5 \pm 0.7) \times 10^{-5}$ | |
| Γ_{38} | $\Sigma(1385)^- \bar{\Sigma}(1385)^+$ | $(8.5 \pm 0.8) \times 10^{-5}$ | |
| Γ_{39} | $\Sigma(1385)^0 \bar{\Sigma}(1385)^0$ | $(6.9 \pm 0.7) \times 10^{-5}$ | |
| Γ_{40} | $\Xi^- \bar{\Xi}^+$ | $(2.87 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{41} | $\Xi^0 \bar{\Xi}^0$ | $(2.3 \pm 0.4) \times 10^{-4}$ | S=4.2 |
| Γ_{42} | $\Xi(1530)^0 \bar{\Xi}(1530)^0$ | $(5.2 \begin{smallmatrix} +3.2 \\ -1.2 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{43} | $K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$ | $(3.9 \pm 0.4) \times 10^{-5}$ | |
| Γ_{44} | $\Xi(1530)^- \bar{\Xi}(1530)^+$ | $(1.15 \pm 0.07) \times 10^{-4}$ | |
| Γ_{45} | $\Xi(1530)^- \bar{\Xi}^+$ | $(7.0 \pm 1.2) \times 10^{-6}$ | |
| Γ_{46} | $\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ | $(5.2 \pm 1.6) \times 10^{-6}$ | |
| Γ_{47} | $\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ | $(1.20 \pm 0.32) \times 10^{-5}$ | |
| Γ_{48} | $K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.}$ | $(3.7 \pm 0.4) \times 10^{-5}$ | |
| Γ_{49} | $\Omega^- \bar{\Omega}^+$ | $(5.2 \pm 0.4) \times 10^{-5}$ | |
| Γ_{50} | $\pi^0 \rho \bar{p}$ | $(1.53 \pm 0.07) \times 10^{-4}$ | |
| Γ_{51} | $N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{52} | $N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}) \times 10^{-5}$ | S=2.5 |
| Γ_{53} | $N(1520) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$ | |
| Γ_{54} | $N(1535) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.5 \pm 1.0) \times 10^{-5}$ | |
| Γ_{55} | $N(1650) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{56} | $N(1720) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{57} | $N(2300) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{58} | $N(2570) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{p}$ | $(2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix}) \times 10^{-5}$ | |
| Γ_{59} | $\pi^0 f_0(2100) \rightarrow \pi^0 \rho \bar{p}$ | $(1.1 \pm 0.4) \times 10^{-5}$ | |
| Γ_{60} | $\eta \rho \bar{p}$ | $(6.0 \pm 0.4) \times 10^{-5}$ | |
| Γ_{61} | $\eta f_0(2100) \rightarrow \eta \rho \bar{p}$ | $(1.2 \pm 0.4) \times 10^{-5}$ | |
| Γ_{62} | $N(1535) \bar{p} \rightarrow \eta \rho \bar{p}$ | $(4.4 \pm 0.7) \times 10^{-5}$ | |
| Γ_{63} | $\omega \rho \bar{p}$ | $(6.9 \pm 2.1) \times 10^{-5}$ | |

| | | | |
|----------------|---|------------------------------------|--------|
| Γ_{64} | $\eta' p\bar{p}$ | $(1.10 \pm 0.13) \times 10^{-5}$ | |
| Γ_{65} | $\phi p\bar{p}$ | $(6.1 \pm 0.6) \times 10^{-6}$ | |
| Γ_{66} | $\phi X(1835) \rightarrow \phi p\bar{p}$ | $< 1.82 \times 10^{-7}$ | CL=90% |
| Γ_{67} | $\pi^+ \pi^- p\bar{p}$ | $(6.0 \pm 0.4) \times 10^{-4}$ | |
| Γ_{68} | $\rho \bar{n} \pi^-$ or c.c. | $(2.48 \pm 0.17) \times 10^{-4}$ | |
| Γ_{69} | $\rho \bar{n} \pi^- \pi^0$ | $(3.2 \pm 0.7) \times 10^{-4}$ | |
| Γ_{70} | $2(\pi^+ \pi^- \pi^0)$ | $(4.8 \pm 1.5) \times 10^{-3}$ | |
| Γ_{71} | $\eta \pi^+ \pi^-$ | $< 1.6 \times 10^{-4}$ | CL=90% |
| Γ_{72} | $\eta \pi^+ \pi^- \pi^0$ | $(9.5 \pm 1.7) \times 10^{-4}$ | |
| Γ_{73} | $2(\pi^+ \pi^-) \eta$ | $(1.2 \pm 0.6) \times 10^{-3}$ | |
| Γ_{74} | $\pi^+ \pi^- \pi^0 \pi^0 \eta$ | $< 4 \times 10^{-4}$ | CL=90% |
| Γ_{75} | $\eta' \pi^+ \pi^- \pi^0$ | $(4.5 \pm 2.1) \times 10^{-4}$ | |
| Γ_{76} | $\omega \pi^+ \pi^-$ | $(7.3 \pm 1.2) \times 10^{-4}$ | S=2.1 |
| Γ_{77} | $b_1^\pm \pi^\mp$ | $(4.0 \pm 0.6) \times 10^{-4}$ | S=1.1 |
| Γ_{78} | $b_1^0 \pi^0$ | $(2.4 \pm 0.6) \times 10^{-4}$ | |
| Γ_{79} | $\omega f_2(1270)$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{80} | $\omega \pi^0 \pi^0$ | $(1.11 \pm 0.35) \times 10^{-3}$ | |
| Γ_{81} | $\pi^0 \pi^0 K^+ K^-$ | $(2.6 \pm 1.3) \times 10^{-4}$ | |
| Γ_{82} | $\pi^+ \pi^- K^+ K^-$ | $(7.3 \pm 0.5) \times 10^{-4}$ | |
| Γ_{83} | $\pi^0 \pi^0 K_S^0 K_L^0$ | $(1.3 \pm 0.6) \times 10^{-3}$ | |
| Γ_{84} | $\rho^0 K^+ K^-$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{85} | $K^*(892)^0 \bar{K}_2^*(1430)^0$ | $(1.9 \pm 0.5) \times 10^{-4}$ | |
| Γ_{86} | $K^+ K^- \pi^+ \pi^- \eta$ | $(1.3 \pm 0.7) \times 10^{-3}$ | |
| Γ_{87} | $K^+ K^- 2(\pi^+ \pi^-) \pi^0$ | $(1.00 \pm 0.31) \times 10^{-3}$ | |
| Γ_{88} | $K^+ K^- 2(\pi^+ \pi^-)$ | $(1.9 \pm 0.9) \times 10^{-3}$ | |
| Γ_{89} | $K_1(1270)^\pm K^\mp$ | $(1.00 \pm 0.28) \times 10^{-3}$ | |
| Γ_{90} | $K_S^0 K_S^0 \pi^+ \pi^-$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{91} | $\rho^0 p\bar{p}$ | $(5.0 \pm 2.2) \times 10^{-5}$ | |
| Γ_{92} | $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$ | $(6.7 \pm 2.5) \times 10^{-4}$ | |
| Γ_{93} | $2(\pi^+ \pi^-)$ | $(2.4 \pm 0.6) \times 10^{-4}$ | S=2.2 |
| Γ_{94} | $\rho^0 \pi^+ \pi^-$ | $(2.2 \pm 0.6) \times 10^{-4}$ | S=1.4 |
| Γ_{95} | $K^+ K^- \pi^+ \pi^- \pi^0$ | $(1.26 \pm 0.09) \times 10^{-3}$ | |
| Γ_{96} | $\omega f_0(1710) \rightarrow \omega K^+ K^-$ | $(5.9 \pm 2.2) \times 10^{-5}$ | |
| Γ_{97} | $K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$ | $(8.6 \pm 2.2) \times 10^{-4}$ | |
| Γ_{98} | $K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$ | $(9.6 \pm 2.8) \times 10^{-4}$ | |
| Γ_{99} | $K^*(892)^+ K^- \rho^0 + \text{c.c.}$ | $(7.3 \pm 2.6) \times 10^{-4}$ | |
| Γ_{100} | $K^*(892)^0 K^- \rho^+ + \text{c.c.}$ | $(6.1 \pm 1.8) \times 10^{-4}$ | |
| Γ_{101} | $\eta K^+ K^-$, no $\eta \phi$ | $(3.1 \pm 0.4) \times 10^{-5}$ | |
| Γ_{102} | $\omega K^+ K^-$ | $(1.62 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{103} | $\omega K^*(892)^+ K^- + \text{c.c.}$ | $(2.07 \pm 0.26) \times 10^{-4}$ | |
| Γ_{104} | $\omega K_2^*(1430)^+ K^- + \text{c.c.}$ | $(6.1 \pm 1.2) \times 10^{-5}$ | |
| Γ_{105} | $\omega \bar{K}^*(892)^0 K^0$ | $(1.68 \pm 0.30) \times 10^{-4}$ | |
| Γ_{106} | $\omega \bar{K}_2^*(1430)^0 K^0$ | $(5.8 \pm 2.2) \times 10^{-5}$ | |

| | | | |
|----------------|---|------------------------------------|--------|
| Γ_{107} | $\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c. | $(1.6 \pm 0.4) \times 10^{-5}$ | |
| Γ_{108} | $\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$ | $(1.09 \pm 0.26) \times 10^{-5}$ | |
| Γ_{109} | $\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c. | $(3.0 \pm 1.0) \times 10^{-6}$ | |
| Γ_{110} | $\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$ | $(1.2 \pm 0.7) \times 10^{-6}$ | |
| Γ_{111} | $3(\pi^+ \pi^-)$ | $(3.5 \pm 2.0) \times 10^{-4}$ | S=2.8 |
| Γ_{112} | $p\bar{p}\pi^+\pi^-\pi^0$ | $(7.3 \pm 0.7) \times 10^{-4}$ | |
| Γ_{113} | $K^+ K^-$ | $(7.5 \pm 0.5) \times 10^{-5}$ | |
| Γ_{114} | $K_S^0 K_L^0$ | $(5.34 \pm 0.33) \times 10^{-5}$ | |
| Γ_{115} | $\pi^+ \pi^- \pi^0$ | $(2.01 \pm 0.17) \times 10^{-4}$ | S=1.7 |
| Γ_{116} | $\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$ | $(1.9 \pm 1.2) \times 10^{-4}$ | |
| Γ_{117} | $\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$ | $(3.2 \pm 1.2) \times 10^{-5}$ | S=1.8 |
| Γ_{118} | $\pi^+ \pi^-$ | $(7.8 \pm 2.6) \times 10^{-6}$ | |
| Γ_{119} | $K_1(1400)^\pm K^\mp$ | $< 3.1 \times 10^{-4}$ | CL=90% |
| Γ_{120} | $K_2^*(1430)^\pm K^\mp$ | $(7.1 \pm 1.3) \times 10^{-5}$ | |
| Γ_{121} | $K^+ K^- \pi^0$ | $(4.07 \pm 0.31) \times 10^{-5}$ | |
| Γ_{122} | $K_S^0 K_L^0 \pi^0$ | $< 3.0 \times 10^{-4}$ | CL=90% |
| Γ_{123} | $K_S^0 K_L^0 \eta$ | $(1.3 \pm 0.5) \times 10^{-3}$ | |
| Γ_{124} | $K^+ K^*(892)^- + \text{c.c.}$ | $(2.9 \pm 0.4) \times 10^{-5}$ | S=1.2 |
| Γ_{125} | $K^*(892)^0 \bar{K}^0 + \text{c.c.}$ | $(1.09 \pm 0.20) \times 10^{-4}$ | |
| Γ_{126} | $\phi \pi^+ \pi^-$ | $(1.18 \pm 0.26) \times 10^{-4}$ | S=1.5 |
| Γ_{127} | $\phi f_0(980) \rightarrow \pi^+ \pi^-$ | $(7.5 \pm 3.3) \times 10^{-5}$ | S=1.6 |
| Γ_{128} | $2(K^+ K^-)$ | $(6.3 \pm 1.3) \times 10^{-5}$ | |
| Γ_{129} | $\phi K^+ K^-$ | $(7.0 \pm 1.6) \times 10^{-5}$ | |
| Γ_{130} | $2(K^+ K^-) \pi^0$ | $(1.10 \pm 0.28) \times 10^{-4}$ | |
| Γ_{131} | $\phi \eta$ | $(3.10 \pm 0.31) \times 10^{-5}$ | |
| Γ_{132} | $\eta \phi(2170), \phi(2170) \rightarrow$ $\phi f_0(980), f_0 \rightarrow \pi^+ \pi^-$ | $< 2.2 \times 10^{-6}$ | CL=90% |
| Γ_{133} | $\phi \eta'$ | $(1.54 \pm 0.20) \times 10^{-5}$ | |
| Γ_{134} | $\phi f_1(1285)$ | $(3.0 \pm 1.3) \times 10^{-5}$ | |
| Γ_{135} | $\phi \eta(1405) \rightarrow \phi \pi^+ \pi^- \eta$ | $(8.5 \pm 1.7) \times 10^{-6}$ | |
| Γ_{136} | $\omega \eta'$ | $(3.2 \pm 2.5) \times 10^{-5}$ | |
| Γ_{137} | $\omega \pi^0$ | $(2.1 \pm 0.6) \times 10^{-5}$ | |
| Γ_{138} | $\rho \eta'$ | $(1.9 \pm 1.7) \times 10^{-5}$ | |
| Γ_{139} | $\rho \eta$ | $(2.2 \pm 0.6) \times 10^{-5}$ | S=1.1 |
| Γ_{140} | $\omega \eta$ | $< 1.1 \times 10^{-5}$ | CL=90% |
| Γ_{141} | $\phi \pi^0$ | $< 4 \times 10^{-7}$ | CL=90% |
| Γ_{142} | $\eta_c \pi^+ \pi^- \pi^0$ | $< 1.0 \times 10^{-3}$ | CL=90% |
| Γ_{143} | $p\bar{p} K^+ K^-$ | $(2.7 \pm 0.7) \times 10^{-5}$ | |
| Γ_{144} | $\bar{\Lambda} n K_S^0 + \text{c.c.}$ | $(8.1 \pm 1.8) \times 10^{-5}$ | |
| Γ_{145} | $\phi f_2'(1525)$ | $(4.4 \pm 1.6) \times 10^{-5}$ | |

| | | | | |
|----------------|--|---------|------------------|--------|
| Γ_{146} | $\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$ | < 8.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{147} | $\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$ | < 1.0 | $\times 10^{-5}$ | CL=90% |
| Γ_{148} | $\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$ | < 7.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{149} | $\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n$ | < 2.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{150} | $\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$ | < 6.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{151} | $K_S^0 K_S^0$ | < 4.6 | $\times 10^{-6}$ | |
| Γ_{152} | $\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$ | < 1.7 | $\times 10^{-6}$ | CL=90% |

Radiative decays

| | | | | |
|----------------|--|---|------------------|--------|
| Γ_{153} | $\gamma \chi_{c0}(1P)$ | $(9.79 \pm 0.20) \%$ | | |
| Γ_{154} | $\gamma \chi_{c1}(1P)$ | $(9.75 \pm 0.24) \%$ | | |
| Γ_{155} | $\gamma \chi_{c2}(1P)$ | $(9.52 \pm 0.20) \%$ | | |
| Γ_{156} | $\gamma \eta_c(1S)$ | $(3.4 \pm 0.5) \times 10^{-3}$ | | S=1.3 |
| Γ_{157} | $\gamma \eta_c(2S)$ | $(7 \pm 5) \times 10^{-4}$ | | |
| Γ_{158} | $\gamma \pi^0$ | $(1.04 \pm 0.22) \times 10^{-6}$ | | S=1.4 |
| Γ_{159} | $\gamma \eta'(958)$ | $(1.24 \pm 0.04) \times 10^{-4}$ | | |
| Γ_{160} | $\gamma f_2(1270)$ | $(2.73^{+0.29}_{-0.25}) \times 10^{-4}$ | | S=1.8 |
| Γ_{161} | $\gamma f_0(1370) \rightarrow \gamma K \bar{K}$ | $(3.1 \pm 1.7) \times 10^{-5}$ | | |
| Γ_{162} | $\gamma f_0(1500)$ | $(9.3 \pm 1.9) \times 10^{-5}$ | | |
| Γ_{163} | $\gamma f_2'(1525)$ | $(3.3 \pm 0.8) \times 10^{-5}$ | | |
| Γ_{164} | $\gamma f_0(1710)$ | | | |
| Γ_{165} | $\gamma f_0(1710) \rightarrow \gamma \pi \pi$ | $(3.5 \pm 0.6) \times 10^{-5}$ | | |
| Γ_{166} | $\gamma f_0(1710) \rightarrow \gamma K \bar{K}$ | $(6.6 \pm 0.7) \times 10^{-5}$ | | |
| Γ_{167} | $\gamma f_0(2100) \rightarrow \gamma \pi \pi$ | $(4.8 \pm 1.0) \times 10^{-6}$ | | |
| Γ_{168} | $\gamma f_0(2200) \rightarrow \gamma K \bar{K}$ | $(3.2 \pm 1.0) \times 10^{-6}$ | | |
| Γ_{169} | $\gamma f_J(2220) \rightarrow \gamma \pi \pi$ | < 5.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{170} | $\gamma f_J(2220) \rightarrow \gamma K \bar{K}$ | < 9.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{171} | $\gamma \gamma$ | < 1.5 | $\times 10^{-4}$ | CL=90% |
| Γ_{172} | $\gamma \eta$ | $(9.2 \pm 1.8) \times 10^{-7}$ | | |
| Γ_{173} | $\gamma \eta \pi^+ \pi^-$ | $(8.7 \pm 2.1) \times 10^{-4}$ | | |
| Γ_{174} | $\gamma \eta(1405)$ | | | |
| Γ_{175} | $\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$ | < 9 | $\times 10^{-5}$ | CL=90% |
| Γ_{176} | $\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-$ | $(3.6 \pm 2.5) \times 10^{-5}$ | | |
| Γ_{177} | $\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^+ \pi^- \pi^0$ | < 5.0 | $\times 10^{-7}$ | CL=90% |
| Γ_{178} | $\gamma \eta(1475)$ | | | |
| Γ_{179} | $\gamma \eta(1475) \rightarrow K \bar{K} \pi$ | < 1.4 | $\times 10^{-4}$ | CL=90% |
| Γ_{180} | $\gamma \eta(1475) \rightarrow \eta \pi^+ \pi^-$ | < 8.8 | $\times 10^{-5}$ | CL=90% |
| Γ_{181} | $\gamma 2(\pi^+ \pi^-)$ | $(4.0 \pm 0.6) \times 10^{-4}$ | | |
| Γ_{182} | $\gamma K^{*0} K^+ \pi^- + \text{c.c.}$ | $(3.7 \pm 0.9) \times 10^{-4}$ | | |
| Γ_{183} | $\gamma K^{*0} \bar{K}^{*0}$ | $(2.4 \pm 0.7) \times 10^{-4}$ | | |
| Γ_{184} | $\gamma K_S^0 K^+ \pi^- + \text{c.c.}$ | $(2.6 \pm 0.5) \times 10^{-4}$ | | |

| | | | |
|----------------|---|---|-------------------------|
| Γ_{185} | $\gamma K^+ K^- \pi^+ \pi^-$ | $(1.9 \pm 0.5) \times 10^{-4}$ | |
| Γ_{186} | $\gamma p \bar{p}$ | $(3.9 \pm 0.5) \times 10^{-5}$ | S=2.0 |
| Γ_{187} | $\gamma f_2(1950) \rightarrow \gamma p \bar{p}$ | $(1.20 \pm 0.22) \times 10^{-5}$ | |
| Γ_{188} | $\gamma f_2(2150) \rightarrow \gamma p \bar{p}$ | $(7.2 \pm 1.8) \times 10^{-6}$ | |
| Γ_{189} | $\gamma X(1835) \rightarrow \gamma p \bar{p}$ | $(4.6 \begin{smallmatrix} +1.8 \\ -4.0 \end{smallmatrix}) \times 10^{-6}$ | |
| Γ_{190} | $\gamma X \rightarrow \gamma p \bar{p}$ | [a] < 2 | $\times 10^{-6}$ CL=90% |
| Γ_{191} | $\gamma \pi^+ \pi^- p \bar{p}$ | $(2.8 \pm 1.4) \times 10^{-5}$ | |
| Γ_{192} | $\gamma 2(\pi^+ \pi^-) K^+ K^-$ | < 2.2 | $\times 10^{-4}$ CL=90% |
| Γ_{193} | $\gamma 3(\pi^+ \pi^-)$ | < 1.7 | $\times 10^{-4}$ CL=90% |
| Γ_{194} | $\gamma K^+ K^- K^+ K^-$ | < 4 | $\times 10^{-5}$ CL=90% |
| Γ_{195} | $\gamma \gamma J/\psi$ | $(3.1 \begin{smallmatrix} +1.0 \\ -1.2 \end{smallmatrix}) \times 10^{-4}$ | |
| Γ_{196} | $e^+ e^- \eta'$ | $(1.90 \pm 0.26) \times 10^{-6}$ | |
| Γ_{197} | $e^+ e^- \chi_{c0}(1P)$ | $(1.06 \pm 0.24) \times 10^{-3}$ | |
| Γ_{198} | $e^+ e^- \chi_{c1}(1P)$ | $(8.5 \pm 0.6) \times 10^{-4}$ | |
| Γ_{199} | $e^+ e^- \chi_{c2}(1P)$ | $(7.0 \pm 0.8) \times 10^{-4}$ | |

Weak decays

| | | | |
|----------------|-----------------------------|-------|-------------------------|
| Γ_{200} | $D^0 e^+ e^- + \text{c.c.}$ | < 1.4 | $\times 10^{-7}$ CL=90% |
|----------------|-----------------------------|-------|-------------------------|

Other decays

| | | | |
|----------------|-----------|-------|----------|
| Γ_{201} | invisible | < 1.6 | % CL=90% |
|----------------|-----------|-------|----------|

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 378.1$ for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

| | | | | | | | | | | |
|-----------|-------|-------|-------|----------|----------|----------|----------|-----------|-----------|-----------|
| x_7 | 3 | | | | | | | | | |
| x_8 | 1 | 0 | | | | | | | | |
| x_{11} | 29 | 11 | 2 | | | | | | | |
| x_{12} | 28 | 6 | 1 | 48 | | | | | | |
| x_{13} | 13 | 4 | 1 | 36 | 15 | | | | | |
| x_{21} | 0 | 0 | 0 | 4 | 3 | 2 | | | | |
| x_{153} | 1 | 0 | 0 | 2 | 1 | 1 | 0 | | | |
| x_{154} | 1 | 0 | 0 | 2 | 1 | 1 | 0 | 0 | | |
| x_{155} | 1 | 0 | 0 | 3 | 1 | 1 | 0 | 0 | 0 | |
| Γ | -81 | -4 | -1 | -38 | -34 | -16 | -7 | -1 | -1 | -1 |
| | x_6 | x_7 | x_8 | x_{11} | x_{12} | x_{13} | x_{21} | x_{153} | x_{154} | x_{155} |

$\psi(2S)$ PARTIAL WIDTHS $\Gamma(\text{hadrons})$ Γ_1

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|--------------|-------------|------|---------------|
| 258 ± 26 | BAI | 02B | BES2 e^+e^- |
| 224 ± 56 | LUTH | 75 | MRK1 e^+e^- |

 $\Gamma(e^+e^-)$ Γ_6

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---|
| 2.33 ± 0.04 OUR FIT | | | |
| 2.29 ± 0.06 OUR AVERAGE | | | |
| $2.23 \pm 0.10 \pm 0.02$ | ¹ ABLIKIM | 15V | BES3 4.0–4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ |
| $2.338 \pm 0.037 \pm 0.096$ | ABLIKIM | 08B | BES2 $e^+e^- \rightarrow \text{hadrons}$ |
| $2.330 \pm 0.036 \pm 0.110$ | ABLIKIM | 06L | BES2 $e^+e^- \rightarrow \text{hadrons}$ |
| 2.44 ± 0.21 | ² BAI | 02B | BES2 e^+e^- |
| 2.14 ± 0.21 | ALEXANDER | 89 | RVUE See Υ mini-review |
| $2.279 \pm 0.015 \pm 0.042$ | ³ ANASHIN | 18 | KEDR e^+e^- |
| $2.282 \pm 0.015 \pm 0.042$ | ⁴ ANASHIN | 18 | KEDR e^+e^- |
| 2.0 ± 0.3 | BRANDELIK | 79C | DASP e^+e^- |
| 2.1 ± 0.3 | ⁵ LUTH | 75 | MRK1 e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$.

³ Combining $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18 with $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12 and assuming lepton universality.

⁴ From the sum of $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12, $\Gamma_{e^+e^-} \cdot B(e^+e^-)$ and $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18, and $\Gamma_{e^+e^-} \cdot B(\tau^+\tau^-)$ from ANASHIN 07.

⁵ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

 $\Gamma(\gamma\gamma)$ Γ_{171}

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------|-----|-------------|------|---------------|
| <43 | 90 | BRANDELIK | 79C | DASP e^+e^- |

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_6/\Gamma$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|--|
| 2.233±0.015±0.042 | ¹ ANASHIN | 12 | KEDR $e^+e^- \rightarrow \text{hadrons}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 2.2 ± 0.4 | ABRAMS | 75 | MRK1 e^+e^- |

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

 $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_6\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|---------------|
| 21.2±0.7±1.2 | ¹ ANASHIN | 18 | KEDR e^+e^- |

¹ From the average of nine scans of the $\psi(2S)$.

 $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|--|
| 19.3±0.3±0.5 | ¹ ANASHIN | 18 | KEDR $\psi(2S) \rightarrow \mu^+\mu^-$ |

¹ From the average of nine scans of the $\psi(2S)$.

 $\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 9.0±2.6 | 79 | ¹ ANASHIN | 07 | KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$ |

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

 $\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_6/\Gamma$


| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------|-------------------|------|---|
| 0.808±0.013 OUR FIT | | | | |
| 0.837±0.025 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.837±0.028±0.005 | | ¹ LEES | 12E | BABR 10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$ |
| 0.852±0.010±0.026 | 19.5k | ADAM | 06 | CLEO 3.773 $e^+e^- \rightarrow \gamma\psi(2S)$ |
| 0.68 ± 0.09 | | ² BAI | 98E | BES e^+e^- |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|-----|---------------------|------|---|
| 0.88 ± 0.08 ± 0.03 | 256 | ³ AUBERT | 07AU | BABR 10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$ |
| 0.755±0.048±0.004 | 544 | ⁴ AUBERT | 05D | BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$ |

¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ²The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ³AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.



| $\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | $\Gamma_{12}\Gamma_6/\Gamma$ |
|---|---|
| VALUE (keV) | EVTS DOCUMENT ID TECN COMMENT |
| 0.425±0.009 OUR FIT | |
| 0.411±0.008±0.018 | 3.6k ADAM 06 CLEO 3.773 $e^+e^- \rightarrow \gamma\psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| 0.48 ±0.09 ±0.02 | 142 ¹ LEES 18E BABR 10.6 $e^+e^- \rightarrow J/\psi\pi^0\pi^0\gamma$ |

¹LEES 18E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0101 \pm 0.0015 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|---------------------|-----------|---|
| 78.6 ± 1.6 OUR FIT | | | | |
| 87 ± 9 OUR AVERAGE | | | | |
| 83 ± 25 ± 5 | 14 | ¹ AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → J/ψ π ⁺ π ⁻ π ⁰ γ |
| 88 ± 6 ± 7 | 291 ± 24 | ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γ ψ(2S) |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi \eta) \cdot B(J/\psi \rightarrow \mu^+ \mu^-) \cdot B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV. | | | | |

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_6/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------|-----|------|-------------|---------|---|
| <8 | 90 | <37 | ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γ ψ(2S) |

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{21}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|----------|---|
| 0.686 ± 0.019 OUR FIT | | | | |
| 0.63 ± 0.05 OUR AVERAGE | | | | Error includes scale factor of 1.2. |
| 0.67 ± 0.12 ± 0.02 | 43 | ¹ LEES | 130 BABR | e ⁺ e ⁻ → p \bar{p} γ |
| 0.74 ± 0.07 ± 0.04 | 142 | ² LEES | 13Y BABR | e ⁺ e ⁻ → p \bar{p} γ |
| 0.579 ± 0.038 ± 0.036 | 2.7k | ANDREOTTI | 07 E835 | p \bar{p} → e ⁺ e ⁻ , J/ψ X |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.70 ± 0.17 ± 0.03 | 22 | ³ AUBERT | 06B BABR | e ⁺ e ⁻ → p \bar{p} γ |
| ¹ ISR photon reconstructed in the detector | | | | |
| ² ISR photon undetected | | | | |
| ³ Superseded by LEES 130 | | | | |

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{30}\Gamma_6/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|-----------|--|
| 1.5 ± 0.4 ± 0.1 | AUBERT | 07BD BABR | 10.6 e ⁺ e ⁻ → Λ $\bar{\Lambda}$ γ |

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{70}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|-------------|----------|---|
| 11.2 ± 3.3 ± 1.3 | 43 | AUBERT | 06D BABR | 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰) γ |

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{81}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|--|
| 0.60 ± 0.31 ± 0.03 | 17 | LEES | 12F BABR | 10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ |

$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{88}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-------------|----------|--|
| 4.4 ± 2.1 ± 0.3 | 26 | AUBERT | 06D BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻) γ |

$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{82}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|-----------------------|-----|------|----------|--|
| 1.92±0.30±0.06 | 133 | LEES | 12F BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |
|-----------------------|-----|------|----------|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------------|----|---------------------|-----------|--|
| 2.56±0.42±0.16 | 85 | ¹ AUBERT | 07AK BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |
|----------------|----|---------------------|-----------|--|

¹Superseded by LEES 12F.

$\Gamma(\pi^0\pi^0K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{83}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|-----------------------|----|------|----------|--|
| 2.92±1.27±0.15 | 14 | LEES | 17A BABR | e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ |
|-----------------------|----|------|----------|--|

$\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{122}\Gamma_6/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|-----|------|-------------|------|---------|
|------------|-----|------|-------------|------|---------|

| | | | | | |
|----------------|----|---|------|----------|--|
| <0.7 | 90 | 8 | LEES | 17A BABR | e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ |
|----------------|----|---|------|----------|--|

$\Gamma(K_S^0K_L^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{123}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|-----------------------|----|------|----------|--|
| 3.14±1.08±0.16 | 16 | LEES | 17A BABR | e ⁺ e ⁻ → K _S ⁰ K _L ⁰ ηγ |
|-----------------------|----|------|----------|--|

$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{127}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|--------------------------|----|-------------------|----------|--|
| 0.345±0.128±0.004 | 12 | ¹ LEES | 12F BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |
|--------------------------|----|-------------------|----------|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|-------|---------------------|-----------|--|
| 0.345±0.168±0.004 | 6 ± 3 | ² AUBERT | 07AK BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ |
|-------------------|-------|---------------------|-----------|--|

¹LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 0.17 ± 0.06 ± 0.02 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 0.17 ± 0.08 ± 0.02 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{128}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|-----------------------|----|------|----------|--|
| 0.22±0.10±0.02 | 13 | LEES | 12F BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ K ⁺ K ⁻ γ |
|-----------------------|----|------|----------|--|

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{126}\Gamma_6/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

| | | | | |
|-----------------------|----|-------------------|----------|--|
| 0.55±0.19±0.01 | 19 | ¹ LEES | 12F BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ |
|-----------------------|----|-------------------|----------|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------------|----|-------------------------|----------|--|
| 0.57±0.23±0.01 | 10 | ² AUBERT, BE | 06D BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ |
|----------------|----|-------------------------|----------|--|

¹LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 0.27 ± 0.09 ± 0.02 eV which we divide by our best value

$B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{17} \Gamma_6 / \Gamma$ |
|---|------|-------------|-----------|--|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 29.7 ± 2.2 ± 1.8 | 410 | AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$ | |

| $\Gamma(\pi^+ \pi^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{19} \Gamma_6 / \Gamma$ |
|--|------|-------------|----------|--|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 12.4 ± 1.8 ± 1.2 | 177 | LEES | 18E BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$ | |

| $\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{20} \Gamma_6 / \Gamma$ |
|---|-----|-------------|----------|--|---------------------------------|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <6.2 | 90 | LEES | 18E BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$ | |

| $\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{76} \Gamma_6 / \Gamma$ |
|---|------|---------------------|-----------|--|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 3.01 ± 0.84 ± 0.02 | 37 | ¹ AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$ | |

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 2.69 \pm 0.73 \pm 0.16$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{80} \Gamma_6 / \Gamma$ |
|---|------|-------------------|----------|--|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 2.58 ± 0.82 ± 0.02 | 33 | ¹ LEES | 18E BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$ | |

¹LEES 18E reports $[\Gamma(\psi(2S) \rightarrow \omega \pi^0 \pi^0) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 2.3 \pm 0.7 \pm 0.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(2(\pi^+ \pi^-) \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{73} \Gamma_6 / \Gamma$ |
|--|------|---------------------|-----------|---|---------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 2.87 ± 1.41 ± 0.01 | 16 | ¹ AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$ | |

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\pi^+ \pi^- \pi^0 \pi^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ | | | | | $\Gamma_{74} \Gamma_6 / \Gamma$ |
|---|-----|-------------|----------|--|---------------------------------|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.85 | 90 | LEES | 18E BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$ | |

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{95} \Gamma_6 / \Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-------------|-----------|---|
| 4.4 ± 1.3 ± 0.3 | 32 | AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ |

 $\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{86} \Gamma_6 / \Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|---------------------|-----------|---|
| 3.04 ± 1.79 ± 0.02 | 7 | ¹ AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ |

¹AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B(η → 2γ)] = 1.2 ± 0.7 ± 0.1 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{113} \Gamma_6 / \Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |

| | | | | |
|-----------------------|----|-------------------|----------|---|
| 0.147 ± 0.035 ± 0.005 | 66 | ¹ LEES | 15J BABR | e ⁺ e ⁻ → K ⁺ K ⁻ γ |
| 0.197 ± 0.035 ± 0.005 | 66 | ² LEES | 15J BABR | e ⁺ e ⁻ → K ⁺ K ⁻ γ |
| 0.35 ± 0.14 ± 0.03 | 11 | ³ LEES | 13Q BABR | e ⁺ e ⁻ → K ⁺ K ⁻ γ |

¹ sin φ > 0.

² sin φ < 0.

³ Interference with non-resonant K⁺K⁻ production not taken into account.

ψ(2S) BRANCHING RATIOS $\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ Γ_1 / Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-------------------|----------|-------------------------------|
| 0.9785 ± 0.0013 OUR AVERAGE | | | |
| 0.9779 ± 0.0015 | ¹ BAI | 02B BES2 | e ⁺ e ⁻ |
| 0.981 ± 0.003 | ¹ LUTH | 75 MRK1 | e ⁺ e ⁻ |

¹ Includes cascade decay into J/ψ(1S).

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$ Γ_2 / Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|-------------------------------------|----------|-------------------------------|
| 0.0173 ± 0.0014 OUR AVERAGE | Error includes scale factor of 1.5. | | |
| 0.0166 ± 0.0010 | ^{1,2} SETH | 04 RVUE | e ⁺ e ⁻ |
| 0.0199 ± 0.0019 | ¹ BAI | 02B BES2 | e ⁺ e ⁻ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | |
|---------------|-------------------|---------|-------------------------------|
| 0.029 ± 0.004 | ¹ LUTH | 75 MRK1 | e ⁺ e ⁻ |
|---------------|-------------------|---------|-------------------------------|

¹ Included in Γ(hadrons)/Γ_{total}.

² Using B(ψ(2S) → ℓ⁺ℓ⁻) = (0.73 ± 0.04)% from RPP-2002 and R = 2.28 ± 0.04 determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_3/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 10.58 ± 1.62 | 2.9 M | ¹ LIBBY | 09 | CLEO $\psi(2S) \rightarrow \text{hadrons}$ |

¹ Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow XJ/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma\eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma\chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+\ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 1.025 ± 0.288 | 200 k | ¹ LIBBY | 09 | CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$ |

¹ Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

 $\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_4/Γ_3

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 9.7 ± 2.6 ± 1.6 | 2.9 M | LIBBY | 09 | CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$ |

 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|---------------------|-------------|------------------------------------|
| 0.154 ± 0.015 | ¹ MENDEZ | 08 | CLEO $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|-------------------|-----|------------------------------------|
| 0.169 ± 0.026 | ² ADAM | 05A | CLEO $e^+e^- \rightarrow \psi(2S)$ |
|---------------|-------------------|-----|------------------------------------|

¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ}\gamma)$, $B(\eta_c\gamma)$ from ATHAR 04 and $B(\ell^+\ell^-)$ from PDG 04. Superseded by MENDEZ 08.

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| 79.3 ± 1.7 OUR FIT | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------|----------------------|----|---------------|
| 88 ± 13 | ¹ FELDMAN | 77 | RVUE e^+e^- |
|---------|----------------------|----|---------------|

¹ From an overall fit assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> |
|---|--------------------|
| 80 ± 6 OUR FIT | |

 $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_7/Γ_6

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|--------------------|-------------|----------------|
| 1.00 ± 0.08 OUR FIT | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|----------|-----|---------------|
| 0.89 ± 0.16 | BOYARSKI | 75C | MRK1 e^+e^- |
|-------------|----------|-----|---------------|

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------------------|---------|-------------------------------|
| 31 ± 4 OUR FIT | | | |
| 30.8 ± 2.1 ± 3.8 | ¹ ABLIKIM | 06W BES | $e^+e^- \rightarrow \psi(2S)$ |

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

————— **DECAYS INTO $J/\psi(1S)$ AND ANYTHING** —————

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------|----------|--------------------------------------|
| 0.614 ± 0.006 OUR FIT | | | | |
| 0.55 ± 0.07 OUR AVERAGE | | | | |
| 0.51 ± 0.12 | | BRANDELIK | 79C DASP | $e^+e^- \rightarrow \mu^+\mu^-X$ |
| 0.57 ± 0.08 | | ABRAMS | 75B MRK1 | $e^+e^- \rightarrow \mu^+\mu^-X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.6254 ± 0.0016 ± 0.0155 | 1.1M | ¹ MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \ell^+\ell^-X$ |
| 0.5950 ± 0.0015 ± 0.0190 | 151k | ADAM | 05A CLEO | Repl. by MENDEZ 08 |


¹ Not independent from other measurements of MENDEZ 08.

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$
 $\Gamma_6/\Gamma_9 = \Gamma_6/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-----------|-------------------------|----------|---|
| 1.291 ± 0.026 OUR FIT | | | | |
| 1.28 ± 0.04 OUR AVERAGE | | | | Error includes scale factor of 1.6. See the ideogram below. |
| 1.22 ± 0.02 ± 0.05 | 5097 ± 73 | ¹ ANDREOTTI | 05 E835 | $\rho\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$ |
| 1.28 ± 0.03 ± 0.02 | | ¹ AMBROGIANI | 00A E835 | $\rho\bar{p} \rightarrow \psi(2S)$ |
| 1.44 ± 0.08 ± 0.02 | | ¹ ARMSTRONG | 97 E760 | $\bar{p}p \rightarrow \psi(2S)$ |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

WEIGHTED AVERAGE
 1.28 ± 0.04 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\mu^+ \mu^-) / \Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_7 / \Gamma_9 = \Gamma_7 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------|
| 0.0130 ± 0.0010 OUR FIT | | | |
| 0.014 ± 0.003 | HILGER | 75 | SPEC $e^+ e^-$ |

$\Gamma(J/\psi(1S)\text{neutrals}) / \Gamma_{\text{total}}$

Γ_{10} / Γ

| VALUE | DOCUMENT ID |
|---|-------------|
| 0.2538 ± 0.0032 OUR FIT | |

$\Gamma(J/\psi(1S)\pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{11} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| 0.3468 ± 0.0030 OUR FIT | | | | |
| 0.348 ± 0.005 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |


| | | | | | |
|--------------------------------|------|---------|-----|------|--|
| $0.3498 \pm 0.0002 \pm 0.0045$ | 20M | ABLIKIM | 13R | BES3 | $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ |
| $0.3504 \pm 0.0007 \pm 0.0077$ | 565k | MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$ |
| 0.323 ± 0.014 | | BAI | 02B | BES2 | $e^+ e^-$ |
| 0.32 ± 0.04 | | ABRAMS | 75B | MRK1 | $e^+ e^- \rightarrow J/\psi \pi^+ \pi^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------------|-----|-------------------|-----|------|--------------------|
| $0.3354 \pm 0.0014 \pm 0.0110$ | 60k | ¹ ADAM | 05A | CLEO | Repl. by MENDEZ 08 |
|--------------------------------|-----|-------------------|-----|------|--------------------|

¹ Not independent from other values reported by ADAM 05A.

WEIGHTED AVERAGE
 0.348 ± 0.005 (Error scaled by 1.3)



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_6/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|---------------------|------|---------------|
| 0.0229 ± 0.0005 OUR FIT | | | |
| 0.0252 ± 0.0028 ± 0.0011 | ¹ AUBERT | 02B | BABR e^+e^- |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_7/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|------------------------|------|---|
| 0.0230 ± 0.0017 OUR FIT | | | |
| 0.0228 ± 0.0018 OUR AVERAGE | | | |
| 0.0230 ± 0.0020 ± 0.0012 | ¹ AAIJ | 16Y | LHCB $\Lambda_b^0 \rightarrow \psi(2S)X$ |
| 0.0216 ± 0.0026 ± 0.0014 | ² AUBERT | 02B | BABR e^+e^- |
| 0.0327 ± 0.0077 ± 0.0072 | ² GRIBUSHIN | 96 | FMPS 515 $\pi^- \text{Be} \rightarrow 2\mu X$ |

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.

² Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_8/Γ_{11}

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|--------------|
| 8.8 ± 1.1 OUR FIT | | | |
| 8.73 ± 1.39 ± 1.57 | BAI | 02 | BES e^+e^- |

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{11}/Γ_9

VALUE EVTS DOCUMENT ID TECN COMMENT

0.5645 ± 0.0026 OUR FIT

0.554 ± 0.008 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

| | | | | | |
|--------------------------|------|------------------------|-----|------|--|
| 0.5604 ± 0.0009 ± 0.0062 | 565k | MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$ |
| 0.525 ± 0.009 ± 0.022 | 4k | ANDREOTTI | 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.536 ± 0.007 ± 0.016 | 20k | ^{1,2} ABLIKIM | 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |
| 0.496 ± 0.037 | | ARMSTRONG | 97 | E760 | $\bar{p}p \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •


| | | | | | |
|--------------------------|-----|------|-----|------|--------------------|
| 0.5637 ± 0.0027 ± 0.0046 | 60k | ADAM | 05A | CLEO | Repl. by MENDEZ 08 |
|--------------------------|-----|------|-----|------|--------------------|

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$.

WEIGHTED AVERAGE

0.554 ± 0.008 (Error scaled by 1.3)



$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})/\Gamma_{11}$

VALUE DOCUMENT ID TECN COMMENT

0.732 ± 0.008 OUR FIT

| | | | | | |
|--------------------|--|-----------|----|------|-----------|
| 0.73 ± 0.09 | | TANENBAUM | 76 | MRK1 | $e^+ e^-$ |
|--------------------|--|-----------|----|------|-----------|

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}} \qquad \Gamma_{12}/\Gamma$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.1824 ± 0.0031 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|-------|---------------------|-----|------|---|
| 0.1769 ± 0.0008 ± 0.0053 | 61k | ¹ MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$ |
| 0.1652 ± 0.0014 ± 0.0058 | 13.4k | ² ADAM | 05A | CLEO | Repl. by MENDEZ 08 |

¹ Not independent from other measurements of MENDEZ 08.² Not independent from other values reported by ADAM 05A.
$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything}) \qquad \Gamma_{12}/\Gamma_9$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.2968 ± 0.0031 OUR FIT**0.320 ± 0.012 OUR AVERAGE**

| | | | | | |
|-----------------------|-----------|------------|-----|------|---------------------------------|
| 0.300 ± 0.008 ± 0.022 | 1655 ± 44 | ANDREOTTI | 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.328 ± 0.013 ± 0.008 | | AMBROGIANI | 00A | E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.323 ± 0.033 | | ARMSTRONG | 97 | E760 | $\bar{p}p \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|-------|--------|-----|------|---|
| 0.2829 ± 0.0012 ± 0.0056 | 61k | MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$ |
| 0.2776 ± 0.0025 ± 0.0043 | 13.4k | ADAM | 05A | CLEO | Repl. by MENDEZ 08 |

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-) \qquad \Gamma_{12}/\Gamma_{11}$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.526 ± 0.008 OUR FIT**0.513 ± 0.022 OUR AVERAGE** Error includes scale factor of 2.2.

| | | | | | |
|--------------------------|-----|----------------------|-----|------|---|
| 0.5047 ± 0.0022 ± 0.0102 | 61k | MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$ |
| 0.570 ± 0.009 ± 0.026 | 14k | ¹ ABLIKIM | 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|-----|------------------------|-----|------|---------------------------------|
| 0.4924 ± 0.0047 ± 0.0086 | 73k | ^{2,3} ADAM | 05A | CLEO | Repl. by MENDEZ 08 |
| 0.571 ± 0.018 ± 0.044 | | ⁴ ANDREOTTI | 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.53 ± 0.06 | | TANENBAUM | 76 | MRK1 | $e^+ e^-$ |
| 0.64 ± 0.15 | | ⁵ HILGER | 75 | SPEC | $e^+ e^-$ |

¹ From a fit to the J/ψ recoil mass spectra.² Not independent from other values reported by ADAM 05A.³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.⁴ Not independent from other values reported by ANDREOTTI 05.⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.
$$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|


33.7 ± 0.5 OUR FIT**32.9 ± 1.7 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

| | | | | | |
|---------------------|-------|------------------------|-----|------|---|
| 33.75 ± 0.17 ± 0.86 | 68.2k | ABLIKIM | 12M | BES3 | $e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$ |
| 29.8 ± 0.9 ± 2.3 | 5.7k | BAI | 04I | BES2 | $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 25.5 ± 2.9 | 386 | ¹ OREGLIA | 80 | CBAL | $e^+ e^- \rightarrow J/\psi 2\gamma$ |
| 45 ± 12 | 17 | ² BRANDELIK | 79B | DASP | $e^+ e^- \rightarrow J/\psi 2\gamma$ |
| 42 ± 6 | 164 | ² BARTEL | 78B | CNTR | $e^+ e^-$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | | | |
|------------------------|-------|---------------------|-----|------|---|
| $34.3 \pm 0.4 \pm 0.9$ | 18.4k | ³ MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ |
| $32.5 \pm 0.6 \pm 1.1$ | 2.8k | ⁴ ADAM | 05A | CLEO | Repl. by MENDEZ 08 |
| 43 ± 8 | 44 | TANENBAUM | 76 | MRK1 | $e^+ e^-$ |

- ¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.
² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
³ Not independent from other measurements of MENDEZ 08.
⁴ Not independent from other values reported by ADAM 05A.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$


Γ_{13}/Γ_9

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------------|--------------|----------------|------|--|
| 0.0549 ± 0.0008 | | | | OUR FIT |
| 0.058 ± 0.007 | | | | OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below. |
| $0.050 \pm 0.006 \pm 0.003$ | 298 ± 20 | ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.072 ± 0.009 | | AMBROGIANI 00A | E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.061 ± 0.015 | | ARMSTRONG 97 | E760 | $\bar{p}p \rightarrow \psi(2S)$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | | | |
|--------------------------------|-------|---------------------|-----|------|---|
| $0.0549 \pm 0.0006 \pm 0.0009$ | 18.4k | ¹ MENDEZ | 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ |
| $0.0546 \pm 0.0010 \pm 0.0007$ | 2.8k | ADAM | 05A | CLEO | Repl. by MENDEZ 08 |

- ¹ Not independent from other measurements of MENDEZ 08.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{13}/Γ_{11}

| VALUE | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|-------|---------------------------|------|---|
| 0.0972 ± 0.0014 | | | | OUR FIT |
| 0.0979 ± 0.0018 | | | | OUR AVERAGE |
| $0.0979 \pm 0.0010 \pm 0.0015$ | 18.4k | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+\ell^-\eta$ |
| $0.098 \pm 0.005 \pm 0.010$ | 2k | ¹ ABLIKIM 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |
| 0.091 ± 0.021 | | ² HIMEL 80 | MRK2 | $e^+e^- \rightarrow \psi(2S)X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $0.0968 \pm 0.0019 \pm 0.0013$ | 2.8k | ³ ADAM 05A | CLEO | Repl. by MENDEZ 08 |
| $0.095 \pm 0.007 \pm 0.007$ | | ⁴ ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{14}/Γ

| VALUE (units 10^{-4}) | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|------|--|
| 12.68 ± 0.32 | | | | OUR AVERAGE |
| $12.6 \pm 0.2 \pm 0.3$ | 4.1k | ABLIKIM 12M | BES3 | $e^+e^- \rightarrow \ell^+\ell^-2\gamma$ |
| $13.3 \pm 0.8 \pm 0.3$ | 530 | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+\ell^-2\gamma$ |
| $14.3 \pm 1.4 \pm 1.2$ | 280 | BAI 04I | BES2 | $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 14 ± 6 | 7 | HIMEL 80 | MRK2 | e^+e^- |
| $9 \pm 2 \pm 1$ | 23 | ¹ OREGLIA 80 | CBAL | $\psi(2S) \rightarrow J/\psi 2\gamma$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |

13 ± 1 ± 1 88 ADAM 05A CLEO Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13}+0.343\Gamma_{154}+0.190\Gamma_{155})$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------------|-----|---------------------|-----|---|
| 0.213 \pm 0.012 \pm 0.003 | 527 | ¹ MENDEZ | 08 | CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$ |
| 0.22 \pm 0.02 \pm 0.01 | | ² ADAM | 05A | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$ |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{14}/Γ_{11}

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------------|-----|---------------------|-----|---|
| 0.380 \pm 0.022 \pm 0.005 | 527 | ¹ MENDEZ | 08 | CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$ |
| 0.39 \pm 0.04 \pm 0.01 | | ² ADAM | 05A | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$ |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

———— HADRONIC DECAYS ————

$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$

Γ_{15}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

8.6 \pm 1.3 OUR AVERAGE

| | | | | |
|-------------------------|-----|-----------------|-----|--|
| 9.0 \pm 1.5 \pm 1.3 | 3k | ¹ GE | 11 | CLEO $\psi(2S) \rightarrow \pi^0$ anything |
| 8.4 \pm 1.3 \pm 1.0 | 11k | ABLIKIM | 10B | BES3 $\psi(2S) \rightarrow \pi^0 h_c$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----------------------------------|--------|-----|--|
| seen | 92 ⁺²³ ₋₂₂ | ADAMS | 09 | CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$ |
| seen | 1282 | DOBBS | 08A | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |
| seen | 168 \pm 40 | ROSNER | 05 | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$

Γ_{16}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|


35 \pm 16

| | | | | |
|---|----------|----|------|------------------------------|
| 6 | FRANKLIN | 83 | MRK2 | $e^+e^- \rightarrow$ hadrons |
|---|----------|----|------|------------------------------|

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{17}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------------|---|-----------|--|
| 29 ± 10 | OUR AVERAGE | Error includes scale factor of 4.7. See the ideogram below. | | |
| 24.9 ± 0.7 ± 3.6 | 2173 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 127 ± 12 ± 2 | 410 | ¹ AUBERT | 07AU BABR | 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$ |
| 26.1 ± 0.7 ± 3.0 | 1703 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$ |
| 30 ± 8 | 42 | FRANKLIN | 83 MRK2 | e^+e^- |

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$ Γ_{18}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|------|-------------|---|---------|
| 2.55 ± 0.73 ± 0.47 | 112 ± 31 | BAI | 04C BES2 | $\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$ | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <2.3 | 90 | BAI | 98J BES | e^+e^- | |

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{21}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|--------------------|----------|--|
| 2.94±0.08 OUR FIT | | | | |
| 3.02±0.08 OUR AVERAGE | | | | |
| 3.05±0.02±0.12 | 19k | ABLIKIM | 18T BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 3.08±0.05±0.18 | 4.5k | ¹ DOBBS | 14 | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 3.36±0.09±0.25 | 1.6k | ABLIKIM | 07C BES | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 2.87±0.12±0.15 | 557 | PEDLAR | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 1.4 ±0.8 | 4 | BRANDELIK | 79C DASP | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 2.3 ±0.7 | | FELDMAN | 77 MRK1 | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{21}/Γ_{11}

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|--------|--|
| 8.49±0.23 OUR FIT | | | |
| 6.98±0.49±0.97 | BAI | 01 BES | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{22}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 3.06±0.06±0.14 | 6k | ABLIKIM | 18T BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$ |

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$ Γ_{23}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|------------------|--------|--|
| 12.8±1.0±3.4 | 157 | ¹ BAI | 01 BES | $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons |

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|----------|---|
| < 0.29 | 90 | ¹ ABLIKIM | 13F BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ |
| ••• | | | | We do not use the following data for averages, fits, limits, etc. ••• |
| <12 | 90 | ² ABLIKIM | 07H BES2 | $e^+e^- \rightarrow \psi(2S)$ |

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{25}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|----------------------|----------|---|
| 2.48±0.34±0.19 | | 60 | ¹ ABLIKIM | 13F BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ |
| ••• | | | | | We do not use the following data for averages, fits, limits, etc. ••• |
| <4.9 | 90 | | ² ABLIKIM | 07H BES2 | $e^+e^- \rightarrow \psi(2S)$ |

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{26}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 1.0±0.1±0.1 | 74.0 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^-$ |

$\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{27}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|-----------|-------------------------------|
| 6.3±0.5±0.5 | 1011 | ABLIKIM | 19AU BES3 | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|--|
| 1.8±0.3±0.3 | 45.8 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$ |

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 2.8±0.4±0.5 | 73.4 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$ |

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{30}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|---|-----------|--|
| 3.81±0.13 OUR AVERAGE | | | Error includes scale factor of 1.4. See the ideogram below. | | |
| 3.97±0.02±0.12 | | 31k | ABLIKIM | 17L BES3 | $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| 3.71±0.05±0.15 | | 6.5k | ¹ DOBBS | 17 | $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| 3.39±0.20±0.32 | | 337 | ABLIKIM | 07C BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| 6.4 ±1.8 ±0.1 | | | ² AUBERT | 07BD BABR | 10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$ |
| 3.28±0.23±0.25 | | 208 | PEDLAR | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 3.75±0.09±0.23 | | 1.9k | ^{1,3} DOBBS | 14 | $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ |
| 1.81±0.20±0.27 | | 80 | ⁴ BAI | 01 BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| < 4 | | 90 | FELDMAN | 77 MRK1 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |


¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by DOBBS 17.

⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

WEIGHTED AVERAGE
 3.81 ± 0.13 (Error scaled by 1.4)



$\Gamma(\Lambda\bar{\Sigma}^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{31}/Γ**

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|----------|---------------------------------------|
| $1.40 \pm 0.03 \pm 0.13$ | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda\bar{\Sigma}^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{32}/Γ**

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|----------|---------------------------------------|
| $1.54 \pm 0.04 \pm 0.13$ | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$ **Γ_{33}/Γ**

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|--|------|-----------------------|--|
| $1.23 \pm 0.23 \pm 0.08$ | 30 | ¹ DOBBS 17 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{34}/Γ**

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|---|
| $1.67 \pm 0.13 \pm 0.12$ | 276 | ¹ ABLIKIM 13D | BES3 | $\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$ |

¹ Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda \gamma) = 100\%$.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|------|--|
| 2.32 ± 0.12 OUR AVERAGE | | | | |
| $2.31 \pm 0.06 \pm 0.10$ | 1.9k | ¹ DOBBS 17 | | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.57 \pm 0.44 \pm 0.68$ | 35 | PEDLAR 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.51 \pm 0.15 \pm 0.16$ 281 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹Using CLEO-c data but not authored by the CLEO Collaboration.

²Superseded by DOBBS 17.

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

Γ_{36}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|-------------------------------------|------|---|
| 2.35 ± 0.09 | OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $2.44 \pm 0.03 \pm 0.11$ | 7k | ABLIKIM | 17L | BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.22 \pm 0.05 \pm 0.11$ | 2.6k | ¹ DOBBS | 17 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.35 \pm 0.36 \pm 0.32$ | 59 | ABLIKIM | 07C | BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.63 \pm 0.35 \pm 0.21$ | 58 | PEDLAR | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $2.25 \pm 0.11 \pm 0.16$ | 439 | ^{1,2} DOBBS | 14 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $1.2 \pm 0.4 \pm 0.4$ | 8 | ³ BAI | 01 | BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹Using CLEO-c data but not authored by the CLEO Collaboration.

²Superseded by DOBBS 17.

³Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

Γ_{37}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|--------------------|------------------|------|---|
| 8.5 ± 0.7 | OUR AVERAGE | | | |
| $8.4 \pm 0.5 \pm 0.5$ | 1.5k | ABLIKIM | 16L | BES3 $\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$ |
| $11 \pm 3 \pm 3$ | 14 | ¹ BAI | 01 | BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

Γ_{38}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $8.5 \pm 0.6 \pm 0.6$ | 1.4K | ABLIKIM | 16L | BES3 $\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$ |

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

Γ_{39}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $0.69 \pm 0.05 \pm 0.05$ | 2.2k | ABLIKIM | 17E | BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

Γ_{40}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|--------------------|-------------------------------------|-------------|------|--|
| 2.87 ± 0.11 | OUR AVERAGE | Error includes scale factor of 1.1. | | | |
| $3.03 \pm 0.05 \pm 0.14$ | 3.6k | ¹ DOBBS | 17 | | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.78 \pm 0.05 \pm 0.14$ | 5k | ABLIKIM | 16L | BES3 | $\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$ |
| $3.03 \pm 0.40 \pm 0.32$ | 67 | ABLIKIM | 07C | BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| $2.38 \pm 0.30 \pm 0.21$ | 63 | PEDLAR | 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$2.66 \pm 0.12 \pm 0.20$ 548 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

$0.94 \pm 0.27 \pm 0.15$ 12 ³ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

< 2 90 FELDMAN 77 MRK1 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹Using CLEO-c data but not authored by the CLEO Collaboration.

²Superseded by DOBBS 17.

³Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

2.3 ± 0.4 OUR AVERAGE Error includes scale factor of 4.2.2.73 ± 0.03 ± 0.13 11k ABLIKIM 17E BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons1.97 ± 0.06 ± 0.11 1.2k ¹ DOBBS 17 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons2.75 ± 0.64 ± 0.61 19 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.02 ± 0.19 ± 0.15 112 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17. $\Gamma(\Xi(1530)^0 \Xi(1530)^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|------|---------|
|--------------------------|-----|------|-------------|------|---------|

5.2 ± 0.3 +3.2 -1.2 527 ¹ ABLIKIM 13S BES3 $\psi(2S) \rightarrow \eta p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 32 90 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ < 8.1 90 ² BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow$
hadrons
hadrons¹ With $N(1535)$ decaying to $p\eta$.² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Xi(1530)^- \Xi(1530)^+)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

11.45 ± 0.40 ± 0.59 5k ABLIKIM 19AT BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$
hadrons $\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

7.0 ± 1.1 ± 0.4 399 ABLIKIM 19AT BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons $\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{43}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

3.86 ± 0.27 ± 0.32 236 ABLIKIM 15I BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$
 $K^- \Lambda \Xi^+ + \text{c.c.}$ $\Gamma(\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

5.21 ± 1.48 ± 0.57 74 ABLIKIM 15I BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$
 $K^- \Lambda \Xi^+ + \text{c.c.}$ $\Gamma(\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{47}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

12.03 ± 2.94 ± 1.22 136 ABLIKIM 15I BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow$
 $K^- \Lambda \Xi^+ + \text{c.c.}$

$\Gamma(K^- \Sigma^0 \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{48}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $3.67 \pm 0.33 \pm 0.28$ | 142 | ABLIKIM 15i | BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \Xi^+ + \text{c.c.}$ |

$\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{49}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------|--------------------|------|---|
| $0.52 \pm 0.03 \pm 0.03$ | | 326 | ¹ DOBBS | 17 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------------|----|----------------------|-----|------|---|
| $0.47 \pm 0.09 \pm 0.05$ | 27 | ^{1,2} DOBBS | 14 | | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <1.5 | 90 | ABLIKIM | 12Q | BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <1.6 | 90 | PEDLAR | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <0.73 | 90 | ³ BAI | 01 | BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{50}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|------|--|
| 1.53 ± 0.07 OUR AVERAGE | | | | |
| $1.65 \pm 0.03 \pm 0.15$ | 4.5k | ABLIKIM 13A | BES3 | $\psi(2S) \rightarrow p \bar{p} \pi^0$ |
| $1.54 \pm 0.06 \pm 0.06$ | 948 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \pi^0 p \bar{p}$ |
| $1.32 \pm 0.10 \pm 0.15$ | 256 | ¹ ABLIKIM 05E | BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \gamma \gamma$ |
| 1.4 ± 0.5 | 9 | FRANKLIN 83 | MRK2 | $e^+ e^-$ |

¹ Computed using $B(\pi^0 \rightarrow \gamma \gamma) = (98.80 \pm 0.03)\%$.

$\Gamma(N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{51}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|------|--|
| 6.42 ± 0.20 $^{+1.78}_{-1.28}$ | 1.9k | ¹ ABLIKIM 13A | BES3 | $\psi(2S) \rightarrow p \bar{p} \pi^0$ |

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

$\Gamma(N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{52}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-------------------------------------|
| 7.3 $^{+1.7}_{-1.5}$ OUR AVERAGE | | | | Error includes scale factor of 2.5. |

| | | | | |
|------------------------------------|------|---------------------------|------|--|
| 3.58 ± 0.25 $^{+1.59}_{-0.84}$ | 1.1k | ¹ ABLIKIM 13A | BES3 | $\psi(2S) \rightarrow p \bar{p} \pi^0$ |
| $8.1 \pm 0.7 \pm 0.3$ | 474 | ² ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \pi^0 p \bar{p}$ |

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

² From a fit of the $p \bar{p}$ and $p \pi^0$ mass distributions to a combination of $N(1440) \bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

$\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{53}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $0.64 \pm 0.05^{+0.22}_{-0.17}$ | 0.2k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $2.47 \pm 0.28^{+0.99}_{-0.97}$ | 0.7k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $3.76 \pm 0.28^{+1.37}_{-1.66}$ | 1.1k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $1.79 \pm 0.10^{+0.24}_{-0.71}$ | 0.5k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{57}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $2.62 \pm 0.28^{+1.12}_{-0.64}$ | 0.9k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{58}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|----------------------|------|--|
| $2.13 \pm 0.08^{+0.40}_{-0.30}$ | 0.8k | ¹ ABLIKIM | 13A | BES3 $\psi(2S) \rightarrow \rho\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{59}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|------------------------|------|---|
| $1.1 \pm 0.4 \pm 0.1$ | 76 | ¹ ALEXANDER | 10 | CLEO $\psi(2S) \rightarrow \pi^0 \rho\bar{p}$ |

¹ From a fit of the $\rho\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

$\Gamma(\eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{60}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------------|------------------------|----------|--|
| 6.0±0.4 OUR AVERAGE | | | | |
| 6.4±0.2±0.6 | 679 | ¹ ABLIKIM | 13S BES3 | $\psi(2S) \rightarrow \eta\rho\bar{p}$ |
| 5.6±0.6±0.3 | 154 | ¹ ALEXANDER | 10 CLEO | $\psi(2S) \rightarrow \eta\rho\bar{p}$ |
| 5.8±1.1±0.7 | 44.8 ± 8.5 | ² ABLIKIM | 05E BES2 | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\gamma\gamma$ |
| 8 ±3 ±3 | 9.8 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$ |

¹ With $N(1535)$ decaying to $p\eta$.² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$. $\Gamma(\eta f_0(2100) \rightarrow \eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{61}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|------------------------|---------|--|
| 1.2±0.4±0.1 | 31 | ¹ ALEXANDER | 10 CLEO | $\psi(2S) \rightarrow \eta\rho\bar{p}$ |

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$. $\Gamma(N(1535)\bar{p} \rightarrow \eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{62}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|------------------------|---------|--|
| 4.4±0.6±0.3 | 123 | ¹ ALEXANDER | 10 CLEO | $\psi(2S) \rightarrow \eta\rho\bar{p}$ |

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$. $\Gamma(\omega\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------------|------------------|---------|--|
| 0.69±0.21 OUR AVERAGE | | | | |
| 0.6 ±0.2 ±0.2 | 21.2 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$ |
| 0.8 ±0.3 ±0.1 | 14.9 ± 0.1 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. $\Gamma(\eta'\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{64}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|----------|---|
| 1.10±0.10±0.08 | 491 | ¹ ABLIKIM | 19N BES3 | $\psi(2S) \rightarrow \eta'\rho\bar{p}$ |

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels. $\Gamma(\phi\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{65}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|-----------|---|
| 6.06±0.38±0.48 | | 753 | ABLIKIM | 19AO BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+K^-$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

| | | | | |
|-----|----|------------------|---------|---|
| <24 | 90 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+K^-$ |
| <26 | 90 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow K^+K^-\rho\bar{p}$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

| $\Gamma(\phi X(1835) \rightarrow \phi p \bar{p})/\Gamma_{\text{total}}$ | | | Γ_{66}/Γ | | |
|---|------------|--------------------|----------------------|---|--|
| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $<1.82 \times 10^{-7}$ | 90 | ABLIKIM | 19AO BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow p \bar{p} K^+ K^-$ | |

| $\Gamma(\pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}}$ | | | Γ_{67}/Γ | | |
|---|-------------|------------------------|----------------------|---|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 6.0 ± 0.4 OUR AVERAGE | | | | | |
| $5.9 \pm 0.2 \pm 0.4$ | 904.5 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \pi^+ \pi^-$ | |
| 8 ± 2 | | ¹ TANENBAUM | 78 MRK1 | e^+e^- | |

¹ Assuming entirely strong decay.

| $\Gamma(p \bar{n} \pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ | | | Γ_{68}/Γ | | |
|---|-------------|--------------------|----------------------|--------------------------------------|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 2.48 ± 0.17 OUR AVERAGE | | | | | |
| $2.45 \pm 0.11 \pm 0.21$ | 851 | ABLIKIM | 06i BES2 | $e^+e^- \rightarrow p \pi^- X$ | |
| $2.52 \pm 0.12 \pm 0.22$ | 849 | ABLIKIM | 06i BES2 | $e^+e^- \rightarrow \bar{p} \pi^+ X$ | |

| $\Gamma(p \bar{n} \pi^- \pi^0)/\Gamma_{\text{total}}$ | | | Γ_{69}/Γ | | |
|---|--------------|--------------------|----------------------|--------------------------------------|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $3.18 \pm 0.50 \pm 0.50$ | 135 ± 21 | ABLIKIM | 06i BES2 | $e^+e^- \rightarrow p \pi^- \pi^0 X$ | |

| $\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ | | | Γ_{71}/Γ | | |
|--|------------|--------------------|----------------------|--|--|
| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| <1.6 | 90 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ | |

| $\Gamma(\eta \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ | | | Γ_{72}/Γ | | |
|---|-------------|---------------------|----------------------|--|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $9.5 \pm 0.7 \pm 1.5$ | | ¹ BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $10.3 \pm 0.8 \pm 1.4$ | 201.7 | ² BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$ | |
| $8.1 \pm 1.4 \pm 1.6$ | 50.0 | ² BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$ | |

¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

² Not independent from other values reported by BRIERE 05.

| $\Gamma(2(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}$ | | | Γ_{73}/Γ | | |
|--|-------------|---------------------|----------------------|--|--|
| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $1.2 \pm 0.6 \pm 0.1$ | 16 | ¹ AUBERT | 07AU BABR | $10.6 e^+e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$ | |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$. | | | | | |


| $\Gamma(\eta' \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ | | | Γ_{75}/Γ | | |
|---|-------------|--------------------|----------------------|---|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $4.5 \pm 1.6 \pm 1.3$ | 12.8 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$ | |

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{76}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|---------------------|-----------|--|
| 7.3±1.2 OUR AVERAGE | | | | Error includes scale factor of 2.1. See the ideogram below. |
| 8.4±0.5±1.2 | 386 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 12.2±2.2±0.7 | 37 | ¹ AUBERT | 07AU BABR | 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$ |
| 8.2±0.5±0.7 | 391 | BRIERE | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| 4.8±0.6±0.7 | 100 ± 22 | ² BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV. | | | | |
| ² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. | | | | |

WEIGHTED AVERAGE
7.3±1.2 (Error scaled by 2.1)



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{77}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|---------|--------------------|----------|---|
| 4.0 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 5.1 ± 0.6 ± 0.8 | 202 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 4.18 ^{+0.43} _{-0.42} ± 0.92 | 170 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 3.2 ± 0.6 ± 0.5 | 61 ± 11 | ^{1,2} BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 5.2 ± 0.8 ± 1.0 | | ¹ BAI | 99C BES | Repl. by BAI 03B |
| ¹ Assuming $B(b_1 \rightarrow \omega\pi) = 1$. | | | | |
| ² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. | | | | |

| $\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{78}/Γ |
|---|------|-------------|------|---------|--------------------------------|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $2.35^{+0.47}_{-0.42} \pm 0.40$ | 45 | ADAM | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

| $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ | | | | | Γ_{79}/Γ |
|---|-----|-------------|------------------|----------|---|
| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 2.2 ± 0.4 OUR AVERAGE | | | | | |
| $2.3 \pm 0.5 \pm 0.4$ | | 57 | ABLIKIM | 07D BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| $2.05 \pm 0.41 \pm 0.38$ | | 62 ± 12 | BAI | 04C BES2 | $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 1.5 | | 90 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| < 1.7 | | 90 | BAI | 98J BES | Repl. by BAI 03B |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

| $\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ | | | | | Γ_{82}/Γ |
|---|------|-------------|------------------------|-----------|--|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 7.3 ± 0.5 OUR AVERAGE | | | | | |
| $8.1 \pm 1.3 \pm 0.3$ | | 133 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| $7.1 \pm 0.3 \pm 0.4$ | | 817.2 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| 16 ± 4 | | | ¹ TANENBAUM | 78 MRK1 | $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $11.0 \pm 1.9 \pm 0.2$ | | 85 | ² AUBERT | 07AK BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$ | | | | | Γ_{84}/Γ |
|--|-------|-------------|------|---------|--|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $2.2 \pm 0.2 \pm 0.4$ | 223.8 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

| $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ | | | | | Γ_{85}/Γ |
|---|-----|-------------|-------------|---------|--|
| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| $1.86 \pm 0.32 \pm 0.43$ | | 93 ± 16 | BAI | 04C | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 1.2 | | 90 | BAI | 98J BES | $e^+ e^-$ |

| $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ | | | | | Γ_{86}/Γ |
|--|------|---------------------|-----------|---------|---|
| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $1.3 \pm 0.7 \pm 0.1$ | 7 | ¹ AUBERT | 07AU BABR | 10.6 | $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$ |
| ¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV. | | | | | |

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{87} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--------------------------------|
| 10.0 ± 2.5 ± 1.8 | 65 | ABLIKIM | 07D BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(K_1(1270)^\pm K^\mp) / \Gamma_{\text{total}}$ Γ_{89} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------------|---------|-----------|
| 10.0 ± 1.8 ± 2.1 | ¹ BAI | 99C BES | $e^+ e^-$ |

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{90} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|--------|-------------|----------|--------------------------------|
| 2.20 ± 0.25 ± 0.37 | 83 ± 9 | ABLIKIM | 050 BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\rho^0 \rho^0) / \Gamma_{\text{total}}$ Γ_{91} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 0.5 ± 0.1 ± 0.2 | 61.1 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \rho^0 \pi^+ \pi^-$ |

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{92} / Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------|------|-----------|
| 6.7 ± 2.5 | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{93} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|--------------|---------|---|
| 2.4 ± 0.6 OUR AVERAGE | Error includes scale factor of 2.2. | | | |
| 2.2 ± 0.2 ± 0.2 | 308 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$ |
| 4.5 ± 1.0 | | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{94} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|--------------|---------|---|
| 2.2 ± 0.6 OUR AVERAGE | Error includes scale factor of 1.4. | | | |
| 2.0 ± 0.2 ± 0.4 | 285.5 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$ |
| 4.2 ± 1.5 | | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{95} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------|---------------------|-----------|--|
| 12.6 ± 0.9 OUR AVERAGE | | | | |
| 18.9 ± 5.7 ± 0.3 | 32 | ¹ AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$ |
| 11.7 ± 1.0 ± 1.5 | 597 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| 12.7 ± 0.5 ± 1.0 | 711.6 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{96} / Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $5.9 \pm 2.0 \pm 0.9$ | 19 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{97} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $8.6 \pm 1.3 \pm 1.8$ | 238 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{98} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $9.6 \pm 2.2 \pm 1.7$ | 133 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{99} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $7.3 \pm 2.2 \pm 1.4$ | 78 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{100} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| $6.1 \pm 1.3 \pm 1.2$ | 125 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(\eta K^+ K^-, \text{no } \eta\phi) / \Gamma_{\text{total}}$ Γ_{101} / Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------|-------------|------|---|
| $3.08 \pm 0.29 \pm 0.25$ | 0.3k | ¹ | ABLIKIM | 12L | BES3 $\psi(2S) \rightarrow K^+ K^- \gamma \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|--------|----|------|--|
| <13 | 90 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
|-----|----|--------|----|------|--|

¹ Excluding $\eta\phi$.

$\Gamma(\omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{102} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|------|---|
| 1.62 ± 0.11 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| $1.56 \pm 0.04 \pm 0.11$ | 2.8k | ABLIKIM | 14G | BES3 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $2.38 \pm 0.37 \pm 0.29$ | 78 | ABLIKIM | 06G | BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.9 \pm 0.3 \pm 0.3$ | 76.8 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.5 \pm 0.3 \pm 0.2$ | 23 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\omega K^*(892)^+ K^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{103} / Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|--|
| 20.7 ± 2.6 OUR AVERAGE | | | | |
| $18.9 \pm 2.9 \pm 2.2$ | 396 | ABLIKIM | 13M | BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| $22.6 \pm 3.0 \pm 2.4$ | 535 | ABLIKIM | 13M | BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{104}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 6.1 ± 1.2 OUR AVERAGE | | | | |
| 6.39 ± 1.50 ± 0.78 | 128 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| 5.86 ± 1.61 ± 0.83 | 143 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

 $\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 16.8 ± 2.5 ± 1.6 | 356 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

 $\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$ Γ_{106}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 5.82 ± 2.08 ± 0.72 | 116 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

 $\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{107}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 1.60 ± 0.27 ± 0.24 | 109 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

 $\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 1.09 ± 0.20 ± 0.16 | 82 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

 $\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{109}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 0.302 ± 0.098 ± 0.027 | 22 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

 $\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{110}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 0.125 ± 0.070 ± 0.013 | 10 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{111}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|-------------------------------------|
| 3.5 ± 2.0 OUR AVERAGE | | | | Error includes scale factor of 2.8. |

| | | | | |
|--------------------|-----|---------|----------|---|
| 5.45 ± 0.42 ± 0.87 | 671 | ABLIKIM | 05H BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$ |
|--------------------|-----|---------|----------|---|

| | | | | |
|-----------|--|------------------------|---------|-----------|
| 1.5 ± 1.0 | | ¹ TANENBAUM | 78 MRK1 | $e^+ e^-$ |
|-----------|--|------------------------|---------|-----------|

¹ Assuming entirely strong decay.

| $\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{112}/Γ |
|---|-------|-------------|------|---------|---|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $7.3 \pm 0.4 \pm 0.6$ | 434.9 | BRIERE | 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-\pi^0$ |

| $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ | | | | | Γ_{113}/Γ |
|--|-----|------|------------------------|------|-------------------------------|
| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| $7.48 \pm 0.23 \pm 0.39$ | | 1.3k | ¹ METREVELI | 12 | $\psi(2S) \rightarrow K^+K^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|------------------------|-----|------|-----------------------------------|
| $6.2 \pm 1.5 \pm 0.2$ | 66 | ^{2,3} LEES | 15J | BABR | $e^+e^- \rightarrow K^+K^-\gamma$ |
| $8.3 \pm 1.5 \pm 0.2$ | 66 | ^{3,4} LEES | 15J | BABR | $e^+e^- \rightarrow K^+K^-\gamma$ |
| $6.3 \pm 0.6 \pm 0.3$ | | ⁵ DOBBS | 06A | CLEO | e^+e^- |
| 10 ± 7 | | ⁵ BRANDELIK | 79C | DASP | e^+e^- |
| < 5 | 90 | FELDMAN | 77 | MRK1 | e^+e^- |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+e^-) = (2.37 \pm 0.04) \text{ keV}$.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant K^+K^- production not taken into account.

| $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ | | | | | Γ_{114}/Γ |
|---|------|-------------|------|---------|-----------------------|
| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT | |

5.34 ± 0.33 OUR AVERAGE

| | | | | |
|--------------------------|--------------|------------------------|-----|---|
| $5.28 \pm 0.25 \pm 0.34$ | 478 ± 23 | ¹ METREVELI | 12 | $\psi(2S) \rightarrow K_S^0 K_L^0$ |
| $5.8 \pm 0.8 \pm 0.4$ | | DOBBS | 06A | CLEO e^+e^- |
| $5.24 \pm 0.47 \pm 0.48$ | 156 ± 14 | ² BAI | 04B | BES2 $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$ |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

| $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{115}/Γ |
|---|------|-------------|------|---------|-----------------------|
| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT | |


2.01 ± 0.17 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.

| | | | | | |
|---------------------------------|--------------|----------------------|-----|------|-------------------------------------|
| $2.14 \pm 0.03^{+0.12}_{-0.11}$ | 7k | ¹ ABLIKIM | 12H | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.81 \pm 0.18 \pm 0.19$ | 260 ± 19 | ² ABLIKIM | 05J | BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.88^{+0.16}_{-0.15} \pm 0.28$ | 194 | ADAM | 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 0.85 ± 0.46 | 4 | FRANKLIN | 83 | MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

WEIGHTED AVERAGE
 2.01 ± 0.17 (Error scaled by 1.7)



$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ **Γ_{116} / Γ**

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|----------|--|
| 1.94 ± 0.25 $^{+1.15}$ $_{-0.34}$ | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$ |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ **Γ_{117} / Γ**

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|----------------------|----------|---|
| 0.32 ± 0.12 OUR AVERAGE | | | | | Error includes scale factor of 1.8. |
| $0.51 \pm 0.07 \pm 0.11$ | | | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$ |
| 0.24 $^{+0.08}$ $_{-0.07} \pm 0.02$ | | 22 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|---|---------------------|---------|-----------|
| <0.83 | 90 | 1 | FRANKLIN | 83 MRK2 | $e^+ e^-$ |
| <10 | 90 | | BARTEL | 76 CNTR | $e^+ e^-$ |
| <10 | 90 | | ² ABRAMS | 75 MRK1 | $e^+ e^-$ |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

² Final state $\rho^0 \pi^0$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{118}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|---------------------------|------|-----------------------------------|
| 0.78 ± 0.26 OUR AVERAGE | | | | | |
| $0.76 \pm 0.25 \pm 0.06$ | | 30 | ¹ METREVELI 12 | | $\psi(2S) \rightarrow \pi^+\pi^-$ |
| 8 ± 5 | | | BRANDELIK 79C | DASP | e^+e^- |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <2.1 | 90 | | DOBBS 06A | CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| <5 | 90 | | FELDMAN 77 | MRK1 | e^+e^- |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{119}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|----------|
| <3.1 | 90 | ¹ BAI 99C | BES | e^+e^- |

¹ Assuming $B(K_1(1400) \rightarrow K^*\pi) = 0.94 \pm 0.06$

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{120}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|-------------|------|-------------------------------|
| $7.12 \pm 0.62 \pm_{-0.61}^{+1.13}$ | 251 ± 22 | ABLIKIM 12L | BES3 | $e^+e^- \rightarrow \psi(2S)$ |

 $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{121}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|-------------------------------------|
| $4.07 \pm 0.16 \pm 0.26$ | | 0.9k | ABLIKIM 12L | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <8.9 | 90 | 1 | FRANKLIN 83 | MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |

 $\Gamma(K^+K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{124}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|---------------|-------------|------|-------------------------------|
| 2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.2. | | | | | |
| $3.18 \pm 0.30 \pm_{-0.31}^{+0.26}$ | | 0.2k | ABLIKIM 12L | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| $2.9 \pm_{-1.7}^{+1.3} \pm 0.4$ | | 9.6 ± 4.2 | ABLIKIM 05I | BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $1.3 \pm_{-0.7}^{+1.0} \pm 0.3$ | | 7 | ADAM 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|--|-------------|------|-------------------------------------|
| <5.4 | 90 | | FRANKLIN 83 | MRK2 | $e^+e^- \rightarrow \text{hadrons}$ |
|------|----|--|-------------|------|-------------------------------------|

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{125}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|----------------|-------------|------|-------------------------------|
| 10.9 ± 2.0 OUR AVERAGE | | | | |
| $13.3 \pm_{-2.8}^{+2.4} \pm 1.7$ | 65.6 ± 9.0 | ABLIKIM 05I | BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| $9.2 \pm_{-2.2}^{+2.7} \pm 0.9$ | 25 | ADAM 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(K^+ K^*(892)^- + c.c.) / \Gamma(K^*(892)^0 \bar{K}^0 + c.c.)$ $\Gamma_{124} / \Gamma_{125}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|-------------------------------------|
| 0.16 ± 0.06 OUR AVERAGE | | | |
| 0.22 ^{+0.10} _{-0.14} | ABLIKIM | 05I | BES2 $e^+ e^- \rightarrow \psi(2S)$ |
| 0.14 ^{+0.08} _{-0.06} | ADAM | 05 | CLEO $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\phi \pi^+ \pi^-) / \Gamma_{total}$ Γ_{126} / Γ


| VALUE (units 10 ⁻⁴) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|-----------------------|------|---|
| 1.18 ± 0.26 OUR AVERAGE | | | | Error includes scale factor of 1.5. See the ideogram below. |
| 2.3 ± 0.8 ± 0.1 | 19 ± 6 | LEES | 12F | BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 0.9 ± 0.2 ± 0.1 | 47.6 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| 1.5 ± 0.2 ± 0.2 | 51.5 ± 8.3 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2.45 ± 0.96 ± 0.04 | 10 ± 4 | ^{2,3} AUBERT | 07AK | BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

WEIGHTED AVERAGE
1.18 ± 0.26 (Error scaled by 1.5)



$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{127} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|-----------------------|-----------|---|
| 0.75 ± 0.33 OUR AVERAGE | | | | Error includes scale factor of 1.6. |
| 1.5 ± 0.5 ± 0.1 | 12 ± 4 | LEES | 12F BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 0.6 ± 0.2 ± 0.1 | 18.4 ± 6.4 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.46 ± 0.71 ± 0.02 | 6 ± 3 | ^{2,3} AUBERT | 07AK BABR | 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

 $\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$ Γ_{128} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|----------|---|
| 0.63 ± 0.13 OUR AVERAGE | | | | |
| 0.9 ± 0.4 ± 0.1 | 13 | LEES | 12F BABR | 10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$ |
| 0.6 ± 0.1 ± 0.1 | 59.2 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |

 $\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$ Γ_{129} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------------|------------------|---------|---|
| 0.70 ± 0.16 OUR AVERAGE | | | | |
| 0.8 ± 0.2 ± 0.1 | 36.8 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |
| 0.6 ± 0.2 ± 0.1 | 16.1 ± 5.0 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow 2(K^+ K^-)$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(2(K^+ K^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{130} / Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 1.1 ± 0.2 ± 0.2 | 44.7 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$ |

 $\Gamma(\phi \eta) / \Gamma_{\text{total}}$ Γ_{131} / Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|----------|--------------------------------|
| 3.10 ± 0.31 OUR AVERAGE | | | | |
| 3.14 ± 0.23 ± 0.23 | 0.2k | ABLIKIM | 12L BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| 2.0 $^{+1.5}_{-1.1}$ ± 0.4 | 6 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
| 3.3 ± 1.1 ± 0.5 | 17 | ABLIKIM | 04k BES | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\eta \phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{132} / Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|-------------|----------|--|
| < 2.2 × 10⁻⁶ | 90 | ABLIKIM | 19i BES3 | $e^+ e^- \rightarrow \eta \phi f_0(980)$ |

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{133}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|----------------------|-----------|-------------------------------|
| 1.54 ± 0.20 OUR AVERAGE | | | | |
| 1.51 ± 0.16 ± 0.12 | 201 | ABLIKIM | 19BA BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| 3.1 ± 1.4 ± 0.7 | 8 | ¹ ABLIKIM | 04K BES | $e^+e^- \rightarrow \psi(2S)$ |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

 $\Gamma(\omega\eta')/\Gamma_{\text{total}}$ Γ_{136}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------------|---------|-------------------------------|
| 3.2^{+2.4}_{-2.0} ± 0.7 | 4 | ¹ ABLIKIM | 04K BES | $e^+e^- \rightarrow \psi(2S)$ |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{137}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|-------------------------------|
| 2.1 ± 0.6 OUR AVERAGE | | | | |
| 2.5 ^{+1.2} _{-1.0} ± 0.2 | 14 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 1.87 ^{+0.68} _{-0.62} ± 0.28 | 14 | ABLIKIM | 04L BES | $e^+e^- \rightarrow \psi(2S)$ |

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$ Γ_{138}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|---------|-------------------------------|
| 1.87^{+1.64}_{-1.11} ± 0.33 | 2 | ABLIKIM | 04L BES | $e^+e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|-----|----------------------|-----------|-------------------------------|
| 1.02 ± 0.11 ± 0.24 | 143 | ¹ ABLIKIM | 17AK BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| 0.569 ± 0.128 ± 0.236 | 80 | ² ABLIKIM | 17AK BES3 | $e^+e^- \rightarrow \psi(2S)$ |

¹ Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

² Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$ Γ_{139}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|-------------------------------------|
| 2.2 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 3.0 ^{+1.1} _{-0.9} ± 0.2 | 18 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 1.78 ^{+0.67} _{-0.62} ± 0.17 | 13 | ABLIKIM | 04L BES | $e^+e^- \rightarrow \psi(2S)$ |

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{140}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|---------|-------------------------------|
| <1.1 | 90 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <3.1 | 90 | ABLIKIM | 04K BES | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{141}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|-------------------------------|
| <0.04 | 90 | ABLIKIM 12L | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.7 | 90 | ADAM 05 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| <0.4 | 90 | ABLIKIM 04K | BES | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{142}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|-------------------------------|
| <1.0 | 90 | PEDLAR 07 | CLEO | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$ Γ_{143}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| $2.7 \pm 0.6 \pm 0.4$ | 30.1 | BRIERE 05 | CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{144}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------------------|------|-----------------------------|
| $0.81 \pm 0.11 \pm 0.14$ | 50 | ¹ ABLIKIM 08C | BES2 | $e^+e^- \rightarrow J/\psi$ |

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$ Γ_{145}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|---------|-------------|------|----------------------------------|
| $0.44 \pm 0.12 \pm 0.11$ | 20 ± 6 | BAI 04C | | | $\psi(2S) \rightarrow 2(K^+K^-)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.45 | 90 | BAI 98J | BES | | $e^+e^- \rightarrow 2(K^+K^-)$ |

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{134}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------------------|------|-------------------------------|
| $3.0 \pm 0.4 \pm 1.3$ | 234 | ¹ ABLIKIM 19BA | BES3 | $e^+e^- \rightarrow \psi(2S)$ |

¹ ABLIKIM 19BA reports $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_{135}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|--------------|------|-------------------------------|
| $8.46 \pm 1.37 \pm 0.92$ | 195 | ABLIKIM 19BA | BES3 | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0\rho K^-\bar{n} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{146}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <0.88 | 90 | BAI 04G | BES2 | e^+e^- |

$\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0\rho K^-\bar{n})/\Gamma_{\text{total}}$ Γ_{147}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|----------|
| <1.0 | 90 | BAI 04G | BES2 | e^+e^- |

$\Gamma(\Theta(1540)K_S^0\bar{p} \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$ Γ_{148}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------------|
| <0.70 | 90 | BAI | 04G | BES2 e^+e^- |

$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$ Γ_{149}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------------|
| <2.6 | 90 | BAI | 04G | BES2 e^+e^- |

$\Gamma(\bar{\Theta}(1540)K_S^0p \rightarrow K_S^0pK^-\bar{n})/\Gamma_{\text{total}}$ Γ_{150}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------------|
| <0.60 | 90 | BAI | 04G | BES2 e^+e^- |

$\Gamma(K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{151}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------------|------|--------------|
| <0.046 | ¹ BAI | 04D | BES e^+e^- |

¹Forbidden by CP.

$\Gamma(\Lambda_c^+ \bar{p}e^+e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{152}/Γ

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|-----|------|-------------|------|------------------------------------|
| <1.7 × 10 ⁻⁶ | 90 | 450M | ABLIKIM | 18Q | BES3 $e^+e^- \rightarrow \psi(2S)$ |

————— RADIATIVE DECAYS —————

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{153}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-----------------------|------|------------------------------------|
| 9.79 ± 0.20 OUR FIT | | | | |
| 9.33 ± 0.26 OUR AVERAGE | | | | |
| 9.389 ± 0.014 ± 0.332 | 4.7M | ABLIKIM | 17U | BES3 $e^+e^- \rightarrow \gamma X$ |
| 9.22 ± 0.11 ± 0.46 | 72k | ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |
| 9.9 ± 0.5 ± 0.8 | | ¹ GAISER | 86 | CBAL $e^+e^- \rightarrow \gamma X$ |
| 7.2 ± 2.3 | | ¹ BIDDICK | 77 | CNTR $e^+e^- \rightarrow \gamma X$ |
| 7.5 ± 2.6 | | ¹ WHITAKER | 76 | MRK1 e^+e^- |

¹Angular distribution $(1+\cos^2\theta)$ assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{154}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|----------------------|------|------------------------------------|
| 9.75 ± 0.24 OUR FIT | | | | |
| 9.54 ± 0.29 OUR AVERAGE | | | | |
| 9.905 ± 0.011 ± 0.353 | 5.0M | ABLIKIM | 17U | BES3 $e^+e^- \rightarrow \gamma X$ |
| 9.07 ± 0.11 ± 0.54 | 76k | ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |
| 9.0 ± 0.5 ± 0.7 | | ¹ GAISER | 86 | CBAL $e^+e^- \rightarrow \gamma X$ |
| 7.1 ± 1.9 | | ² BIDDICK | 77 | CNTR $e^+e^- \rightarrow \gamma X$ |

¹Angular distribution $(1-0.189\cos^2\theta)$ assumed.

²Valid for isotropic distribution of the photon.


$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{155}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|--|
| 9.52 ± 0.20 | | | | OUR FIT |
| 9.42 ± 0.31 | | | | OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below. |
| 9.621 ± 0.013 ± 0.272 | 4.2M | ABLIKIM | 17U | BES3 $e^+e^- \rightarrow \gamma X$ |
| 9.33 ± 0.14 ± 0.61 | 79k | ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |
| 8.0 ± 0.5 ± 0.7 | | ¹ GAISER | 86 | CBAL $e^+e^- \rightarrow \gamma X$ |
| 7.0 ± 2.0 | | ² BIDDICK | 77 | CNTR $e^+e^- \rightarrow \gamma X$ |

¹ Angular distribution $(1 - 0.052 \cos^2\theta)$ assumed.

² Valid for isotropic distribution of the photon.

WEIGHTED AVERAGE
9.42 ± 0.31 (Error scaled by 1.3)



$[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}}$ ($\Gamma_{153} + \Gamma_{154} + \Gamma_{155}$)/ Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|--------------------|------|------------------------------------|
| 27.6 ± 0.3 ± 2.0 | ¹ ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{153}/\Gamma_{154}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------|--------------------|------|------------------------------------|
| 1.02 ± 0.01 ± 0.07 | ¹ ATHAR | 04 | CLEO $e^+e^- \rightarrow \gamma X$ |

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

$\Gamma_{155}/\Gamma_{154}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.03 \pm 0.02 \pm 0.03$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$

$\Gamma_{153}/\Gamma_{155}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.99 \pm 0.02 \pm 0.08$ ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\eta_c(1S))/\Gamma_{total}$

Γ_{156}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

0.34 ± 0.05 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

$0.432 \pm 0.016 \pm 0.060$ MITCHELL 09 CLEO $e^+e^- \rightarrow \gamma X$


$0.32 \pm 0.04 \pm 0.06$ 2.5k ¹ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

0.28 ± 0.06 ² GAISER 86 CBAL $e^+e^- \rightarrow \gamma X$

¹ ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

² GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

WEIGHTED AVERAGE
 0.34 ± 0.05 (Error scaled by 1.3)



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{157}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|-----------------------------|------|--|
| $7 \pm 2 \pm 4$ | | ¹ ABLIKIM 12G | BES3 | $\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$ |
| < 8 | 90 | ² CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K \bar{K} \pi$ |
| < 20 | 90 | ATHAR 04 | CLEO | $e^+ e^- \rightarrow \gamma X$ |
| 20–130 | 95 | EDWARDS 82c | CBAL | $e^+ e^- \rightarrow \gamma X$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K \bar{K} \pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K \bar{K} \pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------------------------------|------|------------------------------------|
| 1.04 ± 0.22 OUR AVERAGE | | | Error includes scale factor of 1.4. | | |
| $0.95 \pm 0.16 \pm 0.05$ | | 423 | ABLIKIM 17X | BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |
| $1.58 \pm 0.40 \pm 0.13$ | | 37 | ABLIKIM 10F | BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------|----|--------------------------|------|---------------------------------|
| < 5 | 90 | PEDLAR 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 5400 | 95 | ¹ LIBERMAN 75 | SPEC | $e^+ e^-$ |
| $< 1 \times 10^4$ | 90 | WIJK 75 | DASP | $e^+ e^-$ |

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$.

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{159}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------|--------------------------|------|---|
| 1.24 ± 0.04 OUR AVERAGE | | | | | |
| $1.251 \pm 0.022 \pm 0.062$ | | 56K | ABLIKIM 17X | BES3 | $\psi(2S) \rightarrow \gamma\pi^+ \pi^- \eta, \gamma\pi^0 \pi^0 \eta$ |
| $1.26 \pm 0.03 \pm 0.08$ | | 2226 | ¹ ABLIKIM 10F | BES3 | $\psi(2S) \rightarrow 3\gamma\pi^+ \pi^-, 2\gamma\pi^+ \pi^-$ |
| $1.19 \pm 0.08 \pm 0.03$ | | | PEDLAR 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| $1.24 \pm 0.27 \pm 0.15$ | | 23 | ABLIKIM 06R | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| $1.54 \pm 0.31 \pm 0.20$ | | ~ 43 | BAI 98F | BES | $\psi(2S) \rightarrow \pi^+ \pi^- 2\gamma, \pi^+ \pi^- 3\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|-----------------------------|------|-----------|
| < 60 | 90 | ² BRAUNSCH... 77 | DASP | $e^+ e^-$ |
| < 11 | 90 | ³ BARTEL 76 | CNTR | $e^+ e^-$ |

¹ Combining the results from $\eta' \rightarrow \pi^+ \pi^- \eta$ and $\eta' \rightarrow \pi^+ \pi^- \gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}} \qquad \Gamma_{160}/\Gamma$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

2.73^{+0.29}_{-0.25} OUR AVERAGE Error includes scale factor of 1.8.

| | | | | |
|---|--------------|-----------|---------|---|
| 2.84 ± 0.15 ^{+0.03} _{-0.10} | 1.9k | 1,2 DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |
| 2.12 ± 0.19 ± 0.32 | | 3,4 BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi \pi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.08 ± 0.19 ± 0.33 | 200.6 ± 18.8 | 3 BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ |
| 2.90 ± 1.08 ± 1.07 | 29.9 ± 11.1 | 3 BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+ \pi^-$ and $\pi^0 \pi^0$ decay modes.

$$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}} \qquad \Gamma_{161}/\Gamma$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|--------------------------|------|-------------|---------|
|--------------------------|------|-------------|---------|

3.1 ± 1.0 ± 1.4 175 ¹ DOBBS 15 $\psi(2S) \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}} \qquad \Gamma_{162}/\Gamma$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|--------------------------|------|-------------|---------|
|--------------------------|------|-------------|---------|

9.3 ± 1.8 ± 0.6 274 ^{1,2} DOBBS 15 $\psi(2S) \rightarrow \gamma \pi \pi$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi \pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi \pi) = (34.5 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}} \qquad \Gamma_{163}/\Gamma$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|--------------------------|------|-------------|---------|
|--------------------------|------|-------------|---------|

3.3 ± 0.8 ± 0.1 136 ^{1,2} DOBBS 15 $\psi(2S) \rightarrow \gamma K \bar{K}$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}} \qquad \Gamma_{165}/\Gamma$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

3.5 ± 0.6 OUR AVERAGE

| | | | | |
|--------------------|------------|---------|---------|---|
| 3.6 ± 0.4 ± 0.5 | 290 | 1 DOBBS | 15 | $\psi(2S) \rightarrow \gamma \pi \pi$ |
| 3.01 ± 0.41 ± 1.24 | 35.6 ± 4.8 | 2 BAI | 03C BES | $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{166}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|-----------|--------------------|---------|---|
| 6.6 ± 0.7 | | | | | OUR AVERAGE |
| 6.7 ± 0.6 ± 0.6 | | 375 | ¹ DOBBS | 15 | $\psi(2S) \rightarrow \gamma K \bar{K}$ |
| 6.04 ± 0.90 ± 1.32 | 39.6 ± 5.9 | | ^{2,3} BAI | 03C BES | $\psi(2S) \rightarrow \gamma K^+ K^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 15.6 | 90 | 6.8 ± 3.1 | ^{2,3} BAI | 03C BES | $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$. $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{167}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | COMMENT |
|--------------------------|------|--------------------|--|
| 4.8 ± 0.5 ± 0.9 | 373 | ¹ DOBBS | 15 $\psi(2S) \rightarrow \gamma \pi \pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{168}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | COMMENT |
|--------------------------|------|--------------------|--|
| 3.2 ± 0.6 ± 0.8 | 207 | ¹ DOBBS | 15 $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{169}/Γ

| VALUE | CL% | DOCUMENT ID | COMMENT |
|-----------------------------------|-----|----------------------|--|
| < 5.8 × 10⁻⁶ | 90 | ^{1,2} DOBBS | 15 $\psi(2S) \rightarrow \gamma \pi \pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively. $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{170}/Γ

| VALUE | CL% | DOCUMENT ID | COMMENT |
|-----------------------------------|-----|----------------------|--|
| < 9.5 × 10⁻⁶ | 90 | ^{1,2} DOBBS | 15 $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively. $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{172}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|----------------------|----------|--|
| 0.92 ± 0.18 | | | | | OUR AVERAGE |
| 0.85 ± 0.18 ± 0.04 | | 382 | ¹ ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |
| 1.38 ± 0.48 ± 0.09 | | 13 | ¹ ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|--------|-----|------|--|
| < 2 | 90 | PEDLAR | 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 90 | 90 | BAI | 98F | BES | $\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$ |
| < 200 | 90 | YAMADA | 77 | DASP | $e^+ e^- \rightarrow 3\gamma$ |

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

$\Gamma(\gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{173}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|--|
| $8.71 \pm 1.25 \pm 1.64$ | 418 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{175}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|--|
| < 0.9 | 90 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3 90 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

< 1.2 90 ¹ SCHARRE 80 MRK1 $e^+ e^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

 $\Gamma(\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{176}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|--|
| $0.36 \pm 0.25 \pm 0.05$ | 10 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{177}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---|
| $< 5.0 \times 10^{-7}$ | 90 | ABLIKIM | 17AJ | BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |

 $\Gamma(\gamma\eta(1475) \rightarrow K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{179}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|------|--|
| < 1.4 | 90 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.5 90 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

 $\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{180}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-----|-------------|------|--|
| < 0.88 | 90 | ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{181}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|-------------------------------------|
| $39.6 \pm 2.8 \pm 5.0$ | 583 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{182}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|-------------------------------------|
| $37.0 \pm 6.1 \pm 7.2$ | 237 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{183}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|-------------------------------------|
| $24.0 \pm 4.5 \pm 5.0$ | 41 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{184}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|-------------------------------------|
| $25.6 \pm 3.6 \pm 3.6$ | 115 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{185}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--------------------------------|
| 19.1 ± 2.7 ± 4.3 | 132 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{186}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------------------------------|------|---|
| 3.9 ± 0.5 OUR AVERAGE | | Error includes scale factor of 2.0. | | |
| 4.18 ± 0.26 ± 0.18 | 348 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p \bar{p}$ |
| 2.9 ± 0.4 ± 0.4 | 142 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

¹From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{187}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------------|------|---|
| 1.2 ± 0.2 ± 0.1 | 111 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p \bar{p}$ |

¹From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{188}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|---------------------------|------|---|
| 0.72 ± 0.18 ± 0.03 | 73 | ¹ ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p \bar{p}$ |

¹From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{189}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------|------|---------------------------------------|
| 4.57 ± 0.36 ^{+1.77}/_{-4.26} | | ABLIKIM 12D | BES3 | $J/\psi \rightarrow \gamma p \bar{p}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|--------------|------|---|
| <1.6 | 90 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p \bar{p}$ |
| <5.4 | 90 | ABLIKIM 07D | BES | $\psi(2S) \rightarrow \gamma p \bar{p}$ |

 $\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{190}/Γ

For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------|------|---|
| <2 | 90 | ALEXANDER 10 | CLEO | $\psi(2S) \rightarrow \gamma p \bar{p}$ |

 $\Gamma(\gamma \pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}}$ Γ_{191}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--------------------------------|
| 2.8 ± 1.2 ± 0.7 | 17 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{192}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|--------------------------------|
| <22 | 90 | ABLIKIM 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{193}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|-------------------------------------|
| <17 | 90 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{194}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|-------------------------------------|
| <4 | 90 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{195}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-------------------------------------|
| $3.1 \pm 0.6^{+0.8}_{-1.0}$ | 1.1k | ABLIKIM | 120 | BES3 $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------|------|----------------------|-----|---|
| 3.2 ± 0.6 | 1.1k | ¹ ABLIKIM | 17N | BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$ |
|---------------|------|----------------------|-----|---|

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.

$\Gamma(e^+ e^- \eta')/\Gamma_{\text{total}}$ Γ_{196}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| 1.90 ± 0.26 OUR AVERAGE | | | | |
| $1.99 \pm 0.33 \pm 0.12$ | 57 | ABLIKIM | 18Z | BES3 $\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \gamma \pi^+ \pi^-$ |
| $1.79 \pm 0.38 \pm 0.11$ | 20 | ABLIKIM | 18Z | BES3 $\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \eta \pi^+ \pi^-$ |

$\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{197}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------------|------|---|
| $10.6 \pm 2.4 \pm 0.4$ | 48 | ¹ ABLIKIM | 17i | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17i reports $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{198}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|---|
| $8.5 \pm 0.6 \pm 0.2$ | 873 | ¹ ABLIKIM | 17i | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17i reports $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(e^+ e^- \chi_{c2}(1P)) / \Gamma_{\text{total}} \qquad \Gamma_{199} / \Gamma$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|----------|--|
| $7.0 \pm 0.7 \pm 0.2$ | 227 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17I reports $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c2}(1P)) / \Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(e^+ e^- \chi_{c0}(1P)) / \Gamma(\gamma \chi_{c0}(1P)) \qquad \Gamma_{197} / \Gamma_{153}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|----------|--|
| $9.4 \pm 1.9 \pm 0.6$ | 48 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$$\Gamma(e^+ e^- \chi_{c1}(1P)) / \Gamma(\gamma \chi_{c1}(1P)) \qquad \Gamma_{198} / \Gamma_{154}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|----------|--|
| $8.3 \pm 0.3 \pm 0.4$ | 873 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$$\Gamma(e^+ e^- \chi_{c2}(1P)) / \Gamma(\gamma \chi_{c2}(1P)) \qquad \Gamma_{199} / \Gamma_{155}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|----------|--|
| $6.6 \pm 0.5 \pm 0.4$ | 227 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

WEAK DECAYS

$$\Gamma(D^0 e^+ e^- + \text{c.c.}) / \Gamma_{\text{total}} \qquad \Gamma_{200} / \Gamma$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|-----------|--------------------------------|
| $< 1.4 \times 10^{-7}$ | 90 | ¹ ABLIKIM | 17AF BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

OTHER DECAYS

$$\Gamma(\text{invisible}) / \Gamma(e^+ e^-) \qquad \Gamma_{201} / \Gamma_6$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----|-------------|----------|----------------------------------|
| < 2.0 | 90 | LEES | 13I BABR | $B \rightarrow K^{(*)} \psi(2S)$ |

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$$\psi(2S) \rightarrow \gamma \chi_{cJ}(1P) \text{ and } \chi_{cJ} \rightarrow \gamma J/\psi(1S)$$

 $a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| 63 ± 7 OUR AVERAGE | | | | |
| 61.7 ± 8.3 | 253k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |
| 67 ⁺¹⁹ ₋₁₃ | 59k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |

¹ Statistical and systematic errors combined.² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09. **$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio**

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| 60 ± 31 OUR AVERAGE | | | | |
| 74 ± 40 | 253k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |
| 37 ⁺⁵³ ₋₄₇ | 59k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$ |

¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09. **$\psi(2S)$ REFERENCES**

| | | | | |
|---------|------|----------------|----------------------------|------------------|
| ABLIKIM | 19AO | PR D99 112010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AT | PR D100 051101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AU | PR D100 052010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19BA | PR D100 092003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19I | PR D99 012014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19N | PR D99 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18Q | PR D97 091102 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18T | PR D98 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18Z | PL B783 452 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ANASHIN | 18 | PL B781 174 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 18E | PR D98 112015 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 17AF | PR D96 111101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AJ | PR D96 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AK | PR D96 112012 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17E | PL B770 217 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17I | PRL 118 221802 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17L | PR D95 052003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17N | PR D95 072004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17U | PR D96 032001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17X | PR D96 052003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DOBBS | 17 | PR D96 092004 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| LEES | 17A | PR D95 052001 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 16Y | JHEP 1605 132 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 16L | PR D93 072003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15I | PR D91 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15V | PL B749 414 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ANASHIN | 15 | PL B749 50 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| DOBBS | 15 | PR D91 052006 | S. Dobbs <i>et al.</i> | (NWES) |
| LEES | 15J | PR D92 072008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 14G | PR D89 112006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |

| | | | | |
|---------------|------|-------------------------------|-----------------------------------|------------------------|
| DOBBS | 14 | PL B739 90 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| ABLIKIM | 13A | PRL 110 022001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13D | PR D87 012007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13F | PR D87 052007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13M | PR D87 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13R | PR D88 032007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13S | PR D88 032010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13W | PR D88 112007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 13I | PR D87 112005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13O | PR D87 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Q | PR D88 032013 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Y | PR D88 072009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 12H | EPJ C72 1972 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 12D | PRL 108 112003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12G | PRL 109 042003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12H | PL B710 594 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12L | PR D86 072011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12M | PR D86 092008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12O | PRL 109 172002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12Q | CP C36 1040 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ANASHIN | 12 | PL B711 280 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 12E | PR D85 112009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 12F | PR D86 012008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| METREVELI | 12 | PR D85 092007 | Z. Metreveli <i>et al.</i> | (NWES, FLOR, WAYN+) |
| GE | 11 | PR D84 032008 | J.Y. Ge <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 10B | PRL 104 132002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 10F | PRL 105 261801 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ALEXANDER | 10 | PR D82 092002 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| CRONIN-HEN... | 10 | PR D81 052002 | D. Cronin-Hennessey <i>et al.</i> | (CLEO Collab.) |
| ADAMS | 09 | PR D80 051106 | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 09 | PR D80 112003 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| LIBBY | 09 | PR D80 072002 | J. Libby <i>et al.</i> | (CLEO Collab.) |
| MITCHELL | 09 | PRL 102 011801 | R.E. Mitchell <i>et al.</i> | (CLEO Collab.) |
| PEDLAR | 09 | PR D79 111101 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 08B | PL B659 74 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 08C | PL B659 789 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| DOBBS | 08A | PRL 101 182003 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| MENDEZ | 08 | PR D78 011102 | H. Mendez <i>et al.</i> | (CLEO Collab.) |
| PDG | 08 | PL B667 1 | C. Amsler <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 07C | PL B648 149 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 07D | PRL 99 011802 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ABLIKIM | 07H | PR D76 092003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ANASHIN | 07 | JETPL 85 347 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| | | Translated from ZETFP 85 429. | | |
| ANDREOTTI | 07 | PL B654 74 | M. Andreotti <i>et al.</i> | (Femilab E835 Collab.) |
| AUBERT | 07AK | PR D76 012008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AU | PR D76 092005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| Also | | PR D77 119902E (errat.) | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BD | PR D76 092006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| PDG | 07 | Unofficial 2007 WWW edition | | (PDG Collab.) |
| PEDLAR | 07 | PR D75 011102 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 06G | PR D73 052004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06I | PR D74 012004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06L | PRL 97 121801 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06R | PR D74 072001 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06W | PR D74 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 06 | PRL 96 082004 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 06B | PR D73 012005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06D | PR D73 052003 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT_BE | 06D | PR D74 091103 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| DOBBS | 06A | PR D74 011105 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 05E | PR D71 072006 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05H | PR D72 012002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05I | PL B614 37 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05J | PL B619 247 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05O | PL B630 21 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 05 | PRL 94 012005 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ADAM | 05A | PRL 94 232002 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ANDREOTTI | 05 | PR D71 032006 | M. Andreotti <i>et al.</i> | (FNAL E835 Collab.) |
| AUBERT | 05D | PR D71 052001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BRIERE | 05 | PRL 95 062001 | R.A. Briere <i>et al.</i> | (CLEO Collab.) |

| | | | | |
|-------------|-----|------------------------------|-------------------------------|-------------------------|
| PEDLAR | 05 | PR D72 051108 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ROSNER | 05 | PRL 95 102003 | J.L. Rosner <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 04B | PR D70 012003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04K | PR D70 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04L | PR D70 112007 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ATHAR | 04 | PR D70 112002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| BAI | 04B | PRL 92 052001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04C | PR D69 072001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04D | PL B589 7 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04G | PR D70 012004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04I | PR D70 012006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 04 | PL B592 1 | S. Eidelman <i>et al.</i> | (PDG Collab.) |
| SETH | 04 | PR D69 097503 | K.K. Seth | |
| AULCHENKO | 03 | PL B573 63 | V.M. Aulchenko <i>et al.</i> | (KEDR Collab.) |
| BAI | 03B | PR D67 052002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 03C | PR D67 032004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AUBERT | 02B | PR D65 031101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BAI | 02 | PR D65 052004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02B | PL B550 24 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02C | PRL 88 101802 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 02 | PR D66 010001 | K. Hagiwara <i>et al.</i> | (PDG Collab.) |
| BAI | 01 | PR D63 032002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AMBROGIANI | 00A | PR D62 032004 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ARTAMONOV | 00 | PL B474 427 | A.S. Artamonov <i>et al.</i> | |
| BAI | 00 | PRL 84 594 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 99C | PRL 83 1918 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98E | PR D57 3854 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98F | PR D58 097101 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98J | PRL 81 5080 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ARMSTRONG | 97 | PR D55 1153 | T.A. Armstrong <i>et al.</i> | (E760 Collab.) |
| GRIBUSHIN | 96 | PR D53 4723 | A. Gribushin <i>et al.</i> | (E672 and E706 Collab.) |
| ARMSTRONG | 93B | PR D47 772 | T.A. Armstrong <i>et al.</i> | (FNAL E760 Collab.) |
| ALEXANDER | 89 | NP B320 45 | J.P. Alexander <i>et al.</i> | (LBL, MICH, SLAC) |
| COHEN | 87 | RMP 59 1121 | E.R. Cohen, B.N. Taylor | (RISC, NBS) |
| GAISER | 86 | PR D34 711 | J. Gaiser <i>et al.</i> | (Crystal Ball Collab.) |
| KURAEV | 85 | SJNP 41 466 | E.A. Kuraev, V.S. Fadin | (NOVO) |
| | | Translated from YAF 41 733. | | |
| FRANKLIN | 83 | PRL 51 963 | M.E.B. Franklin <i>et al.</i> | (LBL, SLAC) |
| EDWARDS | 82C | PRL 48 70 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| LEMOIGNE | 82 | PL 113B 509 | Y. Lemoigne <i>et al.</i> | (SACL, LOIC, SHMP+) |
| HIMEL | 80 | PRL 44 920 | T. Himel <i>et al.</i> | (LBL, SLAC) |
| OREGLIA | 80 | PRL 45 959 | M.J. Oreglia <i>et al.</i> | (SLAC, CIT, HARV+) |
| SCHARRE | 80 | PL 97B 329 | D.L. Scharre <i>et al.</i> | (SLAC, LBL) |
| ZHOLENTZ | 80 | PL 96B 214 | A.A. Zholents <i>et al.</i> | (NOVO) |
| Also | | SJNP 34 814 | A.A. Zholents <i>et al.</i> | (NOVO) |
| | | Translated from YAF 34 1471. | | |
| BRANDELIK | 79B | NP B160 426 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BRANDELIK | 79C | ZPHY C1 233 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BARTEL | 78B | PL 79B 492 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 78 | PR D17 1731 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) |
| BIDDICK | 77 | PRL 38 1324 | C.J. Biddick <i>et al.</i> | (UCSD, UMD, PAVI+) |
| BRAUNSCH... | 77 | PL 67B 249 | W. Braunschweig <i>et al.</i> | (DASP Collab.) |
| BURMESTER | 77 | PL 66B 395 | J. Burmester <i>et al.</i> | (DESY, HAMB, SIEG+) |
| FELDMAN | 77 | PRPL 33C 285 | G.J. Feldman, M.L. Perl | (LBL, SLAC) |
| YAMADA | 77 | Hamburg Conf. 69 | S. Yamada | (DASP Collab.) |
| BARTEL | 76 | PL 64B 483 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 76 | PRL 36 402 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) IG |
| WHITAKER | 76 | PRL 37 1596 | J.S. Whitaker <i>et al.</i> | (SLAC, LBL) |
| ABRAMS | 75 | Stanford Symp. 25 | G.S. Abrams | (LBL) |
| ABRAMS | 75B | PRL 34 1181 | G.S. Abrams <i>et al.</i> | (LBL, SLAC) |
| BOYARSKI | 75C | Palermo Conf. 54 | A.M. Boyarski <i>et al.</i> | (SLAC, LBL) |
| HILGER | 75 | PRL 35 625 | E. Hilger <i>et al.</i> | (STAN, PENN) |
| LIBERMAN | 75 | Stanford Symp. 55 | A.D. Liberman | (STAN) |
| LUTH | 75 | PRL 35 1124 | V. Luth <i>et al.</i> | (SLAC, LBL) JPC |
| WIIK | 75 | Stanford Symp. 69 | B.H. Wiik | (DESY) |