

$B_J(5840)$

$I(J^P) = \frac{1}{2}(??)$
 I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

$B_J(5840)$ MASS

$B_J(5840)^+$ MASS

OUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.

VALUE (MeV)	DOCUMENT ID
-------------	-------------

5851±19 OUR FIT

$m_{B_J(5840)^+} - m_{B^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

571±19 OUR FIT

571±13±14 7k ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

595±26±14 7k ² AAIJ 15AB LHCb $p p$ at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$m_{B_J(5840)^+} - m_{B^{*0}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

565±15±14 7k ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5840)^0$ MASS

OUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.

VALUE (MeV)	DOCUMENT ID
-------------	-------------

5863±9 OUR FIT

$m_{B_J(5840)^0} - m_{B^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
584± 9 OUR FIT				
584± 5±7	12k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				

610±22±7 12k ² AAIJ 15AB LHCb $p p$ at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

 $m_{B_J(5840)^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
584±5±7	12k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
1 AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV				

which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

 $B_J(5840)$ WIDTH **$B_J(5840)^+$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
224±24±80	7k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
215±27±80	7k	² AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
229±27±80	7k	³ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

 $B_J(5840)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
127±17±34	12k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
107±20±34	12k	² AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
119±17±34	12k	³ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

- ² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.
³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.
-

$B_J(5840)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^* \pi$	seen
$\Gamma_2 \quad B \pi$	possibly seen

$B_J(5840)$ BRANCHING RATIOS

$\Gamma(B^* \pi)/\Gamma_{\text{total}}$	Γ_1/Γ				
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	7k	AAIJ	15AB LHCb	\pm	$p p$ at 7, 8 TeV
seen	12k	AAIJ	15AB LHCb	0	$p p$ at 7, 8 TeV

$\Gamma(B \pi)/\Gamma_{\text{total}}$	Γ_2/Γ				
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
possibly seen	7k	¹ AAIJ	15AB LHCb	\pm	$p p$ at 7, 8 TeV
possibly seen		¹ AAIJ	15AB LHCb	0	$p p$ at 7, 8 TeV

¹ A $B \pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^* \pi$ is allowed.

$B_J(5840)$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------
