

# $D_0^*(2300)$

$$I(J^P) = \frac{1}{2}(0^+)$$


was  $D_0^*(2400)$

There is a strong evidence that recent data on  $B \rightarrow D\pi\pi$  (AAIJ 15Y, AAIJ 16AH) and  $B \rightarrow D\pi K$  (AAIJ 14BH, AAIJ 15V, AAIJ 15X) call for two poles in the scalar  $I = 1/2 \pi D$  amplitude in this mass range. The data are consistent with a lower pole at  $(2105^{+6}_{-8}) - i(102^{+10}_{-11})$  MeV and a higher pole at  $(2451^{+35}_{-26}) - i(134^{+7}_{-8})$  MeV (DU 18A, DU 19, DU 21). For details see review on "Heavy Non- $q\bar{q}$  Mesons."

## $D_0^*(2300)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>2343±10 OUR AVERAGE</b>	Error	includes scale factor	of 1.5.	See the	ideogram below.
2360±15±30		1 AAIJ	15X LHCB	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2349± 6± 4		2 AAIJ	15Y LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2297± 8±20	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2308±17±32		ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2354± 7±11		3 AAIJ	15Y LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2403±14±35	18.8k	4 LINK	04A FOCS	+	$\gamma A$
2407±21±35	9.8k	4 LINK	04A FOCS	0	$\gamma A$

WEIGHTED AVERAGE  
2343±10 (Error scaled by 1.5)



<sup>1</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$  S-wave and the  $D\pi$  S- and P-waves.

- <sup>2</sup> Modeling the  $\pi^+\pi^-$  S-wave with the Isobar formalism.  
<sup>3</sup> Modeling the  $\pi^+\pi^-$  S-wave with the K-matrix formalism.  
<sup>4</sup> Possibly the feed-down from another state.

### $D_0^*(2300)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>229±16 OUR AVERAGE</b>					
255±26±51		<sup>1</sup> AAIJ	15X LHCb	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
217±13±13		<sup>2</sup> AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
273±12±48	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
276±21±63		ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
230±15±21		<sup>3</sup> AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
283±24±34	18.8k	<sup>4</sup> LINK	04A FOCS	+	$\gamma A$
240±55±59	9.8k	<sup>4</sup> LINK	04A FOCS	0	$\gamma A$

- <sup>1</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$  S-wave and the  $D\pi$  S- and P-waves.  
<sup>2</sup> Modeling the  $\pi^+\pi^-$  S-wave with the Isobar formalism.  
<sup>3</sup> Modeling the  $\pi^+\pi^-$  S-wave with the K-matrix formalism.  
<sup>4</sup> Possibly the feed-down from another state.

### $D_0^*(2300)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 D\pi^\pm$	seen

$\Gamma(D\pi^\pm)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen		AAIJ	15X LHCb	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$
seen		AAIJ	15Y LHCb	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$
seen	3.4k	AUBERT	09AB BABR	0	$D^*(2300)^0 \rightarrow D^+ \pi^-$
<b>seen</b>		ABE	04D BELL	0	$D^*(2300)^0 \rightarrow D^+ \pi^-$
<b>seen</b>	18.8k	LINK	04A FOCS	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$

### $D_0^*(2300)$ REFERENCES

DU	21	PRL 126 192001	M.-L. Du <i>et al.</i>
DU	19	PR D99 114002	M.-L. Du, F.-K. Guo, U.-G. Meissner
DU	18A	PR D98 094018	M.-L. Du <i>et al.</i>
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i> (LHCb Collab.)
AAIJ	15V	PR D91 092002	R. Aaij <i>et al.</i> (LHCb Collab.)
Also		PR D93 119901 (errat.)	R. Aaij <i>et al.</i> (LHCb Collab.)
AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i> (LHCb Collab.)
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i> (LHCb Collab.)
AAIJ	14BH	PR D90 072003	R. Aaij <i>et al.</i> (LHCb Collab.)
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i> (BABAR Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i> (BELLE Collab.)
LINK	04A	PL B586 11	J.M. Link <i>et al.</i> (FOCUS Collab.)