



$I(J^P) = 0(\frac{3}{2}^+)$ Status: ****

The unambiguous discovery in both production and decay was by BARNES 64. The quantum numbers follow from the assignment of the particle to the baryon decuplet. DEUTSCHMANN 78 and BAUBILLIER 78 rule out $J = 1/2$ and find consistency with $J = 3/2$. AUBERT,BE 06 finds from the decay angular distributions of $\Xi_c^0 \rightarrow \Omega^- K^+$ and $\Omega_c^0 \rightarrow \Omega^- K^+$ that $J = 3/2$; this depends on the spins of the Ξ_c^0 and Ω_c^0 being $J = 1/2$, their supposed values.

ABLIKIM 21E determines the Ω^- spin to be $J = 3/2$ from the decay angular distributions of the complete decay chain $\psi(3686) \rightarrow \Omega^- \bar{\Omega}^+$, with subsequent decays $\Omega^- \rightarrow K^- \Lambda$ and $\bar{\Omega}^+ \rightarrow K^+ \bar{\Lambda}$.

We have omitted some results that have been superseded by later experiments. See our earlier editions.

Ω^- MASS

The fit assumes the Ω^- and $\bar{\Omega}^+$ masses are the same, and averages them together.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1672.45 ± 0.29 OUR FIT				
1672.43 ± 0.32 OUR AVERAGE				
1673 ± 1	100	HARTOUNI	85	SPEC 80–280 GeV $K_L^0 C$
1673.0 ± 0.8	41	BAUBILLIER	78	HBC 8.25 GeV/c $K^- p$
1671.7 ± 0.6	27	HEMINGWAY	78	HBC 4.2 GeV/c $K^- p$
1673.4 ± 1.7	4	¹ DIBIANCA	75	DBC 4.9 GeV/c $K^- d$
1673.3 ± 1.0	3	PALMER	68	HBC $K^- p$ 4.6, 5 GeV/c
1671.8 ± 0.8	3	SCHULTZ	68	HBC $K^- p$ 5.5 GeV/c
1674.2 ± 1.6	5	SCOTTER	68	HBC $K^- p$ 6 GeV/c
1672.1 ± 1.0	1	² FRY	55	EMUL
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1671.43 ± 0.78	13	³ DEUTSCH...	73	HBC $K^- p$ 10 GeV/c
1671.9 ± 1.2	6	³ SPETH	69	HBC See DEUTSCHMANN 73
1673.0 ± 8.0	1	ABRAMS	64	HBC $\rightarrow \Xi^- \pi^0$
1670.6 ± 1.0	1	² FRY	55B	EMUL
1615	1	⁴ EISENBERG	54	EMUL

¹DIBIANCA 75 gives a mass for each event. We quote the average.

²The FRY 55 and FRY 55B events were identified as Ω^- by ALVAREZ 73. The masses assume decay to ΛK^- at rest. For FRY 55B, decay from an atomic orbit could Doppler shift the K^- energy and the resulting Ω^- mass by several MeV. This shift is negligible for FRY 55 because the Ω decay is approximately perpendicular to its orbital velocity, as is known because the Λ strikes the nucleus (L.Alvarez, private communication 1973). We have calculated the error assuming that the orbital n is 4 or larger.

³Excluded from the average; the Ω^- lifetimes measured by the experiments differ significantly from other measurements.

⁴ The EISENBERG 54 mass was calculated for decay in flight. ALVAREZ 73 has shown that the Ω interacted with an Ag nucleus to give $K^- \Xi \text{Ag}$.

$\bar{\Omega}^+$ MASS

The fit assumes the Ω^- and $\bar{\Omega}^+$ masses are the same, and averages them together.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1672.45 ± 0.29 OUR FIT				
1672.5 ± 0.7 OUR AVERAGE				
1672 ± 1	72	HARTOUNI	85	SPEC 80–280 GeV $K_L^0 C$
1673.1 ± 1.0	1	FIRESTONE	71B	HBC 12 GeV/c $K^+ d$

$$(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\Omega^-}$$

A test of CPT invariance.

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
-1.44 ± 7.98		CHAN	98	E756 p Be, 800 GeV

Ω^- MEAN LIFE

Measurements with an error $> 0.1 \times 10^{-10}$ s have been omitted. The fit assumes the Ω^- and $\bar{\Omega}^+$ mean lives are the same, and averages them together.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.821 ± 0.011 OUR FIT				
0.821 ± 0.011 OUR AVERAGE				
0.817 $\pm 0.013 \pm 0.018$	6934	CHAN	98	E756 p Be, 800 GeV
0.811 ± 0.037	1096	LUK	88	SPEC p Be 400 GeV
0.823 ± 0.013	12k	BOURQUIN	84	SPEC SPS hyperon beam
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.822 ± 0.028	2437	BOURQUIN	79B	SPEC See BOURQUIN 84

$\bar{\Omega}^+$ MEAN LIFE

The fit assumes the Ω^- and $\bar{\Omega}^+$ mean lives are the same, and averages them together.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.821 ± 0.011 OUR FIT				
$0.823 \pm 0.031 \pm 0.022$	1801	CHAN	98	E756 p Be, 800 GeV

$(\tau_{\Omega^-} - \tau_{\bar{\Omega}^+}) / \tau_{\Omega^-}$

A test of *CPT* invariance. Our calculation, from the averages in the preceding two data blocks.

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.00±0.05 OUR ESTIMATE	

Ω^- MAGNETIC MOMENT

<u>VALUE (μ_N)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-2.02 ±0.05 OUR AVERAGE				
-2.024±0.056	235k	WALLACE	95	SPEC Ω^- 300–550 GeV
-1.94 ± 0.17 ± 0.14	25k	DIEHL	91	SPEC Spin-transfer production

Ω^- DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Lambda K^-$	(67.7 ± 0.7) %	
$\Gamma_2 \Xi^0 \pi^-$	(24.3 ± 0.7) %	S=1.5
$\Gamma_3 \Xi^- \pi^0$	(8.55 ± 0.33) %	
$\Gamma_4 \Xi^- \pi^+ \pi^-$	(3.7 $^{+0.7}_{-0.6}$) $\times 10^{-4}$	
$\Gamma_5 \Xi(1530)^0 \pi^-$	< 7 $\times 10^{-5}$	CL=90%
$\Gamma_6 \Xi^0 e^- \bar{\nu}_e$	(5.6 ± 2.8) $\times 10^{-3}$	
$\Gamma_7 \Xi^- \gamma$	< 4.6 $\times 10^{-4}$	CL=90%
$\Delta S = 2$ forbidden (S2) modes		
$\Gamma_8 \Lambda \pi^-$	$S2 < 2.9 \times 10^{-6}$	CL=90%

Ω^- BRANCHING RATIOS

The BOURQUIN 84 values (which include results of BOURQUIN 79B, a separate experiment) are much more accurate than any other results, and so the other results have been omitted.

<u>$\Gamma(\Lambda K^-)/\Gamma_{\text{total}}$</u>	<u>Γ_1/Γ</u>
67.7±0.7 OUR AVERAGE	
66.3±0.8±2.0	ABLIKIM 23BJ BES3 $e^+ e^- \rightarrow \psi(3686) \rightarrow \Omega \bar{\Omega}$
67.8±0.7	BOURQUIN 84 SPEC SPS hyperon beam
• • • We do not use the following data for averages, fits, limits, etc. • • •	
68.6±1.3	BOURQUIN 79B SPEC See BOURQUIN 84

$\Gamma(\Xi^0\pi^-)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
24.3 ± 0.7 OUR AVERAGE		Error includes scale factor of 1.5.			
25.03 ± 0.44 ± 0.53	5.4k	ABLIKIM 23BJ BES3	e ⁺ e ⁻ → ψ(3686) → Ω̄Ω		■
23.6 ± 0.7	1947	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
23.4 ± 1.3	317	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^-\pi^0)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.55 ± 0.33 OUR AVERAGE					
8.43 ± 0.52 ± 0.28	794	ABLIKIM 23BJ BES3	e ⁺ e ⁻ → ψ(3686) → Ω̄Ω		■
8.6 ± 0.4	759	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
8.0 ± 0.8	145	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^-\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.74^{+0.67}_{-0.56}	100	5 KAMAEV	10	HYCP p Cu, 800 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.3 ^{+3.4} _{-1.3}	4	BOURQUIN 84	SPEC	SPS hyperon beam	

⁵ This KAMAEV 10 value uses 76 $\Omega^- \rightarrow \Xi^-\pi^+\pi^-$ and 24 $\bar{\Omega}^+ \rightarrow \Xi^+\pi^-\pi^+$ decays. The Ω^- and $\bar{\Omega}^+$ branching fractions measurements are statistically equal. The errors given combine statistical and systematic contributions. The *CP* branching-fraction asymmetry, $(\Omega^- - \bar{\Omega}^+)/\text{sum}$, is $+0.12 \pm 0.20$.

$\Gamma(\Xi(1530)^0\pi^-)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (units 10 ⁻⁴)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.7	90		KAMAEV 10	HYCP	p Cu, 800 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.4 ^{+5.1} _{-2.0}	4	6 BOURQUIN 84	SPEC	SPS hyperon beam	

⁶ The same 4 events as in the previous mode, with the isospin factor to take into account $\Xi(1530)^0 \rightarrow \Xi^0\pi^0$ decays included. BOURQUIN 84 adopted a theoretical assumption that $\Xi(1530)^0\pi^-$ would dominate $\Xi^-\pi^+\pi^-$ decay.

$\Gamma(\Xi^0e^-\bar{\nu}_e)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.6±2.8	14	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~10	3	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^-\gamma)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (units 10 ⁻⁴)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 4.6	90	0	ALBUQUERQ...94	E761	Ω^- 375 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<22	90	9	BOURQUIN 84	SPEC	SPS hyperon beam
<31	90	0	BOURQUIN 79B	SPEC	See BOURQUIN 84

$\Gamma(\Lambda\pi^-)/\Gamma_{\text{total}}$	Γ_8/Γ
$\Delta S=2$. Forbidden in first-order weak interaction.	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>
< 2.9	90
WHITE 05 HYCP p Cu, 800 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 190	90
BOURQUIN 84 SPEC SPS hyperon beam	
<1300	90
BOURQUIN 79B SPEC See BOURQUIN 84	

Ω^- DECAY PARAMETERS

$\alpha(\Omega^-) \alpha_-(\Lambda)$ FOR $\Omega^- \rightarrow \Lambda K^-$

Some early results have been omitted.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0115±0.0015 OUR AVERAGE				
0.0133±0.0033±0.0052	960k	7 CHEN 05	HYCP	p Cu, 800 GeV
0.0114±0.0012±0.0010	4.5M	7 LU 05A	HYCP	p Cu, 800 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.018 ± 0.030	6953	CHAN 98	E756	p Be, 800 GeV
-0.022 ± 0.051	1743	LUK 88	SPEC	p Be 400 GeV
-0.016 ± 0.018	12k	BOURQUIN 84	SPEC	SPS hyperon beam

⁷ The results of CHEN 05 and LU 05A are from different experimental runs.

α FOR $\Omega^- \rightarrow \Lambda K^-$

The above average, $\alpha(\Omega^-)\alpha_-(\Lambda) = 0.0115 \pm 0.0015$, divided by our current average $\alpha_-(\Lambda) = 0.747 \pm 0.009$ gives $\alpha(\Omega^-)$:

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.0154±0.0020 OUR EVALUATION	

$\bar{\alpha}$ FOR $\bar{\Omega}^+ \rightarrow \bar{\Lambda} K^+$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.0181±0.0028±0.0026	1.89M	LU 06	HYCP	p Cu, 800 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.017 ± 0.077	1823	CHAN 98	E756	p Be, 800 GeV

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Omega^- \rightarrow \Lambda K^-$, $\bar{\Omega}^+ \rightarrow \bar{\Lambda} K^+$

Zero if CP is conserved.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.016±0.092±0.089	8 LU 06	HYCP	p Cu, 800 GeV

⁸ This value uses the results of CHEN 05, LU 05A, and LU 06.

α FOR $\Omega^- \rightarrow \Xi^0 \pi^-$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.09±0.14	1630	BOURQUIN 84	SPEC	SPS hyperon beam

α FOR $\Omega^- \rightarrow \Xi^- \pi^0$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.05±0.21	614	BOURQUIN 84	SPEC	SPS hyperon beam

Ω^- REFERENCES

We have omitted some papers that have been superseded by later experiments. See our earlier editions.

ABLIKIM	23BJ	PR D108 L091101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21E	PRL 126 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
KAMAEV	10	PL B693 236	O. Kamaev <i>et al.</i>	(FNAL HyperCP Collab.)
AUBERT,BE	06	PRL 97 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LU	06	PRL 96 242001	L.C. Lu <i>et al.</i>	(FNAL HyperCP Collab.)
CHEN	05	PR D71 051102	Y.C. Chen <i>et al.</i>	(FNAL HyperCP Collab.)
LU	05A	PL B617 11	L.C. Lu <i>et al.</i>	(FNAL HyperCP Collab.)
WHITE	05	PRL 94 101804	C.G. White <i>et al.</i>	(FNAL HyperCP Collab.)
CHAN	98	PR D58 072002	A.W. Chan <i>et al.</i>	(FNAL E756 Collab.)
WALLACE	95	PRL 74 3732	N.B. Wallace <i>et al.</i>	(MINN, ARIZ, MICH+)
ALBUQUERQ...	94	PR D50 18	I.F. Albuquerque <i>et al.</i>	(FNAL E761 Collab.)
DIEHL	91	PRL 67 804	H.T. Diehl <i>et al.</i>	(RUTG, FNAL, MICH+)
LUK	88	PR D38 19	K.B. Luk <i>et al.</i>	(RUTG, WISC, MICH, MINN)
HARTOUNI	85	PRL 54 628	E.P. Hartouni <i>et al.</i>	(COLU, ILL, FNAL)
BOURQUIN	84	NP B241 1	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
Also		PL 87B 297	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
BOURQUIN	79B	PL 88B 192	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
BAUBILLIER	78	PL 78B 342	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+) J
DEUTSCH...	78	PL 73B 96	M. Deutschmann <i>et al.</i>	(AACH3, BERL, CERN+) J
HEMINGWAY	78	NP B142 205	R.J. Hemingway <i>et al.</i>	(CERN, ZEEM, NIJM+)
DIBIANCA	75	NP B98 137	F.A. Dibianca, R.J. Endorf	(CMU)
ALVAREZ	73	PR D8 702	L.W. Alvarez	(LBL)
DEUTSCH...	73	NP B61 102	M. Deutschmann <i>et al.</i>	(ABCLV Collab.)
FIRESTONE	71B	PRL 26 410	I. Firestone <i>et al.</i>	(LRL)
SPETH	69	PL 29B 252	R. Speth <i>et al.</i>	(AACH, BERL, CERN, LOIC+)
PALMER	68	PL 26B 323	R.B. Palmer <i>et al.</i>	(BNL, SYRA)
SCHULTZ	68	PR 168 1509	P.F. Schultz <i>et al.</i>	(ILL, ANL, NWES+)
SCOTTER	68	PL 26B 474	D. Scotter <i>et al.</i>	(BIRM, GLAS, LOIC+)
ABRAMS	64	PRL 13 670	G.S. Abrams <i>et al.</i>	(UMD, NRL)
BARNES	64	PRL 12 204	V.E. Barnes <i>et al.</i>	(BNL)
FRY	55	PR 97 1189	W.F. Fry, J. Schneps, M.S. Swami	(WISC)
FRY	55B	NC 2 346	W.F. Fry, J. Schneps, M.S. Swami	(WISC)
EISENBERG	54	PR 96 541	Y. Eisenberg	(CORN)