

$T_{cc}(3875)^+$ $I(J^P) = ?(??)$

OMITTED FROM SUMMARY TABLE

Observed with large significance by AAIJ 22E in the doubly-charmed
($C = 2$) decay mode $D^0 D^0 \pi^+$ using inclusive pp collisions at 7, 8,
and 13 TeV.

 $T_{cc}(3875)^+$ MASS

OUR FIT value comes from the measurement of $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$
below and $m_{D^{*+}} + m_{D^0}$ values.

VALUE (MeV)	DOCUMENT ID
3874.83 ± 0.11 OUR FIT	

 $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.27 ± 0.06 OUR FIT				
$-0.273 \pm 0.061^{+0.012}_{-0.015}$	117	¹ AAIJ	22E LHCb	$pp \rightarrow D^0 D^0 \pi^+ X$

¹ The fit assumes a relativistic P -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV^{-1} . The significance for $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0}) < 0$ is 4.3σ .

 $T_{cc}(3875)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.410 \pm 0.165^{+0.047}_{-0.057}$	117	¹ AAIJ	22E LHCb	$pp \rightarrow D^0 D^0 \pi^+ X$

¹ The fit assumes a relativistic P -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV^{-1} .

 $T_{cc}(3875)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad D^0 D^0 \pi^+$	seen

 $T_{cc}(3875)^+$ BRANCHING RATIOS

$\Gamma(D^0 D^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	117	AAIJ	22E LHCb	$pp \rightarrow D^0 D^0 \pi^+ X$

 $T_{cc}(3875)^+$ REFERENCES

AAIJ	22E NATP 18 751	R. Aaij et al.	(LHCb Collab.)
------	-----------------	----------------	----------------