
 $I^G(J^{PC}) = 0^-(1^{--})$

also known as $Y(4360)$; was $X(4360)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

$\psi(4360)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4374 ± 7 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
4371.6 ± 2.5 ± 9.2	1	ABLIKIM	22AL BES3	$e^+e^- \rightarrow \pi^+\pi^-D^+D^-$
4298 ± 12 ± 26	2	ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
4390.3 ± 6.0 ± 0.7	3	ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4371.7 ± 7.5 ± 1.8	4	ABLIKIM	21AK BES3	$e^+e^- \rightarrow \gamma\chi_c2 \rightarrow \gamma\gamma J/\psi$
4382.0 ± 13.3 ± 1.7	5	ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$
4391.5 ± 6.3 ± 1.0		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$
4347 ± 6 ± 3	279	6 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4340 ± 16 ± 9	37	7 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4406.9 ± 17.2 ± 4.5	8	ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
4320.0 ± 10.4 ± 7.0	9	ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
4383.8 ± 4.2 ± 0.8	10	ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4383.7 ± 2.9 ± 6.2	11	ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4386.4 ± 2.1 ± 6.4	12	ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
4355 ± 9 ± 9	74	13 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4324 ± 24		14 AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4361 ± 9 ± 9	47	7 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

¹ From a fit to the cross section for $e^+e^- \rightarrow D^+D^-\pi^+\pi^-$ in the range $\sqrt{s} = 4.190\text{--}4.946$ GeV.

² From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730\text{--}4.7008$ GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.

³ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.

⁴ From a five-resonance fit to the cross section for $e^+e^- \rightarrow \gamma\gamma J/\psi \rightarrow \gamma\gamma\ell^+\ell^-$.

⁵ From a fit of the measured cross section in the range $\sqrt{s} = 3.808\text{--}4.600$ GeV.

⁶ From a two-resonance fit. Supersedes WANG 07D.


⁷ From a two-resonance fit.

⁸ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.

⁹ From a three-resonance fit. Superseded by ABLIKIM 22AM.

¹⁰ From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.

- 11 From a three-resonance fit.
 12 From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.
 13 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.
 14 From a single-resonance fit. Systematic errors not estimated.



psi(4360) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
118 \pm12 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
167 \pm 4 \pm 29		1 ABLIKIM	22AL BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D^+ D^-$
127 \pm 17 \pm 10		2 ABLIKIM	22AM BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
143.3 \pm 10.0 \pm 0.5		3 ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
51.1 \pm 17.6 \pm 1.9		4 ABLIKIM	21AK BES3	$e^+ e^- \rightarrow \gamma \chi_{c2} \rightarrow \gamma \gamma J/\psi$
135.8 \pm 60.8 \pm 22.5		5 ABLIKIM	20O BES3	$e^+ e^- \rightarrow \eta J/\psi$
139.5 $^{+16.2}_{-20.6}$ \pm 0.6		ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$
103 \pm 9 \pm 5 279		6 WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
94 \pm 32 \pm 13 37		7 LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
128.1 \pm 37.2 \pm 2.3		8 ABLIKIM	22R BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
101.4 $^{+25.3}_{-19.7}$ \pm 10.2		9 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
84.2 \pm 12.5 \pm 2.1		10 ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
94.2 \pm 7.3 \pm 2.0		11 ZHANG	17B RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$

$96.0 \pm 6.7 \pm 2.7$	12	ZHANG	17C	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
$103 \begin{array}{l} +17 \\ -15 \end{array} \pm 11$	74	LIU	08H	RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
172 ± 33		AUBERT	07S	BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$74 \pm 15 \pm 10$	47	WANG	07D	BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

¹ From a fit to the cross section for $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$ in the range $\sqrt{s} = 4.190\text{--}4.946$ GeV.

² From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730\text{--}4.7008$ GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.

³ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.

⁴ From a five-resonance fit to the cross section for $e^+ e^- \rightarrow \gamma \gamma J/\psi \rightarrow \gamma \gamma \ell^+ \ell^-$.

⁵ From a fit of the measured cross section in the range $\sqrt{s} = 3.808\text{--}4.600$ GeV.

⁶ From a two-resonance fit. Supersedes WANG 07D.

⁷ From a two-resonance fit.

⁸ From a fit to the $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.

⁹ From a three-resonance fit. Superseded by ABLIKIM 22AM.


¹⁰ From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.

¹¹ From a three-resonance fit.

¹² From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.

¹³ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

¹⁴ From a single-resonance fit. Systematic errors not estimated.



$\psi(4360)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	seen
$\Gamma_2 h_c \pi^+ \pi^-$	seen
$\Gamma_3 J/\psi \pi^+ \pi^-$	seen
$\Gamma_4 \psi(2S) \pi^+ \pi^-$	seen
$\Gamma_5 \psi(3770) \pi^+ \pi^-$	possibly seen
$\Gamma_6 \psi_2(3823) \pi^+ \pi^-$	seen
$\Gamma_7 J/\psi \eta$	seen
$\Gamma_8 D^0 D^{*-} \pi^+$	not seen
$\Gamma_9 D^+ D^- \pi^+ \pi^-$	seen
$\Gamma_{10} D_1(2420) \bar{D} + \text{c.c.}$	possibly seen
$\Gamma_{11} \phi \eta$	not seen
$\Gamma_{12} \omega \pi^0$	not seen
$\Gamma_{13} \omega \eta$	not seen
$\Gamma_{14} p \bar{p} \eta$	not seen
$\Gamma_{15} p \bar{p} \omega$	not seen
$\Gamma_{16} \chi_{c1} \gamma$	not seen
$\Gamma_{17} \chi_{c2} \gamma$	not seen
$\Gamma_{18} \Xi^- \Xi^+$	not seen
$\Gamma_{19} p K^- \bar{\Lambda} + \text{c.c.}$	not seen

$$\psi(4360) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$$

$\Gamma(h_c \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1 / \Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$11.6^{+5.0}_{-4.4} \pm 1.9$	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$

$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$		$\Gamma_4 \Gamma_1 / \Gamma$		
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.7 \pm 4.1		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
20.7 \pm 2.5		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
9.9 \pm 4.1		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
19.4 \pm 2.0		⁴ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
7.3 \pm 2.8		⁵ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.0 \pm 3.8		⁶ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
9.2 \pm 0.6 \pm 0.6	279	⁷ WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
10.9 \pm 0.6 \pm 0.7	279	⁸ WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
6.0 \pm 1.0 \pm 0.5	37	⁵ LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.2 \pm 1.0 \pm 0.6	37	⁶ LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
11.1 \pm 1.3	74	⁹ LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

12.3 ± 1.2	74	¹⁰ LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$10.4 \pm 1.7 \pm 1.5$	47	⁵ WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$11.8 \pm 1.8 \pm 1.4$	47	⁶ WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

¹ Solution I of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

² Solution II of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

³ Solution III of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

⁴ Solution IV of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

⁵ Solution I of two equivalent solutions in a fit using two interfering resonances.

⁶ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁷ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

⁸ Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

⁹ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

¹⁰ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_7\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

¹ 3.4 ± 2.2 ABLIKIM 200 BES3 $e^+e^- \rightarrow \eta J/\psi$

² 1.5 ± 1.0 ABLIKIM 200 BES3 $e^+e^- \rightarrow \eta J/\psi$

³ 1.7 ± 1.1 ABLIKIM 200 BES3 $e^+e^- \rightarrow \eta J/\psi$

<6.8 90 WANG 13B BELL $e^+e^- \rightarrow J/\psi\eta\gamma$

¹ Solution 1 of three equivalent fit solutions using three resonant structures.

² Solution 2 of three equivalent fit solutions using three resonant structures.

³ Solution 3 of three equivalent fit solutions using three resonant structures.

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{16}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.57	90	¹ HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{17}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	¹ HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{18}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.0448	90	¹ ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4360)$

¹ From a fit to $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ cross sections.

$\Gamma(pK^-\bar{\Lambda}+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{19}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<4.7 \times 10^{-3}$	90	¹ ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4360)$

¹ From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda}+c.c.$ cross sections.

$\psi(4360)$ BRANCHING RATIOS

$\Gamma(h_c\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	ABLIKIM	17G	$e^+e^- \rightarrow \pi^+\pi^- h_c$	

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	¹ ABLIKIM	17v	BES3 $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	

¹ From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb⁻¹.

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
-------	-------------	------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(0.81 \pm 0.12 \pm 0.13)$ to $(42 \pm 15 \pm 15)$	¹ ZHANG	17C	RVUE $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$	
--	--------------------	-----	---	--

¹ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

$\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
possibly seen	¹ ABLIKIM	19AR	BES3 $e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$	

¹ Observe $e^+e^- \rightarrow \pi^+\pi^-\psi(3770)$ at $\sqrt{s} = 4.26, 4.36,$ and 4.42 GeV but cannot establish if continuum or resonant.

$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
seen		¹ ABLIKIM	22R	BES3 $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	19	² ABLIKIM	15S	BES3 $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$	
---------------	----	----------------------	-----	--	--

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.

² From a fit of $e^+e^- \rightarrow \pi^+\pi^-\psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1}\gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4360)$ line shape.

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
seen	¹ ABLIKIM	200	BES3 $e^+e^- \rightarrow \eta J/\psi$	

¹ With a significance of 6.0 σ .

$\Gamma(D^0 D^{*-}\pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma \times \Gamma_1/\Gamma$
$<0.72 \times 10^{-6}$	90	¹ PAKHLOVA	09	BELL $e^+e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-}\pi^+$	

¹ Using $4355^{+9}_{-10} \pm 9$ MeV for the mass of $\psi(4360)$.

$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$					Γ_8/Γ_4
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<8	90	PAKHLOVA	09	BELL	$e^+ e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-} \pi^+$
$\Gamma(D^+ D^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	1 ABLIKIM	22AL BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D^+ D^-$		
1 From a fit to the cross section for $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$ in the range $\sqrt{s} = 4.190\text{--}4.946$ GeV.					
$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
possibly seen	1 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$		
1 Evidence for $e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.					
$\Gamma(\phi\eta)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	23BT BES3	$e^+ e^- \rightarrow \phi\eta$		
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega\pi^0$		
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega\eta$		
$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$					Γ_{14}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	21AN BES3	$e^+ e^- \rightarrow p\bar{p}\eta$		
$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$					Γ_{15}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	21AN BES3	$e^+ e^- \rightarrow p\bar{p}\omega$		

$\psi(4360)$ REFERENCES

ABLIKIM	23BK JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BL JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BT PR D108 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AL PR D106 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AM PR D106 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22K JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22R PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AJ PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AK PR D104 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AN PR D104 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20O PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19K PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17B PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	17G	PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan	
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)