

$\Sigma(2030) \ 7/2^+$ $I(J^P) = 1(\frac{7}{2}^+)$ Status: ***

Discovered by COOL 66 and by WOHL 66. For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and invariant-mass distributions around 2030 MeV may be found in our 1984 edition, Reviews of Modern Physics **56** S1 (1984).

 $\Sigma(2030)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2010 to 2030 (\approx 2020) OUR ESTIMATE			
2014 \pm 6	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
2025 $^{+10}_{-5}$	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1993	ZHANG 13A	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

-2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
130 to 190 (\approx 160) OUR ESTIMATE			
172 \pm 12	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
130 $^{+6}_{-24}$	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
176	ZHANG 13A	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

 $\Sigma(2030)$ POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.20 \pm 0.04	-38 \pm 8	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.220	-38	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma\pi$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.07 \pm 0.02	165 \pm 12	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0807 135 ¹ KAMANO 15 DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda\pi$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.04	-22 ± 12	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.138 -24 ¹ KAMANO 15 DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Xi K$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.01 ± 0.01		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0348 129 ¹ KAMANO 15 DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma(1385)\pi$, F-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.04 ± 0.03		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.089 -23 ¹ KAMANO 15 DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma(1385)\pi$, H-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.0245	132	¹ KAMANO 15	DPWA	Multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, D-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.03 ± 0.02	-100 ± 40	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, G-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.02 ± 0.02		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta\bar{K}$, F-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06	-130 ± 20	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta\bar{K}$, H-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.04 ± 0.02	-130 ± 35	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=1/2$, F-wave

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 ± 0.02		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.193	38	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=3/2$, F-wave

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.16 ± 0.09	-160 ± 40	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.320	37	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=3/2$, H-wave

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.00358	22	¹ KAMANO 15	DPWA	Multichannel

¹ From the preferred solution A in KAMANO 15.

 $\Sigma(2030)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2025 to 2040 (≈ 2030) OUR ESTIMATE			
2032 ± 6	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
2030 ± 5	ZHANG 13A	DPWA	$\bar{K}N$ multichannel
2036 ± 5	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2038 ± 10	CORDEN 77B		$K^- N \rightarrow N\bar{K}^*$
2030 ± 3	¹ CORDEN 76	DPWA	$K^- n \rightarrow \Lambda\pi^-$
2035 ± 15	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
2038 ± 10	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$
2042 ± 11	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
2020 ± 6	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
2035 ± 10	LITCHFIELD 74B	DPWA	$K^- p \rightarrow \Lambda(1520)\pi^0$
2020 ± 30	LITCHFIELD 74C	DPWA	$K^- p \rightarrow \Delta(1232)\bar{K}$
2025 ± 10	LITCHFIELD 74D	DPWA	$K^- p \rightarrow \Lambda(1820)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2040 ± 5	GOPAL 77	DPWA	$\bar{K}N$ multichannel
2027 to 2057	GOYAL 77	DPWA	$K^- N \rightarrow \Sigma\pi$
2030	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$

¹ Preferred solution 3; see CORDEN 76 for other possibilities.

 $\Sigma(2030)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
150 to 200 (≈ 180) OUR ESTIMATE			
177 ± 12	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
207 ± 17	ZHANG 13A	DPWA	$\bar{K}N$ multichannel
172 ± 10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
137 ± 40	CORDEN 77B		$K^- N \rightarrow N\bar{K}^*$

201 ± 9	1 CORDEN	76	DPWA	$K^- n \rightarrow \Lambda \pi^-$
180 ± 20	BAILLON	75	IPWA	$\bar{K} N \rightarrow \Lambda \pi$
172 ± 15	HEMINGWAY	75	DPWA	$K^- p \rightarrow \bar{K} N$
178 ± 13	VANHORN	75	DPWA	$K^- p \rightarrow \Lambda \pi^0$
111 ± 5	KANE	74	DPWA	$K^- p \rightarrow \Sigma \pi$
160 ± 20	LITCHFIELD	74B	DPWA	$K^- p \rightarrow \Lambda(1520) \pi^0$
200 ± 30	LITCHFIELD	74C	DPWA	$K^- p \rightarrow \Delta(1232) \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

260	DECLAIS	77	DPWA	$\bar{K} N \rightarrow \bar{K} N$
190 ± 10	GOPAL	77	DPWA	$\bar{K} N$ multichannel
126 to 195	GOYAL	77	DPWA	$K^- N \rightarrow \Sigma \pi$
160	DEBELLEFON	76	IPWA	$K^- p \rightarrow \Lambda \pi^0$
70 to 125	LITCHFIELD	74D	DPWA	$K^- p \rightarrow \Lambda(1820) \pi^0$

¹ Preferred solution 3; see CORDEN 76 for other possibilities.

$\Sigma(2030)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N \bar{K}$	17–23 %
$\Gamma_2 \Lambda \pi$	17–23 %
$\Gamma_3 \Sigma \pi$	5–10 %
$\Gamma_4 \Xi K$	<2 %
$\Gamma_5 \Sigma(1385) \pi$	5–15 %
$\Gamma_6 \Sigma(1385) \pi$, <i>F</i> -wave	(1.0 ± 1.0) %
$\Gamma_7 \Sigma(1385) \pi$, <i>H</i> -wave	
$\Gamma_8 \Lambda(1520) \pi$	10–20 %
$\Gamma_9 \Lambda(1520) \pi$, <i>D</i> -wave	
$\Gamma_{10} \Lambda(1520) \pi$, <i>G</i> -wave	
$\Gamma_{11} \Delta(1232) \bar{K}$	10–20 %
$\Gamma_{12} \Delta(1232) \bar{K}$, <i>F</i> -wave	(15 ± 5) %
$\Gamma_{13} \Delta(1232) \bar{K}$, <i>H</i> -wave	(1.0 ± 1.0) %
$\Gamma_{14} N \bar{K}^*(892)$	
$\Gamma_{15} N \bar{K}^*(892)$, $S=1/2$, <i>F</i> -wave	
$\Gamma_{16} N \bar{K}^*(892)$, $S=3/2$, <i>F</i> -wave	(14 ± 8) %
$\Gamma_{17} N \bar{K}^*(892)$, $S=3/2$, <i>H</i> -wave	
$\Gamma_{18} \Lambda(1820) \pi$, <i>P</i> -wave	

$\Sigma(2030)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.17 to 0.23 OUR ESTIMATE			
0.20 ± 0.04	SARANTSEV 19	DPWA	$\bar{K} N$ multichannel
0.13 ± 0.01	ZHANG 13A	DPWA	$\bar{K} N$ multichannel

0.19 ± 0.03	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.18 ± 0.03	HEMINGWAY	75	DPWA $K^- p \rightarrow \bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.269	¹ KAMANO	15	DPWA Multichannel
0.15	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.24 ± 0.02	GOPAL	77	DPWA See GOPAL 80

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.04	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.080	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.025 ± 0.008	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.037	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Xi K)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.006	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda(1520)\pi, D\text{-wave})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
~ 0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

$\Gamma(\Lambda(1520)\pi, G\text{-wave})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

$\Gamma(\Sigma(1385)\pi, F\text{-wave})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.01 ± 0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.030	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma(1385)\pi, H\text{-wave})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.003	¹ KAMANO	15	Multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Delta(1232)\bar{K}, F\text{-wave})/\Gamma_{\text{total}}$	Γ_{12}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.15±0.05	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

$\Gamma(\Delta(1232)\bar{K}, H\text{-wave})/\Gamma_{\text{total}}$	Γ_{13}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.01±0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

$\Gamma(N\bar{K}^*(892), S=1/2, F\text{-wave})/\Gamma_{\text{total}}$	Γ_{15}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.154	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, F\text{-wave})/\Gamma_{\text{total}}$	Γ_{16}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14 ±0.08	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.422	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, H\text{-wave})/\Gamma_{\text{total}}$	Γ_{17}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
not seen	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel
¹ From the preferred solution A in KAMANO 15.			

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda\pi$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.15 ±0.01	ZHANG 13A	DPWA	Multichannel
+0.18 ±0.02	GOPAL 77	DPWA	$\bar{K}N$ multichannel
+0.20 ±0.01	¹ CORDEN 76	DPWA	$K^- n \rightarrow \Lambda\pi^-$
+0.18 ±0.02	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
+0.20 ±0.01	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
+0.195±0.053	DEVENISH 74B		Fixed- <i>t</i> dispersion rel.
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.20	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$

¹ Preferred solution 3; see CORDEN 76 for other possibilities.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma\pi$	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.08 ±0.01	ZHANG 13A	DPWA	Multichannel
-0.09 ±0.01	¹ CORDEN 77C		$K^- n \rightarrow \Sigma\pi$
-0.06 ±0.01	¹ CORDEN 77C		$K^- n \rightarrow \Sigma\pi$
-0.15 ±0.03	GOPAL 77	DPWA	$\bar{K}N$ multichannel
-0.10 ±0.01	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.085 ± 0.02 2 GOYAL 77 DPWA $K^- N \rightarrow \Sigma \pi$

¹ The two entries for CORDEN 77C are from two different acceptable solutions.

² This coupling is extracted from unnormalized data.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Xi K$ $(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.023	MULLER 69B	DPWA	$K^- p \rightarrow \Xi K$
<0.05	BURGUN 68	DPWA	$K^- p \rightarrow \Xi K$
<0.05	TRIPP 67	RVUE	$K^- p \rightarrow \Xi K$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma(1385)\pi$, F-wave $(\Gamma_1 \Gamma_6)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.16 \pm 0.01$	ZHANG 13A	DPWA	Multichannel
$+0.153 \pm 0.026$	¹ CAMERON 78	DPWA	$K^- p \rightarrow \Sigma(1385)\pi$

¹ The published sign has been changed to be in accord with the baryon-first convention.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, D-wave $(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.114 \pm 0.010$	¹ CAMERON 77	DPWA	$K^- p \rightarrow \Lambda(1520)\pi^0$
0.14 ± 0.03	LITCHFIELD 74B	DPWA	$K^- p \rightarrow \Lambda(1520)\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.10 ± 0.03 2 CORDEN 75B DBC $K^- n \rightarrow N\bar{K}\pi^-$

¹ The published sign has been changed to be in accord with the baryon-first convention.

² An upper limit.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, G-wave $(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.146 \pm 0.010$	¹ CAMERON 77	DPWA	$K^- p \rightarrow \Lambda(1520)\pi^0$
0.02 ± 0.02	LITCHFIELD 74B	DPWA	$K^- p \rightarrow \Lambda(1520)\pi^0$

¹ The published sign has been changed to be in accord with the baryon-first convention.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}$, F-wave $(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.12 \pm 0.02$	ZHANG 13A	DPWA	Multichannel
0.16 ± 0.03	LITCHFIELD 74C	DPWA	$K^- p \rightarrow \Delta(1232)\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.03 1 CORDEN 75B DBC $K^- n \rightarrow N\bar{K}\pi^-$

¹ An upper limit.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}$, H-wave $(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.00 ± 0.02	LITCHFIELD 74C	DPWA	$K^- p \rightarrow \Delta(1232)\bar{K}$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=1/2$, F -wave

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_{15})^{1/2} / \Gamma$
$+0.06 \pm 0.02$	ZHANG 13A	DPWA	Multichannel	
$+0.06 \pm 0.03$	¹ CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$	
-0.02 ± 0.01	CORDEN 77B		$K^- d \rightarrow NN\bar{K}^*$	

¹ The published sign has been changed to be in accord with the baryon-first convention.

 $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=3/2$, F -wave

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_{16})^{1/2} / \Gamma$
$+0.05 \pm 0.01$	ZHANG 13A	DPWA	Multichannel	
$+0.04 \pm 0.03$	¹ CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$	
-0.12 ± 0.02	CORDEN 77B		$K^- d \rightarrow NN\bar{K}^*$	

¹ The upper limit on the G_3 wave is 0.03.

 $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1820)\pi$, P -wave

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_{18})^{1/2} / \Gamma$
0.14 ± 0.02	CORDEN 75B	DBC	$K^- n \rightarrow N\bar{K}\pi^-$	
0.18 ± 0.04	LITCHFIELD 74D	DPWA	$K^- p \rightarrow \Lambda(1820)\pi^0$	

 $\Sigma(2030)$ REFERENCES

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO	15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
PDG	84	RMP 56 S1	C.G. Wohl <i>et al.</i>	(LBL, CIT, CERN)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
CAMERON	78	NP B143 189	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
CAMERON	78B	NP B146 327	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
CORDEN	77B	NP B121 365	M.J. Corden <i>et al.</i>	(BIRM) IJP
CORDEN	77C	NP B125 61	M.J. Corden <i>et al.</i>	(BIRM) IJP
DECLAIS	77	CERN 77-16	Y. Declais <i>et al.</i>	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
GOYAL	77	PR D16 2746	D.P. Goyal, A.V. Sodhi	(DELH) IJP
CORDEN	76	NP B104 382	M.J. Corden <i>et al.</i>	(BIRM) IJP
DEBELLEFON	76	NP B109 129	A. de Bellefon, A. Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	P.H. Baillon, P.J. Litchfield	(CERN, RHEL) IJP
CORDEN	75B	NP B92 365	M.J. Corden <i>et al.</i>	(BIRM) IJP
HEMINGWAY	75	NP B91 12	R.J. Hemingway <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
VANHORN	75	NP B87 145	A.J. van Horn	(LBL) IJP
Also		NP B87 157	A.J. van Horn	(LBL) IJP
DEVENISH	74B	NP B81 330	R.C.E. Devenish, C.D. Froggatt, B.R. Martin	(DESY+) IJP
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
LITCHFIELD	74B	NP B74 19	P.J. Litchfield <i>et al.</i>	(CERN, HEIDH) IJP
LITCHFIELD	74C	NP B74 39	P.J. Litchfield <i>et al.</i>	(CERN, HEIDH) IJP
LITCHFIELD	74D	NP B74 12	P.J. Litchfield <i>et al.</i>	(CERN, HEIDH) IJP
MULLER	69B	Thesis UCRL 19372	R.A. Muller	(LRL) IJP
BURGUN	68	NP B8 447	G. Burgun <i>et al.</i>	(SACL, CDEF, RHEL) IJP
TRIPP	67	NP B3 10	R.D. Tripp <i>et al.</i>	(LRL, SLAC, CERN+) IJP
COOL	66	PRL 16 1228	R.L. Cool <i>et al.</i>	(BNL) IJP
WOHL	66	PRL 17 107	C.G. Wohl, F.T. Solmitz, M.L. Stevenson	(LRL) IJP