

**$D(3000)^0$**  $I(J^P) = \frac{1}{2}(??)$ 

## OMITTED FROM SUMMARY TABLE

Both natural- and unnatural-parity components observed depending on the decay mode (AAIJ 13CC).

 **$D(3000)^0$  MASS**

| VALUE (MeV)   | EVTS  | DOCUMENT ID         | TECN      | COMMENT                           |
|---|-------|---------------------|-----------|-----------------------------------|
| <b>3214 ± 29 ± 49</b>   | 28k   | <sup>1</sup> AAIJ   | 16AH LHCb | $B^- \rightarrow D^+ \pi^- \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                     |           |                                   |
| 2971.8 ± 8.7  | 9.5k  | <sup>2,3</sup> AAIJ | 13CC LHCb | $p p \rightarrow D^{*+} \pi^- X$  |
| 3008.1 ± 4.0  | 17.6k | <sup>2,4</sup> AAIJ | 13CC LHCb | $p p \rightarrow D^+ \pi^- X$     |

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

<sup>2</sup> Systematic uncertainty not estimated.

<sup>3</sup> Unnatural parity preferred.

<sup>4</sup> Natural parity state. A state  $D(3000)^+$  is possibly seen in  $D^0 \pi^+$  final state.

 **$D(3000)^0$  WIDTH**

| VALUE (MeV)   | EVTS  | DOCUMENT ID         | TECN      | COMMENT                           |
|---|-------|---------------------|-----------|-----------------------------------|
| <b>186 ± 38 ± 72</b>  | 28k   | <sup>5</sup> AAIJ   | 16AH LHCb | $B^- \rightarrow D^+ \pi^- \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                     |           |                                   |
| 188.1 ± 44.8  | 9.5k  | <sup>6,7</sup> AAIJ | 13CC LHCb | $p p \rightarrow D^{*+} \pi^- X$  |
| 110.5 ± 11.5  | 17.6k | <sup>6,8</sup> AAIJ | 13CC LHCb | $p p \rightarrow D^+ \pi^- X$     |

<sup>5</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

<sup>6</sup> Systematic uncertainty not estimated.

<sup>7</sup> Unnatural parity preferred.

<sup>8</sup> Natural parity state. A state  $D(3000)^+$  is possibly seen in  $D^0 \pi^+$  final state.

 **$D(3000)^0$  DECAY MODES**

| Mode                      | Fraction ( $\Gamma_i/\Gamma$ ) |
|---------------------------|--------------------------------|
| $\Gamma_1$ $D^{*+} \pi^-$ | seen                           |

## **$D(3000)^0$ POLARIZATION AMPLITUDE $A_{D_J}$**

A polarization amplitude  $A_{D_J}$  is a parameter that depends on the initial polarization of the  $D_J$ . For  $D_J$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_J} \cos^2(\theta_H)$ , where  $\theta_H$  is the angle in the  $D_J$  rest frame between the two pions emitted in the  $D_J \rightarrow D^* \pi$  and  $D^* \rightarrow D \pi$  decays.

| <u>VALUE</u>   | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u>                              | <u>COMMENT</u> |
|--|-------------|--------------------|--|----------------|
| <b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b> |             |                    |  |                |
| $1.5 \pm 0.9$  | 9.5k        | <sup>9</sup> AAIJ  | 13CC LHCb $p p \rightarrow D^*+ \pi^- X$ |                |
| <sup>9</sup> Systematic uncertainty not estimated.                                   |             |                    |  |                |

## **$D(3000)^0$ REFERENCES**

|      |                    |                       |                |
|------|--------------------|-----------------------|----------------|
| AAIJ | 16AH PR D94 072001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13CC JHEP 1309 145 | R. Aaij <i>et al.</i> | (LHCb Collab.) |