

**$f_2(1270)$**  $I^G(J^{PC}) = 0^+(2^{++})$  **$f_2(1270)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma = -2 \operatorname{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1260–1283) – <math>i</math> (90–110) OUR ESTIMATE</b>			
(1268 ± 8) – $i$ (101 ± 6)	RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
(1263.3 ± 0.2 ± 1.5) – $i$ (96.9 ± 0.2 ± 0.8)	ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
(1270 ± 8) – $i$ (97 ± 8)	<sup>1</sup> ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$
(1278 ± 5) – $i$ (102 ± 10)	<sup>1</sup> BERTIN	97C	OBLX 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> Amplitude did not include dispersive corrections. **$f_2(1270)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1275.4 ± 0.8 OUR AVERAGE</b>				
1275.8 ± 1.0 ± 0.4		<sup>1</sup> BOGOLYUB...	13	SPEC $7\pi^+(K^+, p)A \rightarrow n\gamma + X$
1262 ± 1 ± 8		2 ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1275 ± 15		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
1283 ± 5		ALDE	98	GAM4 $100\pi^-p \rightarrow \pi^0\pi^0n$
1272 ± 8	200k	PROKOSHKIN	94	GAM2 $38\pi^-p \rightarrow \pi^0\pi^0n$
1269.7 ± 5.2	5730	AUGUSTIN	89	DM2 $e^+e^- \rightarrow 5\pi$
1283 ± 8	400	<sup>3</sup> ALDE	87	GAM4 $100\pi^-p \rightarrow 4\pi^0n$
1274 ± 5		3 AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
1283 ± 6		<sup>4</sup> LONGACRE	86	MPS $22\pi^-p \rightarrow n2K_S^0$
1276 ± 7		COURAU	84	DLCO $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
1273.3 ± 2.3		5 CHABAUD	83	ASPK $17\pi^-p$ polarized
1280 ± 4		<sup>6</sup> CASON	82	STRC $8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$
1281 ± 7	11600	GIDAL	81	MRK2 $J/\psi$ decay
1282 ± 5		<sup>7</sup> CORDEN	79	OMEG $12\text{--}15\pi^-p \rightarrow n2\pi$
1269 ± 4	10k	APEL	75	NICE $40\pi^-p \rightarrow n2\pi^0$
1272 ± 4	4600	ENGLER	74	DBC $6\pi^+n \rightarrow \pi^+\pi^-p$
1277 ± 4	5300	FLATTE	71	HBC $7.0\pi^+p$
1273 ± 8		<sup>3</sup> STUNTEBECK	70	HBC $8\pi^-p, 5.4\pi^+d$
1265 ± 8		BOESEBECK	68	HBC $8\pi^+p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1257 ± 6		<sup>8</sup> KLEMPT	22	RVUE $J/\psi(1S) \rightarrow \gamma\pi^0\pi^0, \gamma K_S^0 K_S^0$
1263 ± 12		CARVER	21	CLAS $\gamma p \rightarrow \pi^0\pi^0p$
1259 ± 4 ± 4	1.7k	<sup>9,10</sup> DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1267 ± 4 ± 3	1.5k	<sup>9,10</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1277 ± 6	870	<sup>11</sup> SCHEGELSKY	06A	RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
1251 ± 10		TIKHOMIROV	03	SPEC $40.0\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$

1260	$\pm 10$	12 ALDE	97	GAM2	450 $p p \rightarrow p p \pi^0 \pi^0$
1278	$\pm 6$	12 GRYGOREV	96	SPEC	40 $\pi^- N \rightarrow K_S^0 K_S^0 X$
1262	$\pm 11$	AGUILAR-...	91	EHS	400 $p p$
1275	$\pm 10$	AKER	91	CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
1220	$\pm 10$	BREAKSTONE	90	SFM	$p p \rightarrow p p \pi^+ \pi^-$
1288	$\pm 12$	ABACHI	86B	HRS	$e^+ e^- \rightarrow \pi^+ \pi^- X$
1284	$\pm 30$	3k	BINON	GAM2	38 $\pi^- p \rightarrow n 2\eta$
1280	$\pm 20$	3k	APEL	CNTR	25 $\pi^- p \rightarrow n 2\pi^0$
1284	$\pm 10$	16000	DEUTSCH...	HBC	16 $\pi^+ p$
1258	$\pm 10$	600	TAKAHASHI	HBC	8 $\pi^- p \rightarrow n 2\pi^-$
1275	$\pm 13$	ARMENISE	70	HBC	9 $\pi^+ n \rightarrow p \pi^+ \pi^-$
1261	$\pm 5$	1960	3 ARMENISE	DBC	5.1 $\pi^+ n \rightarrow p \pi^+ MM^-$
1270	$\pm 10$	360	3 ARMENISE	DBC	5.1 $\pi^+ n \rightarrow p \pi^0 MM$
1268	$\pm 6$	13 JOHNSON	68	HBC	3.7–4.2 $\pi^- p$

<sup>1</sup> Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.

<sup>2</sup> Breit-Wigner mass.

<sup>3</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>4</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>5</sup> From an energy-independent partial-wave analysis.

<sup>6</sup> From an amplitude analysis of the reaction  $\pi^+ \pi^- \rightarrow 2\pi^0$ .

<sup>7</sup> From an amplitude analysis of  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$  scattering data.

<sup>8</sup> Fit of the tensor partial waves from BES3 in the multipole basis.

<sup>9</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>10</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 185$  MeV.

<sup>11</sup> From analysis of L3 data at 91 and 183–209 GeV.

<sup>12</sup> Systematic uncertainties not estimated.

<sup>13</sup> JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

## $f_2(1270)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>186.6<math>^{+ 2.8}_{- 2.2}</math> OUR FIT</b>				Error includes scale factor of 1.5.
<b>185.8<math>^{+ 2.8}_{- 2.1}</math> OUR AVERAGE</b>				Error includes scale factor of 1.6. See the ideogram below.
190.3 $\pm 1.9 \pm 1.8$	1 BOGOLYUB...	13	SPEC	$7\pi^+(K^+, p)A \rightarrow n\gamma + X$
175 $^{+ 6}_{- 4} \pm 10$	2 ABLIKIM	06V	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
190 $\pm 20$	ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
171 $\pm 10$	ALDE	98	GAM4	$100\pi^- p \rightarrow \pi^0\pi^0 n$
192 $\pm 5$	200k	PROKOSHIN	94	GAM2 $38\pi^- p \rightarrow \pi^0\pi^0 n$
180 $\pm 24$	AGUILAR-...	91	EHS	400 $p p$
169 $\pm 9$	5730	3 AUGUSTIN	89	DM2 $e^+ e^- \rightarrow 5\pi$
150 $\pm 30$	400	3 ALDE	87	GAM4 $100\pi^- p \rightarrow 4\pi^0 n$
186 $^{+ 9}_{- 2}$	4 LONGACRE	86	MPS	$22\pi^- p \rightarrow n 2K_S^0$
179.2 $^{+ 6.9}_{- 6.6}$	5 CHABAUD	83	ASPK	$17\pi^- p$ polarized
160 $\pm 11$	DENNEY	83	LASS	$10\pi^+ N$

196	$\pm 10$	3k	APEL	82	CNTR	$25 \pi^- p \rightarrow n 2\pi^0$
152	$\pm 9$		<sup>6</sup> CASON	82	STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
186	$\pm 27$	11600	GIDAL	81	MRK2	$J/\psi$ decay
216	$\pm 13$		<sup>7</sup> CORDEN	79	OMEG	$12\text{--}15 \pi^- p \rightarrow n 2\pi^0$
190	$\pm 10$	10k	APEL	75	NICE	$40 \pi^- p \rightarrow n 2\pi^0$
192	$\pm 16$	4600	ENGLER	74	DBC	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
183	$\pm 15$	5300	FLATTE	71	HBC	$7 \pi^+ p \rightarrow \Delta^{++} f_2$
196	$\pm 30$		<sup>3</sup> STUNTEBECK	70	HBC	$8 \pi^- p, 5.4 \pi^+ d$
216	$\pm 20$	1960	<sup>3</sup> ARMENISE	68	DBC	$5.1 \pi^+ n \rightarrow p \pi^+ \text{MM}^-$
128	$\pm 27$		<sup>3</sup> BOESEBECK	68	HBC	$8 \pi^+ p$
176	$\pm 21$		<sup>3,8</sup> JOHNSON	68	HBC	$3.7\text{--}4.2 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

168	$\pm 7$		<sup>9</sup> KLEMPPT	22	RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0,$ $\gamma K_S^0 K_S^0$
183	$\pm 2$		CARVER	21	CLAS	$\gamma p \rightarrow \pi^0 \pi^0 p$
195	$\pm 15$	870	<sup>10</sup> SCHEGELSKY	06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$
121	$\pm 26$		TIKHOMIROV	03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
187	$\pm 20$		<sup>11</sup> ALDE	97	GAM2	$450 p p \rightarrow p p \pi^0 \pi^0$
184	$\pm 10$		<sup>11</sup> GRYGOREV	96	SPEC	$40 \pi^- N \rightarrow K_S^0 K_S^0 X$
200	$\pm 10$		AKER	91	CBAR	$0.0 \bar{p} p \rightarrow 3\pi^0$
240	$\pm 40$	3k	BINON	83	GAM2	$38 \pi^- p \rightarrow n 2\eta$
187	$\pm 30$	650	<sup>3</sup> ANTIPOV	77	CIBS	$25 \pi^- p \rightarrow p 3\pi$
225	$\pm 38$	16000	DEUTSCH...	76	HBC	$16 \pi^+ p$
166	$\pm 28$	600	<sup>3</sup> TAKAHASHI	72	HBC	$8 \pi^- p \rightarrow n 2\pi$
173	$\pm 53$		<sup>3</sup> ARMENISE	70	HBC	$9 \pi^+ n \rightarrow p \pi^+ \pi^-$

<sup>1</sup> Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.

<sup>2</sup> Breit-Wigner width

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>4</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>5</sup> From an energy-independent partial-wave analysis.

<sup>6</sup> From an amplitude analysis of the reaction  $\pi^+ \pi^- \rightarrow 2\pi^0$ .


<sup>7</sup> From an amplitude analysis of  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$  scattering data.

<sup>8</sup> JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

<sup>9</sup> Fit of the tensor partial waves from BES3 in the multipole basis.

<sup>10</sup> From analysis of L3 data at 91 and 183–209 GeV.

<sup>11</sup> Systematic uncertainties not estimated.

 $f_2(1270)$  width (MeV) **$f_2(1270)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1 \pi\pi$	$(84.3 \pm 2.8) \%$	S=1.2
$\Gamma_2 \pi^+\pi^- 2\pi^0$	$(7.7 \pm 1.2) \%$	S=1.2
$\Gamma_3 K\bar{K}$	$(4.6 \pm 0.4) \%$	S=2.7
$\Gamma_4 2\pi^+ 2\pi^-$	$(2.8 \pm 0.4) \%$	S=1.2
$\Gamma_5 \eta\eta$	$(4.0 \pm 0.8) \times 10^{-3}$	S=2.1
$\Gamma_6 4\pi^0$	$(3.0 \pm 1.0) \times 10^{-3}$	
$\Gamma_7 \gamma\gamma$	$(1.42 \pm 0.24) \times 10^{-5}$	S=1.4

$\Gamma_8$	$\eta\pi\pi$	< 8	$\times 10^{-3}$	CL=95%
$\Gamma_9$	$K^0 K^- \pi^+ + \text{c.c.}$	< 3.4	$\times 10^{-3}$	CL=95%
$\Gamma_{10}$	$e^+ e^-$	< 6	$\times 10^{-10}$	CL=90%

## CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 44 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 82.3$  for 37 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-91					
$x_3$	10	-39				
$x_4$	10	-38	1			
$x_5$	1	-6	0	0		
$x_6$	0	-7	0	0	0	
$x_7$	4	1	-15	0	0	0
$\Gamma$	-72	66	-10	-7	-1	0
	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
						$x_7$

	Mode	Rate (MeV)	Scale factor
$\Gamma_1$	$\pi\pi$	157.2	$+5.0$ $-1.0$
$\Gamma_2$	$\pi^+ \pi^- 2\pi^0$	14.3	$+2.3$ $-6.0$
$\Gamma_3$	$K\bar{K}$	8.5	$\pm 0.8$
$\Gamma_4$	$2\pi^+ 2\pi^-$	5.2	$\pm 0.7$
$\Gamma_5$	$\eta\eta$	0.75	$\pm 0.14$
$\Gamma_6$	$4\pi^0$	0.56	$\pm 0.19$
$\Gamma_7$	$\gamma\gamma$	0.0026 $\pm 0.0005$	1.4

## $f_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$	$\Gamma_1$
VALUE (MeV)	EVTS
<b>157.2<math>^{+5.0}_{-1.0}</math> OUR FIT</b>	
<b>157.0<math>^{+6.0}_{-1.0}</math></b>	<sup>1</sup> LONGACRE 86 MPS $22\pi^- p \rightarrow n2K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
152 $\pm 8$	<sup>2</sup> SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

### $\Gamma(K\bar{K})$

$\Gamma_3$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**8.5±0.8 OUR FIT** Error includes scale factor of 2.8.

**9.0<sup>+0.7</sup><sub>-0.3</sub>** <sup>1</sup> LONGACRE 86 MPS  $22 \pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.5 \pm 2.0$  870 <sup>2</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

### $\Gamma(\eta\eta)$

$\Gamma_5$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.75±0.14 OUR FIT** Error includes scale factor of 2.1.

**1.0 ±0.1** <sup>1</sup> LONGACRE 86 MPS  $22 \pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.8 \pm 0.4$  870 <sup>2</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

### $\Gamma(\gamma\gamma)$

$\Gamma_7$

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an S-wave contribution, values are systematically higher (typically around 3 keV).

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**2.6 ±0.5 OUR FIT** Error includes scale factor of 1.4.

**2.93±0.40** <sup>1</sup> DAI 14A RVUE Compilation

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.14 \pm 0.20$  2,3 PENNINGTON 08 RVUE Compilation

$3.82 \pm 0.30$  3,4 PENNINGTON 08 RVUE Compilation

$2.55 \pm 0.15$  870 <sup>5</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

$2.84 \pm 0.35$  BOGLIONE 99 RVUE  $\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$

$2.93 \pm 0.23 \pm 0.32$  <sup>6</sup> YABUKI 95 VNS

$2.58 \pm 0.13^{+0.36}_{-0.27}$  <sup>7</sup> BEHREND 92 CELL  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$3.10 \pm 0.35 \pm 0.35$  <sup>8</sup> BLINOV 92 MD1  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$2.27 \pm 0.47 \pm 0.11$  ADACHI 90D TOPZ  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$3.15 \pm 0.04 \pm 0.39$  BOYER 90 MRK2  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$3.19 \pm 0.16^{+0.29}_{-0.28}$  MARSISKE 90 CBAL  $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

$2.35 \pm 0.65$  <sup>9</sup> MORGAN 90 RVUE  $\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$

$3.19 \pm 0.09^{+0.22}_{-0.38}$  2177 OEST 90 JADE  $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

$3.2 \pm 0.1 \pm 0.4$  <sup>10</sup> AIHARA 86B TPC  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$2.5 \pm 0.1 \pm 0.5$  BEHREND 84B CELL  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$2.85 \pm 0.25 \pm 0.5$  <sup>11</sup> BERGER 84 PLUT  $e^+ e^- \rightarrow e^+ e^- 2\pi$

$2.70 \pm 0.05 \pm 0.20$  COURAU 84 DLCO  $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

$2.52 \pm 0.13 \pm 0.38$	<sup>12</sup> SMITH EDWARDS	84C 82F	MRK2 CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ $e^+ e^- \rightarrow e^+ e^- 2\pi^0$
$2.7 \pm 0.2 \pm 0.6$	<sup>13</sup> EDWARDS	82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
$2.9^{+0.6}_{-0.4} \pm 0.6$	BRANDELIK	81B	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
$3.2 \pm 0.2 \pm 0.6$	ROUSSARIE	81	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
$3.6 \pm 0.3 \pm 0.5$	<sup>14</sup> BERGER	80B	PLUT	$e^+ e^-$

<sup>1</sup> Based on a  $K$ -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.

<sup>2</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).

<sup>3</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

<sup>4</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

<sup>5</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

<sup>6</sup> With a narrow scalar state around 1220 MeV.

<sup>7</sup> Using a unitarized model with a 300 – 500 keV wide scalar at 1100 MeV.

<sup>8</sup> Using the unitarized model of LYTH 85.

<sup>9</sup> Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with  $\gamma\gamma$  width of  $f_0(1370)$ :  $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$  KeV.

<sup>10</sup> Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes  $2.66 \pm 0.21$  in the calculation of LANDRO 86.

<sup>11</sup> Using the MENNESSIER 83 model.

<sup>12</sup> Superseded by BOYER 90.

<sup>13</sup> If helicity = 2 assumption is not made.

<sup>14</sup> Using mass, width and  $B(f_2(1270) \rightarrow 2\pi)$  from PDG 78.

$\Gamma(e^+ e^-)$					$\Gamma_{10}$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.11	90	ACHASOV	00K	SND	$e^+ e^- \rightarrow \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.7	90	VOROBIEV	88	ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

### $f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_7/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
<b>0.121 ± 0.020 OUR FIT</b>	Error includes scale factor of 1.3.				
<b>0.091 ± 0.007 ± 0.027</b>					
1 ALBRECHT	90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.104 ± 0.007 ± 0.072	2 ALBRECHT	90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	

<sup>1</sup> Using an incoherent background.

<sup>2</sup> Using a coherent background.

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_5\Gamma_7/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
<b>11.5<sup>+1.8+4.5</sup><sub>-2.0-3.7</sub></b>	1 UEHARA	10A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$	

<sup>1</sup> Including interference with the  $f'_2(1525)$  (parameters fixed to the values from the 2008 edition of this review, PDG 08) and  $f_0(Y)$ .

**Helicity-0/Helicity-2 RATIO IN  $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$** 

<u>VALUE</u> (units $10^{-2}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.7 \pm 0.3^{+15.9}_{-2.9}</math></b>	UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
9.5 $\pm 1.8$	<sup>1</sup> DAI	14A RVUE	Compilation
13	2,3 PENNINGTON 08	RVUE	Compilation
26	3,4 PENNINGTON 08	RVUE	Compilation

<sup>1</sup> Based on a  $K$ -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.  
<sup>2</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).  
<sup>3</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.  
<sup>4</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

 **$f_2(1270)$  BRANCHING RATIOS**

<u><math>\Gamma(\pi\pi)/\Gamma_{\text{total}}</math></u>	<u><math>\Gamma_1/\Gamma</math></u>			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.843^{+0.028}_{-0.010}</math> OUR FIT</b>	Error includes scale factor of 1.2.			
<b><math>0.837 \pm 0.020</math> OUR AVERAGE</b>				
0.849 $\pm 0.025$	CHABAUD	83 ASPK	17 $\pi^- p$ polarized	
0.85 $\pm 0.05$	250 BEAUPRE	71 HBC	$8 \pi^+ p \rightarrow \Delta^{++} f_2$	
0.8 $\pm 0.04$	600 OH	70 HBC	$1.26 \pi^- p \rightarrow \pi^+ \pi^- n$	
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
0.856 $\pm 0.001 \pm 0.05$	<sup>1</sup> ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$	

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

<u><math>\Gamma(\pi^+\pi^- 2\pi^0)/\Gamma(\pi\pi)</math></u>	<u><math>\Gamma_2/\Gamma_1</math></u>			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.091^{+0.015}_{-0.040}</math> OUR FIT</b>	Error includes scale factor of 1.2.			
<b><math>0.15 \pm 0.06</math></b>	600 EISENBERG	74 HBC	$4.9 \pi^+ p \rightarrow \Delta^{++} f_2$	
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
0.07	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$	

<u><math>\Gamma(K\bar{K})/\Gamma_{\text{total}}</math></u>	<u><math>\Gamma_3/\Gamma</math></u>		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
0.033 $\pm 0.001 \pm 0.005$	<sup>1</sup> ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ )).

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$  $\Gamma_3/\Gamma_1$ 

We average only experiments which either take into account  $f_2(1270)$ - $a_2(1320)$  interference explicitly or demonstrate that  $a_2(1320)$  production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.054<sup>+0.005</sup><sub>-0.006</sub> OUR FIT</b>				Error includes scale factor of 2.7.

**0.041<sup>+0.004</sup><sub>-0.005</sub> OUR AVERAGE**

0.045 $\pm$ 0.01		<sup>1</sup> BARGIOTTI	03	OBLX $\bar{p}p$
0.037 $^{+0.008}$ $^{-0.021}$		ETKIN	82B	MPS $23 \pi^- p \rightarrow n2K_S^0$
0.045 $\pm$ 0.009		CHABAUD	81	ASPK $17 \pi^- p$ polarized
0.039 $\pm$ 0.008		LOVERRE	80	HBC $4 \pi^- p \rightarrow K\bar{K}N$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
0.052 $\pm$ 0.025		ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
0.036 $\pm$ 0.005		<sup>2</sup> COSTA	80	OMEG $1-2.2 \pi^- p \rightarrow K^+ K^- n$
0.030 $\pm$ 0.005		<sup>3</sup> MARTIN	79	RVUE
0.027 $\pm$ 0.009		<sup>4</sup> POLYCHRO...	79	STRC $7 \pi^- p \rightarrow n2K_S^0$
0.025 $\pm$ 0.015		EMMS	75D	DBC $4 \pi^+ n \rightarrow pf_2$
0.031 $\pm$ 0.012	20	ADERHOLZ	69	HBC $8 \pi^+ p \rightarrow K^+ K^- \pi^+ p$

<sup>1</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .

<sup>2</sup> Re-evaluated by CHABAUD 83.

<sup>3</sup> Includes PAWLICKI 77 data.

<sup>4</sup> Takes into account the  $f_2(1270)$ - $f'_2(1525)$  interference.

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$  $\Gamma_4/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.033<math>\pm</math>0.005 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.033<math>\pm</math>0.004 OUR AVERAGE</b>				Error includes scale factor of 1.1.
0.024 $\pm$ 0.006	160	EMMS	75D	DBC $4 \pi^+ n \rightarrow pf_2$
0.051 $\pm$ 0.025	70	EISENBERG	74	HBC $4.9 \pi^+ p \rightarrow \Delta^{++} f_2$
0.043 $^{+0.007}$ $^{-0.011}$	285	<sup>1</sup> LOUIE	74	HBC $3.9 \pi^- p \rightarrow nf_2$
0.037 $\pm$ 0.007	154	ANDERSON	73	DBC $6 \pi^+ n \rightarrow pf_2$
0.047 $\pm$ 0.013		OH	70	HBC $1.26 \pi^- p \rightarrow \pi^+ \pi^- n$

<sup>1</sup> LOUIE 74 was quoted as 0.065 in PDG 74. Factor 2/3 to go from  $\pi^+ \pi^- \rightarrow \pi\pi$  forgotten. Mike L.

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma_1$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.0<math>\pm</math>0.8 OUR FIT</b>			Error includes scale factor of 2.1.
<b>2.9<math>\pm</math>0.5 OUR AVERAGE</b>			
2.7 $\pm$ 0.7		BINON	05 GAMS $33 \pi^- p \rightarrow \eta\eta n$
2.8 $\pm$ 0.7		ALDE	86D GAM4 $100 \pi^- p \rightarrow 2\eta n$
5.2 $\pm$ 1.7		BINON	83 GAM2 $38 \pi^- p \rightarrow 2\eta n$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			

4.0 $\pm$ 1.0 $\pm$ 2.0	<sup>1</sup> ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
-------------------------	-----------------------	---------	--

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

$\Gamma(\eta\eta)/\Gamma(\pi\pi)$				$\Gamma_5/\Gamma_1$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.003±0.001</b>		BARBERIS 00E	450	$p p \rightarrow p_f \eta\eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.05	95	EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\eta$
<0.016	95	EMMS 75D	DBC	$4\pi^+ n \rightarrow p f_2$
<0.09	95	EISENBERG 74	HBC	$4.9\pi^+ p \rightarrow \Delta^{++} f_2$
$\Gamma(4\pi^0)/\Gamma_{\text{total}}$				$\Gamma_6/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0030±0.0010 OUR FIT</b>				
<b>0.003 ± 0.001</b>	400 ± 50	ALDE 87	GAM4	$100\pi^- p \rightarrow 4\pi^0 n$
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_7/\Gamma$
<u>VALUE (units <math>10^{-5}</math>)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.57±0.01 <sup>+1.39</sup> <sub>-0.14</sub>		UEHARA 08A	BELL	$10.6e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$				$\Gamma_8/\Gamma_1$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.010</b>	95	EMMS 75D	DBC	$4\pi^+ n \rightarrow p f_2$
$\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$				$\Gamma_9/\Gamma_1$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.004</b>	95	EMMS 75D	DBC	$4\pi^+ n \rightarrow p f_2$
$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{10}/\Gamma$
<u>VALUE (units <math>10^{-10}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;6</b>	90	ACHASOV 00K	SND	$e^+ e^- \rightarrow \pi^0 \pi^0$

## f<sub>2</sub>(1270) REFERENCES

KLEMPT	22	PL B830 137171	E. Klempert <i>et al.</i>	(BONN)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
CARVER	21	PRL 126 082002	M. Carver <i>et al.</i>	(CLAS Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
BOGOLYUB...	13	PAN 76 1324	M.Yu. Bogolyubsky <i>et al.</i>	(HYPERON-M Collab.)
		Translated from YAF 76 1389.		
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		

ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington	
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)
		Translated from YAF 59 2187.		
YABUKI	95	JPSJ 64 435	F. Yabuki <i>et al.</i>	(VENUS Collab.)
PROKOSHKIN	94	PD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)
		Translated from DANS 336 613.		
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)
BLINOV	92	ZPHY C53 33	A.E. Blinov <i>et al.</i>	(NOVO)
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)
ADACHI	90D	PL B234 185	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48 436.		
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAZO, CLER, FRAS+)
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
LYTH	85	JP G11 459	D.H. Lyth	
BEHREND	84B	ZPHY C23 223	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BERGER	84	ZPHY C26 199	C. Berger <i>et al.</i>	(PLUTO Collab.)
COURAU	84	PL 147B 227	A. Courau <i>et al.</i>	(CIT, SLAC)
SMITH	84C	PR D30 851	J.R. Smith <i>et al.</i>	(SLAC, LBL, HARV)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
		Translated from YAF 38 934.		
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>	
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)

ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)
PDG	74	PL 50B 1	V. Chaloupka <i>et al.</i>	
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)