

$f_1(1285)$

$I^G(J^{PC}) = 0^+(1^{++})$

$f_1(1285)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1281.8 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
1280.2 ± 0.6 ± 1.2	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1281.0 ± 0.8		DICKSON	16 CLAS	$\gamma p \rightarrow \eta \pi^+ \pi^- p$
1287.4 ± 3.0	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
1281.16 ± 0.39 ± 0.45		¹ LEES	12X BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$
1285.1 ± 1.0 ± 1.6		² ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
1281 ± 2 ± 1		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
1276.1 ± 8.1 ± 8.0	203	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
1274 ± 6	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1280 ± 4		ACCIARRI	01G L3	
1288 ± 4 ± 5	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1284 ± 6	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1281 ± 1		BARBERIS	97B OMEG	$450 pp \rightarrow pp 2(\pi^+ \pi^-)$
1281 ± 1		BARBERIS	97C OMEG	$450 pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
1280 ± 2		³ ANTINORI	95 OMEG	$300,450 pp \rightarrow pp 2(\pi^+ \pi^-)$
1282.2 ± 1.5		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
1279 ± 5		FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1278 ± 2	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K\bar{K} \pi pp$
1278 ± 2		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi \pi p,$ $pp \rightarrow 4\pi pp$
1280.1 ± 2.1	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750	⁴ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1280 ± 1	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
1280 ± 4		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1277 ± 2	420	REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
1285 ± 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow N K\bar{K} \pi$
1279 ± 2	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K\bar{K} \pi \pi p,$ $pp \rightarrow K\bar{K} \pi pp$
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 pp
1278 ± 4		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
1283 ± 3	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K\bar{K} \pi n$
1282 ± 2	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K} 3\pi$

1279	± 5	210	GRASSLER	77	HBC	16 $\pi^\mp p$
1286	± 3	180	DUBOC	72	HBC	1.2 $\bar{p}p \rightarrow 2K4\pi$
1283	± 5		DAHL	67	HBC	1.6–4.2 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1289.3	± 2.8	234	ABLIKIM	19BA	BES3	$e^+ e^- \rightarrow \psi(2S)$
1284.2	± 2.2		⁵ AAIJ	14Y	LHCb	$\overline{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
1281.9	± 0.5		⁵ SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
1282.8	± 0.6		⁵ SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1270	± 10		AMELIN	95	VES	$37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280	± 2		ABATZIS	94	OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$
1282	± 4		ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270	± 6	± 10	ARMSTRONG	92C	OMEG	300 $p p \rightarrow p p \pi^+ \pi^- \gamma$
1281	± 1		ARMSTRONG	89E	OMEG	300 $p p \rightarrow p p 2(\pi^+ \pi^-)$
1279	± 6	± 10	BECKER	87	MRK3	$e^+ e^- \rightarrow \phi K \overline{K} \pi$
1286	± 9		GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
1287	± 5	353	BITYUKOV	84B	SPEC	32 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
~ 1279			⁶ TORNQVIST	82B	RVUE	
1275	± 6	31	BROMBERG	80	SPEC	100 $\pi^- p \rightarrow K \overline{K} \pi X$
1288	± 9	200	GURTU	79	HBC	4.2 $K^- p \rightarrow n \eta 2\pi$
~ 1275.0		46	⁷ STANTON	79	CNTR	8.5 $\pi^- p \rightarrow n 2\gamma 2\pi$
1271	± 10	34	CORDEN	78	OMEG	12–15 $\pi^- p \rightarrow K^+ K^- \pi n$
1295	± 12	85	CORDEN	78	OMEG	12–15 $\pi^- p \rightarrow n 5\pi$
1292	± 10	150	DEFOIX	72	HBC	0.7 $\bar{p}p \rightarrow 7\pi$
1280	± 3	500	⁸ THUN	72	MMS	13.4 $\pi^- p$
1303	± 8		BARDADIN...	71	HBC	8 $\pi^+ p \rightarrow p 6\pi$
1283	± 6		BOESEBECK	71	HBC	16.0 $\pi p \rightarrow p 5\pi$
1270	± 10		CAMPBELL	69	DBC	2.7 $\pi^+ d$
1285	± 7		LORSTAD	69	HBC	0.7 $\bar{p}p$, 4,5-body
1290	± 7		D'ANDLAU	68	HBC	1.2 $\bar{p}p$, 5–6 body

¹ Using the $2\pi^+ 2\pi^-$ and $\pi^+ \pi^- \eta$ modes of $f_1(1285)$ decay.

² The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

³ Supersedes ABATZIS 94, ARMSTRONG 89E.


⁴ From partial wave analysis of $K^+ \overline{K}^0 \pi^-$ system.

⁵ No systematic error given.

⁶ From a unitarized quark-model calculation.

⁷ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁸ Seen in the missing mass spectrum.

 $f_1(1285)$ mass (MeV) **$f_1(1285)$ WIDTH**

Only experiments giving width error less than 20 MeV are kept for averaging.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
23.0± 1.1 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.			
28.2± 1.1 ^{+ 5.5} _{- 2.9}	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
18.4± 1.4		DICKSON	16 CLAS	2.55 $\gamma p \rightarrow \eta \pi^+ \pi^- p$
18.3± 6.3	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
22.0± 3.1 ^{+ 2.0} _{- 1.5}		¹ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$

35 \pm 6 \pm 4		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
40.0 \pm 8.6 \pm 9.3	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
29 \pm 12	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
45 \pm 9 \pm 7	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
55 \pm 18	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta\pi^0\pi^0 n$
24 \pm 3		BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$
20 \pm 2		BARBERIS	97C OMEG	$450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
36 \pm 5		² ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+\pi^-)$
29.0 \pm 4.1		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
25 \pm 4	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K\bar{K}\pi pp$
22 \pm 2	4750	³ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 \pm 4	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
19 \pm 5		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta\pi^+\pi^- n$
32 \pm 8	420	REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
22 \pm 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow N K \bar{K} \pi$
32 \pm 3	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K\bar{K}\pi\pi p, p\bar{p} \rightarrow K\bar{K}\pi pp$
24 \pm 3		CHAUVAT	84 SPEC	ISR 31.5 pp
29 \pm 10	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K\bar{K}\pi n$
28.3 \pm 6.7	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
17.1 \pm 3.4	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
32.4 \pm 5.8		⁴ AAIJ	14Y LHCb	$\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+\pi^-)$
18.2 \pm 1.2		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$
19.4 \pm 1.5		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
40 \pm 5		ABATZIS	94 OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$
31 \pm 5		ARMSTRONG	89E OMEG	$300 pp \rightarrow pp2(\pi^+\pi^-)$
41 \pm 12		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$
17.9 \pm 10.9	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
14 \pm 20 \pm 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K\bar{K}\pi$
26 \pm 12		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
25 \pm 15	200	GURTU	79 HBC	$4.2 K^- p \rightarrow n\eta 2\pi$
~ 10		⁵ STANTON	79 CNTR	$8.5 \pi^- p \rightarrow n 2\gamma 2\pi$
24 \pm 18	210	GRASSLER	77 HBC	$16 \pi^\mp p$
28 \pm 5	150	⁶ DEFOIX	72 HBC	$0.7 \bar{p}p \rightarrow 7\pi$
46 \pm 9	180	⁶ DUBOC	72 HBC	$1.2 \bar{p}p \rightarrow 2K4\pi$
37 \pm 5	500	⁷ THUN	72 MMS	$13.4 \pi^- p$
10 \pm 10		BOESEBECK	71 HBC	$16.0 \pi p \rightarrow p5\pi$
30 \pm 15		CAMPBELL	69 DBC	$2.7 \pi^+ d$
60 \pm 15		⁶ LORSTAD	69 HBC	$0.7 \bar{p}p, 4,5\text{-body}$
35 \pm 10		⁶ DAHL	67 HBC	$1.6\text{--}4.2 \pi^- p$

¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.

² Supersedes ABATZIS 94, ARMSTRONG 89E.


³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

⁴ No systematic error given.

⁵ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁶ Resolution is not unfolded.

⁷ Seen in the missing mass spectrum.



$f_1(1285)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \quad 4\pi$	$(32.7 \pm 1.8) \%$	$S=1.2$
$\Gamma_2 \quad \pi^0 \pi^0 \pi^+ \pi^-$	$(21.8 \pm 1.2) \%$	$S=1.2$
$\Gamma_3 \quad 2\pi^+ 2\pi^-$	$(10.9 \pm 0.6) \%$	$S=1.2$
$\Gamma_4 \quad \rho^0 \pi^+ \pi^-$	$(10.9 \pm 0.6) \%$	$S=1.2$
$\Gamma_5 \quad \rho^0 \rho^0$	seen	
$\Gamma_6 \quad 4\pi^0$	$< 7 \times 10^{-4}$	$CL=90\%$
$\Gamma_7 \quad \eta \pi^+ \pi^-$	$(35 \pm 15) \%$	

Γ_8	$\eta\pi\pi$	(52.2 \pm 1.9) %	S=1.2
Γ_9	$a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$]	(38 \pm 4) %	
Γ_{10}	$\eta\pi\pi$ [excluding $a_0(980)\pi$]	(14 \pm 4) %	
Γ_{11}	$K\bar{K}\pi$	(9.0 \pm 0.4) %	S=1.1
Γ_{12}	$K\bar{K}^*(892)$	not seen	
Γ_{13}	$\pi^+\pi^-\pi^0$	(3.0 \pm 0.9) $\times 10^{-3}$	
Γ_{14}	$\rho^\pm\pi^\mp$	< 3.1 $\times 10^{-3}$	CL=95%
Γ_{15}	$\gamma\rho^0$	(6.1 \pm 1.0) %	S=1.7
Γ_{16}	$\phi\gamma$	(7.4 \pm 2.6) $\times 10^{-4}$	
Γ_{17}	e^+e^-	< 9.4 $\times 10^{-9}$	CL=90%
Γ_{18}	$\gamma\gamma^*$		
Γ_{19}	$\gamma\gamma$		

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.0$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	-29			
x_{10}	-12	-89		
x_{11}	22	-9	-4	
x_{15}	-24	-8	-3	-27
	x_1	x_9	x_{10}	x_{11}

$f_1(1285) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_{19}/\Gamma = (\Gamma_9 + \Gamma_{10})\Gamma_{19}/\Gamma$$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.62	95	GIDAL	87	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

$$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_{18}/\Gamma = (\Gamma_9 + \Gamma_{10})\Gamma_{18}/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.4 \pm 0.4 OUR AVERAGE		Error includes scale factor of 1.4.		
1.18 \pm 0.25 \pm 0.20	26	1, ² AIHARA	88B TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

2.30 \pm 0.61 \pm 0.42	1, ³ GIDAL	87	MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
----------------------------	-----------------------	----	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 \pm 0.3 \pm 0.3	420	⁴ ACHARD	02B L3	183–209 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
-------------------------	-----	---------------------	--------	---

¹ Assuming a ρ -pole form factor.

² Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

³ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.⁴ Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52. **$f_1(1285)$ BRANCHING RATIOS**

$$\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$$

$$\Gamma_{11}/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.274±0.017 OUR FIT	Error includes scale factor of 1.4.		
0.271±0.016 OUR AVERAGE	Error includes scale factor of 1.2.		
0.265±0.014	1 BARBERIS 97C OMEG 450 $p p \rightarrow p p K_S^0 K^\pm \pi^\mp$		
0.28 ± 0.05	2 ARMSTRONG 89E OMEG 300 $p p \rightarrow p p f_1(1285)$		
0.37 ± 0.03 ± 0.05	3 ARMSTRONG 89G OMEG 85 $\pi p \rightarrow 4\pi X$		

¹ Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.³ 4π consistent with being entirely $\rho\pi\pi$.

$$\Gamma(\pi^0\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma = \frac{2}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
0.218±0.012 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
0.109±0.006 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_4/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
0.109±0.006 OUR FIT	Error includes scale factor of 1.2.

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2\pi^+ 2\pi^-)$$

$$\Gamma_4/\Gamma_3$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0±0.4	GRASSLER 77 HBC	16 GeV	$\pi^\pm p$

$$\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	COMMENT
seen	BARBERIS 00C	$450 p p \rightarrow p_f 4\pi p_s$

$$\Gamma(4\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma$$

VALUE (units 10^{-4})	CL %	DOCUMENT ID	TECN	COMMENT
<7	90	ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\eta\pi^+\pi^-)$$

$$\Gamma_{13}/\Gamma_7$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.16±0.20	2.3k	¹ DOROFEEV 11 VES		$\pi^- N \rightarrow \pi^- f_1(1285) N$

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum.

$$\Gamma(\eta\pi\pi)/\Gamma_{\text{total}}$$

$$\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$$

VALUE	DOCUMENT ID
0.522±0.019 OUR FIT	Error includes scale factor of 1.2.

$\Gamma(4\pi)/\Gamma(\eta\pi\pi)$	$\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9+\Gamma_{10})$			
VALUE	DOCUMENT ID	TECN	COMMENT	
0.63±0.06 OUR FIT	Error includes scale factor of 1.2.			
0.41±0.14 OUR AVERAGE				
0.37±0.11±0.11	BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.64±0.40	GURTU	79	HBC	$4.2 K^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.93±0.30	¹ GRASSLER	77	HBC	$16 \pi^\mp p$

¹ Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(\eta\pi\pi)$	Γ_3/Γ_8			
VALUE	DOCUMENT ID	TECN	COMMENT	
0.28±0.02±0.02	¹ LEES	12X	BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$
¹ Assuming $B(f_1(1285) \rightarrow \pi\pi\eta) = 3/2 B(f_1(1285) \rightarrow \pi^+ \pi^- \eta)$.				
$\Gamma(a_0(980)\pi \text{ [ignoring } a_0(980) \rightarrow K\bar{K}])/\Gamma(\eta\pi\pi)$	$\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9+\Gamma_{10})$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.72±0.08 OUR FIT				
0.72±0.07 OUR AVERAGE				
0.74±0.02±0.09	DICKSON	16	CLAS	$\gamma p \rightarrow f_1(1285) p$
0.72±0.15	GURTU	79	HBC	$4.2 K^- p$
$0.6^{+0.3}_{-0.2}$	CORDEN	78	OMEG	$12-15 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.69	95	ACHARD	02B L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
0.28±0.07	ALDE	97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1.0 ± 0.3	GRASSLER	77	HBC	$16 \pi^\mp p$

$\Gamma(K\bar{K}\pi)/\Gamma(\eta\pi\pi)$	$\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9+\Gamma_{10})$			
VALUE	DOCUMENT ID	TECN	COMMENT	
0.172±0.011 OUR FIT				
0.176±0.012 OUR AVERAGE				
0.216±0.010±0.031	DICKSON	16	CLAS	$\gamma p \rightarrow f_1(1285) p$
0.166±0.01 ± 0.008	BARBERIS	98C	OMEG	$450 pp \rightarrow p_f f_1(1285) p_s$
0.42 ± 0.15	GURTU	79	HBC	$4.2 K^- p$
0.5 ± 0.2	¹ CORDEN	78	OMEG	$12-15 \pi^- p$
0.20 ± 0.08	² DEFOIX	72	HBC	$0.7 \bar{p}p \rightarrow 7\pi$
0.16 ± 0.08	CAMPBELL	69	DBC	$2.7 \pi^+ d$

¹ CORDEN 78 assumes low-mass $\eta\pi\pi$ region is dominantly 1^{++} . See BARBERIS 98C and MANAK 00A for discussion.

² $K\bar{K}$ system characterized by the $I = 1$ threshold enhancement. (See under $a_0(980)$).

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_{12}/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	NACASCH	78	HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	¹ ACHARD	07	L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$

¹ A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_{13}/Γ			
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.30±0.055±0.074	2.3k	1 DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

$\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$	Γ_{14}/Γ			
<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.31	95	DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$	Γ_{15}/Γ			
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

6.1±1.0 OUR FIT Error includes scale factor of 1.7.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.8±0.7±0.6	1 AMELIN	95 VES	37 $\pi^- N \rightarrow \pi^-\pi^+\pi^-\gamma N$
<5	95 BITYUKOV	91B SPEC	32 $\pi^- p \rightarrow \pi^+\pi^-\gamma n$

¹ Not an independent measurement.

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$	$\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

0.55±0.10 OUR FIT Error includes scale factor of 1.5.

0.45±0.18	1 COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
------------------	-----------	---------	---

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.


$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$	$\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

8.6±1.6 OUR FIT Error includes scale factor of 1.9.

8.5±2.0 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

21.3±4.4	DICKSON	16 CLAS	$\gamma p \rightarrow f_1(1285) p$
10.0±1.0±2.0	BARBERIS	98C OMEG	$450 pp \rightarrow p_f f_1(1285) p_s$
7.5±1.0	¹ ARMSTRONG	92C OMEG	$300 pp \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$

¹ Published value multiplied by 1.5.

 $\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$ Γ_{15}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.035	90	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma K\bar{K}\pi) = < 0.72 \times 10^{-3}$.				

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{16}/Γ_{11}

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.82±0.21±0.20		19	BITYUKOV	88	SPEC $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.50	95		BARBERIS	98C	OMEG $450 pp \rightarrow p_f f_1(1285) p_s$
<0.93	95		AMELIN	95	VES $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.4 \times 10^{-9}$	90	¹ ACHASOV	20	$e^+ e^- \rightarrow \eta\pi^0\pi^0$

¹ ACHASOV 20 reports two candidate events corresponding to a significance of 2.5σ and the branching fraction of $(5.1^{+3.7}_{-2.7}) \times 10^{-9}$.

$f_1(1285)$ REFERENCES

ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	20	PL B800 135074	M.N. Achasov <i>et al.</i>	(SND Collab.)
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DICKSON	16	PR C93 065202	R. Dickson <i>et al.</i>	(JLab CLAS Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	14Y	PRL 112 091802	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
SOSA	99	PRL 83 913	M. Sosa <i>et al.</i>	
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 60	458.	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BITYUKOV	91B	SJNP 54 318	S.I. Bityukov <i>et al.</i>	(SERP)
		Translated from YAF 54	529.	
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+) JPC
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
AIHARA	88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
BITYUKOV	88	PL B203 327	S.I. Bityukov <i>et al.</i>	(SERP)
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)
Conference				
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIR, SAGA+) IJP
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP
ARMSTRONG	84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP
BITYUKOV	84B	PL 144B 133	S.I. Bityukov <i>et al.</i>	(SERP)
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
NACASCH	78	NP B135 203	R. Nacasch <i>et al.</i>	(PARIS, MADR, CERN)
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LIVP)
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)
BARDADIN-...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(VARS)
BOESEBECK	71	PL 34B 659	K. Boesebeck	(AACH, BERL, BONN, CERN, CRAC+)

CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+) IJP
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP
