

**$\Sigma(1775) \text{ } 5/2^-$**  $I(J^P) = 1(\frac{5}{2}^-)$  Status: \*\*\*

Discovered by GALTIERI 63, this resonance plays the same role as cornerstone for isospin-1 analyses in this region as the  $\Lambda(1820)F_{05}$  does in the isospin-0 channel.

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

 **$\Sigma(1775) \text{ POLE POSITION}$** **REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1760 to 1780 (<math>\approx 1770</math>) OUR ESTIMATE</b>			
1767 $\pm 4$	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
1767 $^{+2}_{-2}$	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1759	ZHANG 13A	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**-2xIMAGINARY PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>45 to 65 (<math>\approx 55</math>) OUR ESTIMATE</b>			
122 $\pm 8$	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
128 $^{+4}_{-2}$	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118	ZHANG 13A	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

 **$\Sigma(1775) \text{ POLE RESIDUES}$** 

The normalized residue is the residue divided by  $\Gamma_{pole}/2$ .

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}$** 

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.44 $\pm 0.09$	$-17 \pm 10$	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.371	-32	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma\pi$** 

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
0.13 $\pm 0.03$	$10 \pm 12$	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.115	-24	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda\pi$** 

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.47 ± 0.10	130 ± 15	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.325	157	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$ , *D*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.391	137	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$ , *G*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0129	-58	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892)$ ,  $S=1/2$ , *D*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.04 ± 0.02	-100 ± 60	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892)$ ,  $S=3/2$ , *D*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.09 ± 0.06	10 ± 50	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892)$ ,  $S=3/2$ , *G*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.04 ± 0.02	-100 ± 60	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Xi K$** 

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 ± 0.01	-90 ± 35	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520)\pi$ , *P*-wave**

<u>MODULUS</u>	<u>PHASE (°)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.09 ± 0.03	10 ± 30	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520)\pi$ , *F*-wave**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.01 ± 0.01	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Delta\bar{K}$ , *D*-wave**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 ± 0.02	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

## $\Sigma(1775)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1770 to 1780 (<math>\approx 1775</math>) OUR ESTIMATE</b>			
1776 $\pm$ 4	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
1778 $\pm$ 1	ZHANG 13A	DPWA	$\bar{K}N$ multichannel
1778 $\pm$ 5	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1777 $\pm$ 5	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1775 $\pm$ 10	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
1774 $\pm$ 10	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
1772 $\pm$ 6	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1774 $\pm$ 5	GOPAL 77	DPWA	$\bar{K}N$ multichannel
1772 or 1777	<sup>1</sup> MARTIN 77	DPWA	$\bar{K}N$ multichannel
1765	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$
1 The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.			

## $\Sigma(1775)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>105 to 135 (<math>\approx 120</math>) OUR ESTIMATE</b>			
124 $\pm$ 8	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
131 $\pm$ 3	ZHANG 13A	DPWA	$\bar{K}N$ multichannel
137 $\pm$ 10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
116 $\pm$ 10	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
125 $\pm$ 15	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
146 $\pm$ 18	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
154 $\pm$ 10	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
130 $\pm$ 10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
102 or 103	<sup>1</sup> MARTIN 77	DPWA	$\bar{K}N$ multichannel
120	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$
1 The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.			

## $\Sigma(1775)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 N\bar{K}$	37–43%
$\Gamma_2 \Lambda\pi$	14–20%
$\Gamma_3 \Sigma\pi$	2–5%
$\Gamma_4 \Sigma(1385)\pi$	8–12%
$\Gamma_5 \Sigma(1385)\pi$ , <i>D</i> -wave	
$\Gamma_6 \Sigma(1385)\pi$ , <i>G</i> -wave	
$\Gamma_7 \Lambda(1520)\pi$ , <i>P</i> -wave	17–23%

$\Gamma_8$	$\Sigma \pi \pi$
$\Gamma_9$	$\Delta(1232) \bar{K}$ , <i>D</i> -wave
$\Gamma_{10}$	$N \bar{K}^*(892)$ , $S=1/2$
$\Gamma_{11}$	$N \bar{K}^*(892)$ , $S=1/2$ , <i>D</i> -wave
$\Gamma_{12}$	$N \bar{K}^*(892)$ , $S=3/2$ , <i>D</i> -wave
$\Gamma_{13}$	$N \bar{K}^*(892)$ , $S=3/2$ , <i>G</i> -wave

## $\Sigma(1775)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on  $\Lambda$  and  $\Sigma$  Resonances. Also, the errors quoted do not include uncertainties due to the parametrization used in the partial-wave analyses and are thus too small.

### $\Gamma(N\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>0.37 to 0.43 OUR ESTIMATE</b>				
0.43 $\pm$ 0.09	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel	
0.40 $\pm$ 0.01	ZHANG 13A	DPWA	$\bar{K}N$ multichannel	
0.40 $\pm$ 0.02	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.37 $\pm$ 0.03	ALSTON-...	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.402	<sup>1</sup> KAMANO 15	DPWA	Multichannel	
0.41 $\pm$ 0.03	GOPAL 77	DPWA	See GOPAL 80	
0.37 or 0.36	<sup>2</sup> MARTIN 77	DPWA	$\bar{K}N$ multichannel	

<sup>1</sup> From the preferred solution A in KAMANO 15.

<sup>2</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

### $\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
0.49 $\pm$ 0.10	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.244	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel	

<sup>1</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Lambda\pi)/\Gamma(N\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_1$
0.33 $\pm$ 0.05	UHLIG 67	HBC	$K^- p$ 0.9 GeV/c	

### $\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
0.035 $\pm$ 0.010	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.042	<sup>1</sup> KAMANO 15	DPWA	$\bar{K}N$ multichannel	

<sup>1</sup> From the preferred solution A in KAMANO 15.

### $\Gamma(\Sigma(1385)\pi)/\Gamma(N\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma_1$
0.25 $\pm$ 0.09	UHLIG 67	HBC	$K^- p$ 0.9 GeV/c	

$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
0.309	<sup>1</sup> KAMANO	15	DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

 $\Gamma(\Sigma(1385)\pi, G\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	<sup>1</sup> KAMANO	15	DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

 $\Gamma(\Lambda(1520)\pi, P\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02±0.01	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

 $\Gamma(\Lambda(1520)\pi, P\text{-wave})/\Gamma(N\bar{K})$   $\Gamma_7/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.28±0.05	UHLIG	67	HBC $K^- p$ 0.9 GeV/c

 $\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			

0.12 <sup>1</sup> ARMENTEROS68C HDPC  $K^- N \rightarrow \Sigma\pi\pi$

<sup>1</sup> For about 3/4 of this, the  $\Sigma\pi$  system has  $I = 0$  and is almost entirely  $\Lambda(1520)$ . For the rest, the  $\Sigma\pi$  has  $I = 1$ , which is about what is expected from the known  $\Sigma(1775) \rightarrow \Sigma(1385)\pi$  rate, as seen in  $\Lambda\pi\pi$ .

 $\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			

not seen <sup>1</sup> KAMANO 15 DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			

0.003 <sup>1</sup> KAMANO 15 DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=3/2, G\text{-wave})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			

not seen <sup>1</sup> KAMANO 15 DPWA Multichannel

<sup>1</sup> From the preferred solution A in KAMANO 15.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda\pi$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.31 ± 0.01	ZHANG	13A	DPWA Multichannel
-0.28 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel
-0.25 ± 0.02	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
-0.28 $^{+0.04}_{-0.05}$	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
-0.259 ± 0.048	DEVENISH	74B	Fixed- $t$ dispersion rel.
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.29 or -0.28	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
-0.30	DEBELLEFON	76	IPWA $K^- p \rightarrow \Lambda\pi^0$

<sup>1</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma\pi$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.08 ± 0.01	ZHANG	13A	DPWA Multichannel
+0.13 ± 0.02	GOPAL	77	DPWA $\bar{K}N$ multichannel
0.09 ± 0.01	KANE	74	DPWA $K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.08 or +0.08	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel

<sup>1</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$ , <i>D</i> -wave	$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.12 ± 0.01	ZHANG	13A	DPWA Multichannel
-0.184 ± 0.011	<sup>1</sup> CAMERON	78	DPWA $K^- p \rightarrow \Sigma(1385)\pi$
+0.20 ± 0.02	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385)\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.32 ± 0.06	SIMS	68	DBC $K^- N \rightarrow \Lambda\pi\pi$
0.24 ± 0.03	ARMENTEROS67C	HBC	$K^- p \rightarrow \Lambda\pi\pi$

<sup>1</sup> The CAMERON 78 upper limit on *G*-wave decay is 0.03.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520)\pi$ , <i>P</i> -wave	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.06 ± 0.01	ZHANG	13A	DPWA Multichannel
-0.305 ± 0.010	<sup>1</sup> CAMERON	77	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$
0.31 ± 0.02	BARLETTA	72	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$
0.27 ± 0.03	ARMENTEROS65C	HBC	$K^- p \rightarrow \Lambda(1520)\pi^0$

<sup>1</sup> This rate combines *P*-wave- and *F*-wave decays. The CAMERON 77 results for the separate *P*-wave- and *F*-wave decays are  $-0.303 \pm 0.010$  and  $-0.037 \pm 0.014$ . The published signs have been changed here to be in accord with the baryon-first convention.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Delta(1232)\bar{K}$ , <i>D</i> -wave	$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.06 ± 0.03	ZHANG	13A	DPWA Multichannel

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892), S=1/2 \quad (\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.04 ± 0.01	ZHANG	13A	DPWA Multichannel

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892), S=3/2, D\text{-wave}$$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.04 ± 0.01	ZHANG	13A	DPWA Multichannel

## $\Sigma(1775)$ REFERENCES

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO	15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
ALSTON-...	78	PR D18 182	M. Alston-Garnjost <i>et al.</i>	(LBL, MTDO+) IJP
Also		PRL 38 1007	M. Alston-Garnjost <i>et al.</i>	(LBL, MTDO+) IJP
CAMERON	78	NP B143 189	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	B.R. Martin, M.K. Pidcock, R.G. Moorhouse	(LOUC+) IJP
Also		NP B126 266	B.R. Martin, M.K. Pidcock	(LOUC)
Also		NP B126 285	B.R. Martin, M.K. Pidcock	(LOUC) IJP
DEBELLEFON	76	NP B109 129	A. de Bellefon, A. Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	P.H. Baillon, P.J. Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145	A.J. van Horn	(LBL) IJP
Also		NP B87 157	A.J. van Horn	(LBL) IJP
DEVENISH	74B	NP B81 330	R.C.E. Devenish, C.D. Froggatt, B.R. Martin	(DESY+) IJP
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
PREVOST	74	NP B69 246	J. Prevost <i>et al.</i>	(SACL, CERN, HEID)
BARLETTA	72	NP B40 45	W.A. Barletta	(EFI) IJP
Also		PRL 17 841	S. Fenster <i>et al.</i>	(CHIC, ANL, CERN) IJP
ARMENTEROS	68C	NP B8 216	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL) I
SIMS	68	PRL 21 1413	W.H. Sims <i>et al.</i>	(FSU, TUFTS, BRAN)
ARMENTEROS	67C	ZPHY 202 486	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL)
UHLIG	67	PR 155 1448	R.P. Uhlig <i>et al.</i>	(UMD, NRL)
ARMENTEROS	65C	PL 19 338	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL) IJP
GALTIERI	63	PL 6 296	A. Galtieri, A. Hussain, R. Tripp	(LRL) IJ