

$\Upsilon(2S)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\Upsilon(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10023.4±0.5	¹ SHAMOV 23	RVUE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10022.7±0.4	² SHAMOV 23	RVUE	$e^+ e^- \rightarrow$ hadrons
10023.5±0.5	^{3,4} ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10023.6±0.5	^{5,6} BARU 86B	MD1	$e^+ e^- \rightarrow$ hadrons
10023.1±0.4	⁷ BARBER 84	ARG	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects. ² Obtained by reanalysing ARGUS and Crystal Ball data (BARBER 84), but not authored by the ARGUS and Crystal Ball collaboration. ³ Reanalysis of BARU 86B using new electron mass (COHEN 87). ⁴ Superseded by SHAMOV 23. ⁵ Reanalysis of ARTAMONOV 84. ⁶ Superseded by ARTAMONOV 00. ⁷ Reanalysed by SHAMOV 23.			

$m_{\Upsilon(3S)} - m_{\Upsilon(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

$\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
31.98±2.63 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

$\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \quad \Upsilon(1S)\pi^+\pi^-$	(17.85± 0.26) %	
$\Gamma_2 \quad \Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
$\Gamma_3 \quad \tau^+\tau^-$	(2.00± 0.21) %	
$\Gamma_4 \quad \mu^+\mu^-$	(1.93± 0.17) %	S=2.2
$\Gamma_5 \quad e^+e^-$	(1.91± 0.16) %	
$\Gamma_6 \quad \Upsilon(1S)\pi^0$	< 4 $\times 10^{-5}$	CL=90%
$\Gamma_7 \quad \Upsilon(1S)\eta$	(2.9 ± 0.4) $\times 10^{-4}$	S=2.0
$\Gamma_8 \quad J/\psi(1S)$ anything	< 6 $\times 10^{-3}$	CL=90%
$\Gamma_9 \quad J/\psi(1S)\eta_c$	< 5.4 $\times 10^{-6}$	CL=90%
$\Gamma_{10} \quad J/\psi(1S)\chi_{c0}$	< 3.4 $\times 10^{-6}$	CL=90%

Γ_{11}	$J/\psi(1S)\chi_{c1}$	< 1.2	$\times 10^{-6}$	CL=90%
Γ_{12}	$J/\psi(1S)\chi_{c2}$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{13}	$J/\psi(1S)\eta_c(2S)$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{14}	$J/\psi(1S)X(3940)$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{15}	$J/\psi(1S)X(4160)$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{16}	χ_{c1} anything	(2.2 \pm 0.5)	$\times 10^{-4}$	
Γ_{17}	$\chi_{c1}(1P)^0 X_{tetra}$	< 3.67	$\times 10^{-5}$	CL=90%
Γ_{18}	χ_{c2} anything	(2.3 \pm 0.8)	$\times 10^{-4}$	
Γ_{19}	$\psi(2S)\eta_c$	< 5.1	$\times 10^{-6}$	CL=90%
Γ_{20}	$\psi(2S)\chi_{c0}$	< 4.7	$\times 10^{-6}$	CL=90%
Γ_{21}	$\psi(2S)\chi_{c1}$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{22}	$\psi(2S)\chi_{c2}$	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{23}	$\psi(2S)\eta_c(2S)$	< 3.3	$\times 10^{-6}$	CL=90%
Γ_{24}	$\psi(2S)X(3940)$	< 3.9	$\times 10^{-6}$	CL=90%
Γ_{25}	$\psi(2S)X(4160)$	< 3.9	$\times 10^{-6}$	CL=90%
Γ_{26}	$T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$	< 1.0	$\times 10^{-6}$	CL=90%
Γ_{27}	$T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$	< 1.67	$\times 10^{-5}$	CL=90%
Γ_{28}	$T_{c\bar{c}1}(3900)^{\pm} T_{c\bar{c}1}(4200)^{\mp}$	< 7.3	$\times 10^{-6}$	CL=90%
Γ_{29}	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	< 1.35	$\times 10^{-5}$	CL=90%
Γ_{30}	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	< 2.67	$\times 10^{-5}$	CL=90%
Γ_{31}	$T_{c\bar{c}}(4050)^{\pm} T_{c\bar{c}}(4250)^{\mp}$	< 2.72	$\times 10^{-5}$	CL=90%
Γ_{32}	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%
Γ_{33}	$T_{c\bar{c}}(4055)^{\pm} T_{c\bar{c}}(4055)^{\mp}$	< 1.11	$\times 10^{-5}$	CL=90%
Γ_{34}	$T_{c\bar{c}}(4055)^{\pm} T_{c\bar{c}1}(4430)^{\mp}$	< 2.11	$\times 10^{-5}$	CL=90%
Γ_{35}	$\overline{^2H}$ anything	(2.78 \pm 0.30)	$\times 10^{-5}$	S=1.2
Γ_{36}	hadrons	(94 \pm 11)	%	
Γ_{37}	ggg	(58.8 \pm 1.2)	%	
Γ_{38}	γgg	(1.87 \pm 0.28)	%	
Γ_{39}	$\phi K^+ K^-$	(1.6 \pm 0.4)	$\times 10^{-6}$	
Γ_{40}	$\omega \pi^+ \pi^-$	< 2.58	$\times 10^{-6}$	CL=90%
Γ_{41}	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(2.3 \pm 0.7)	$\times 10^{-6}$	
Γ_{42}	$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$	CL=90%
Γ_{43}	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%
Γ_{44}	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%
Γ_{45}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	(1.5 \pm 0.6)	$\times 10^{-6}$	
Γ_{46}	$K_1(1270)^{\pm} K^{\mp}$	< 3.22	$\times 10^{-6}$	CL=90%
Γ_{47}	$K_1(1400)^{\pm} K^{\mp}$	< 8.3	$\times 10^{-7}$	CL=90%
Γ_{48}	$b_1(1235)^{\pm} \pi^{\mp}$	< 4.0	$\times 10^{-7}$	CL=90%
Γ_{49}	$\rho \pi$	< 1.16	$\times 10^{-6}$	CL=90%
Γ_{50}	$\pi^+ \pi^- \pi^0$	< 8.0	$\times 10^{-7}$	CL=90%
Γ_{51}	$\omega \pi^0$	< 1.63	$\times 10^{-6}$	CL=90%
Γ_{52}	$\pi^+ \pi^- \pi^0 \pi^0$	(1.30 \pm 0.28)	$\times 10^{-5}$	
Γ_{53}	$K_S^0 K^+ \pi^- + \text{c.c.}$	(1.14 \pm 0.33)	$\times 10^{-6}$	

Γ_{54}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$< 4.22 \times 10^{-6}$	CL=90%
Γ_{55}	$K^*(892)^- K^+ + \text{c.c.}$	$< 1.45 \times 10^{-6}$	CL=90%
Γ_{56}	$f_1(1285) \text{anything}$	$(2.2 \pm 1.6) \times 10^{-3}$	
Γ_{57}	$f_1(1285) X_{\text{tetra}}$	$< 6.47 \times 10^{-5}$	CL=90%
Γ_{58}	$D_s^+ D_{s1}(2536)^-$		
Γ_{59}	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{60}	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.4 \pm 2.3) \times 10^{-6}$	
Γ_{61}	$D_s^{*+} D_{s1}(2536)^-$		
Γ_{62}	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.4 \pm 0.4) \times 10^{-5}$	
Γ_{63}	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.2 \pm 3.1) \times 10^{-6}$	
Γ_{64}	$D_s^+ D_{s2}^*(2573)^-$		
Γ_{65}	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(1.4 \pm 0.4) \times 10^{-5}$	
Γ_{66}	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(6.9 \pm 3.0) \times 10^{-6}$	
Γ_{67}	$D_s^{*+} D_{s2}^*(2573)^-$		
Γ_{68}	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(9 \pm 5) \times 10^{-6}$	
Γ_{69}	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(5 \pm 6) \times 10^{-6}$	
Γ_{70}	Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$	

Radiative decays

Γ_{71}	$\gamma \chi_{b1}(1P)$	$(6.9 \pm 0.4) \%$	
Γ_{72}	$\gamma \chi_{b2}(1P)$	$(7.15 \pm 0.35) \%$	
Γ_{73}	$\gamma \chi_{b0}(1P)$	$(3.8 \pm 0.4) \%$	
Γ_{74}	$\gamma f_0(1710)$	$< 5.9 \times 10^{-4}$	CL=90%
Γ_{75}	$\gamma f'_2(1525)$	$< 5.3 \times 10^{-4}$	CL=90%
Γ_{76}	$\gamma f_2(1270)$	$< 2.41 \times 10^{-4}$	CL=90%
Γ_{77}	$\gamma f_J(2220)$		
Γ_{78}	$\gamma \eta_c(1S)$	$< 2.7 \times 10^{-5}$	CL=90%
Γ_{79}	$\gamma \chi_{c0}$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{80}	$\gamma \chi_{c1}$	$< 3.6 \times 10^{-6}$	CL=90%
Γ_{81}	$\gamma \chi_{c2}$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{82}	$\gamma \chi_{c1}(3872)$	$< 2.3 \times 10^{-5}$	CL=90%
Γ_{83}	$\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	$< 2.4 \times 10^{-6}$	CL=90%
Γ_{84}	$\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi$	$< 2.8 \times 10^{-6}$	CL=90%

Γ_{85}	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%
Γ_{86}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{87}	$\gamma\eta_b(1S)$	(5.5 \pm 1.1) $\times 10^{-4}$		S=1.2
Γ_{88}	$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{89}	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%
Γ_{90}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%
Γ_{91}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%
Γ_{92}	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	< 8.3	$\times 10^{-6}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{93}	$e^\pm\tau^\mp$	<i>LF</i>	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{94}	$\mu^\pm\tau^\mp$	<i>LF</i>	< 3.3	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements to determine 2 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

$$\begin{array}{c} x_7 \\ \boxed{x_2} \\ x_1 \end{array}$$

$\Gamma(2S)\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_5/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$6.5 \pm 1.5 \pm 1.0$	KOBEL	92	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\Gamma(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_5/\Gamma$		
VALUE (eV)	EVTS	DOCUMENT ID	TECN
$105.4 \pm 1.0 \pm 4.2$	11.8k	¹ AUBERT	08BP BABR
¹ Using $B(\Gamma(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.			

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{36}\Gamma_5/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.577 ± 0.009 OUR AVERAGE			
0.581 $\pm 0.004 \pm 0.009$	¹ ROSNER	06	CLEO $10.0 \text{ e}^+\text{e}^- \rightarrow \text{hadrons}$
0.552 $\pm 0.031 \pm 0.017$	¹ BARU	96	MD1 $\text{e}^+\text{e}^- \rightarrow \text{hadrons}$
0.54 $\pm 0.04 \pm 0.02$	¹ JAKUBOWSKI	88	CBAL $\text{e}^+\text{e}^- \rightarrow \text{hadrons}$

0.58 $\pm 0.03 \pm 0.04$	² GILES	84B	CLEO	$e^+ e^- \rightarrow$ hadrons
0.60 $\pm 0.12 \pm 0.07$	² ALBRECHT	82	DASP	$e^+ e^- \rightarrow$ hadrons
0.54 $\pm 0.07 + 0.09 - 0.05$	² NICZYPORUK	81C	LENA	$e^+ e^- \rightarrow$ hadrons
0.41 ± 0.18	² BOCK	80	CNTR	$e^+ e^- \rightarrow$ hadrons

¹ Radiative corrections evaluated following KURAEV 85.² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Upsilon(2S)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$	Γ_5
VALUE (keV)	<u>DOCUMENT ID</u>
0.612 ± 0.011 OUR EVALUATION	

$\Upsilon(2S)$ BRANCHING RATIOS

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
Abbreviation MM in the <i>COMMENT</i> field below stands for missing mass.	

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.85 ± 0.26 OUR FIT				

17.92 ± 0.26 OUR AVERAGE

16.8 $\pm 1.1 \pm 1.3$	906k	¹ LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$
17.80 $\pm 0.05 \pm 0.37$	170k	² LEES	11L	BABR $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
18.02 $\pm 0.02 \pm 0.61$	851k	³ BHARI	09	CLEO $e^+ e^- \rightarrow \pi^+ \pi^-$ MM
17.22 $\pm 0.17 \pm 0.75$	11.8k	⁴ AUBERT	08BP	BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$
19.2 $\pm 0.2 \pm 1.0$	52.6k	⁵ ALEXANDER	98	CLE2 $\pi^+ \pi^- \ell^+ \ell^-$, $\pi^+ \pi^-$ MM
18.1 $\pm 0.5 \pm 1.0$	11.6k	ALBRECHT	87	ARG $e^+ e^- \rightarrow \pi^+ \pi^-$ MM
16.9 ± 4.0		GELPHMAN	85	CBAL $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
19.1 $\pm 1.2 \pm 0.6$		BESSON	84	CLEO $\pi^+ \pi^-$ MM
18.9 ± 2.6		FONSECA	84	CUSB $e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
21 ± 7	7	NICZYPORUK	81B	LENA $e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$

¹ LEES 11C reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

³ A weighted average of the inclusive and exclusive results.

⁴ Using $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

⁵ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_2/Γ
8.6 ± 0.4 OUR AVERAGE	

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 0.4 OUR AVERAGE				
8.43 $\pm 0.16 \pm 0.42$	38k	¹ BHARI	09	CLEO $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.2 $\pm 0.6 \pm 0.8$	275	² ALEXANDER	98	CLE2 $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.5 $\pm 1.9 \pm 1.9$	25	ALBRECHT	87	ARG $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
8.0 ± 1.5		GELPHMAN	85	CBAL $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
10.3 ± 2.3		FONSECA	84	CUSB $e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

² Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.462 ± 0.037 ¹ BHARI 09 CLEO $e^+ e^- \rightarrow \Upsilon(2S)$

¹ Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.00±0.21 OUR AVERAGE

$2.00 \pm 0.12 \pm 0.18$ 22k ¹ BESSON 07 CLEO $e^+ e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+ \tau^-$
 $1.7 \pm 1.5 \pm 0.6$ HAAS 84B CLEO $e^+ e^- \rightarrow \tau^+ \tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+ \mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.0193±0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.


$0.0203 \pm 0.0003 \pm 0.0008$	120k	ADAMS	05	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
$0.0122 \pm 0.0028 \pm 0.0019$		¹ KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$
$0.0138 \pm 0.0025 \pm 0.0015$		KAARSBERG	89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$
$0.009 \pm 0.006 \pm 0.006$		² ALBRECHT	85	ARG	$e^+ e^- \rightarrow \mu^+ \mu^-$
$0.018 \pm 0.008 \pm 0.005$		HAAS	84B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038 90 NICZYPORUK 81C LENA $e^+ e^- \rightarrow \mu^+ \mu^-$

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 0.026$.



$\Gamma(\tau^+ \tau^-)/\Gamma(\mu^+ \mu^-)$	Γ_3/Γ_4
<u>VALUE</u> 1.04±0.04±0.05	<u>EVTS</u> 22k <u>DOCUMENT ID</u> BESSON <u>TECN</u> CLEO <u>COMMENT</u> $e^+ e^- \rightarrow \gamma(2S)$

$\Gamma(\gamma(1S)\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	¹ TAMPONI	13	BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^0$
< 18	90	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<110	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<800	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+\ell^-\gamma\gamma$

¹ TAMPONI 13 reports $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^0)/\Gamma_{\text{total}}] / [\mathcal{B}(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)]$

< 2.3×10^{-4} which we multiply by our best value $\mathcal{B}(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.

² Authors assume $\mathcal{B}(\gamma(1S) \rightarrow e^+ e^-) + \mathcal{B}(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

$\Gamma(\gamma(1S)\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$	Γ_6/Γ_1
<u>VALUE (units 10^{-4})</u> <2.3	<u>CL%</u> 90 <u>DOCUMENT ID</u> TAMPONI <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \gamma(1S)\pi^0$

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

2.9 ± 0.4 OUR FIT Error includes scale factor of 2.0.**2.9 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

$2.39 \pm 0.31 \pm 0.14$	112	¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$
$2.1 \begin{array}{l} +0.7 \\ -0.6 \end{array} \pm 0.3$	14	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

$3.55 \pm 0.32 \pm 0.05$	241	³ TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\eta$
--------------------------	-----	----------------------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •


< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$
< 28	90	ALEXANDER98	CLE2		$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 70	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$
< 20	90	FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

² Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

³ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.



$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_1

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.64±0.25 OUR FIT					Error includes scale factor of 2.0.
1.99±0.14±0.11	241	TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.35 \pm 0.17 \pm 0.08$	1	LEES	11L	BABR	$\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
< 5.2	90	2	AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

¹ Not independent of other values reported by LEES 11L.² Not independent of other values reported by AUBERT 08BP.
 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$ Γ_6/Γ_7

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.13	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.006	90	MASCHMANN	90	CBAL $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 5.4 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 3.4 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.2 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.5 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.0 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c1} \text{anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.24 \pm 0.44 \pm 0.20$	376	JIA	17	$\gamma(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<36.7 \times 10^{-6}$	90	1 JIA	17A	$e^+ e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 36.7×10^{-6} .

 $\Gamma(\chi_{c2} \text{anything})/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.28 \pm 0.73 \pm 0.34$		JIA	17	$\gamma(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.1 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.7 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.5 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.9 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.3 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.9 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.9 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$

 $\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.0 \times 10^{-6}$	90	1 JIA	18	$\gamma(2S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$.

$\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<16.7 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.3 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$.

 $\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<13.5 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<26.7 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$ Γ_{31}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<27.2 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

 $\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<20.3 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2P) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<11.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}}$ Γ_{34}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<21.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{35}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.78^{+0.30}_{-0.26}$ OUR AVERAGE				Error includes scale factor of 1.2.

$2.64 \pm 0.11^{+0.26}_{-0.21}$ LEES 14G BABR $e^+ e^- \rightarrow \overline{2H} X$

$3.37 \pm 0.50 \pm 0.25$ 58 ASNER 07 CLEO $e^+ e^- \rightarrow \overline{2H} X$

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_{37}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
58.8 ± 1.2	6M	1 BESSON	06A CLEO	$\gamma(2S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+ \pi^- \gamma(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0 \pi^0 \gamma(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_{38}/Γ_{37}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A CLEO	$\gamma(2S) \rightarrow (\gamma +) \text{hadrons}$

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

$\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.58	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{41}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.32 \pm 0.40 \pm 0.54$	135	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.33	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.57	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$ Γ_{44}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{45}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.53 \pm 0.52 \pm 0.19$	32	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{46}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.22	90	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{47}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.83	90	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.40	90	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.16	90	SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.80	90	SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.63	90	SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$13.0 \pm 1.9 \pm 2.1$	261 ± 37	SHEN	13	BELL $\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.14 \pm 0.30 \pm 0.13$	40 ± 10	SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2 90 ¹ DOBBS 12A $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.22	90	SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.45	90	SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(f_1(1285)\text{anything})/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.20 \pm 1.50 \pm 0.63$	2.9k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(f_1(1285)X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<64.7 \times 10^{-6}$	90	¹ JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 7.8×10^{-6} to 64.7×10^{-6} .

$$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{\text{total}} \quad \Gamma_{59}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.6 \pm 0.3 \pm 0.2$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{\text{total}} \quad \Gamma_{60}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.84 \pm 0.18 \pm 0.15$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{\text{total}} \quad \Gamma_{62}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.4 \pm 0.4 \pm 0.2$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{\text{total}} \quad \Gamma_{63}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.82 \pm 0.25 \pm 0.19$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{\text{total}} \quad \Gamma_{65}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.4 \pm 0.4 \pm 0.2$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{\text{total}} \quad \Gamma_{66}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.69 \pm 0.20 \pm 0.22$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{\text{total}} \quad \Gamma_{68}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.9 \pm 0.5 \pm 0.2$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{\text{total}} \quad \Gamma_{69}/\Gamma$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.54 \pm 0.31 \pm 0.47$	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

$$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}} \quad \Gamma_{70}/\Gamma$$

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>COMMENT</u>
0.29 ± 0.03	1, ² DOBBS	12A $\gamma(2S) \rightarrow \text{hadrons}$

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$$\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.069 ± 0.004 OUR AVERAGE				
$0.0693 \pm 0.0012 \pm 0.0041$	407k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
$0.069 \pm 0.005 \pm 0.009$		EDWARDS	99	CLE2 $\gamma(2S) \rightarrow \gamma \chi(1P)$
$0.091 \pm 0.018 \pm 0.022$		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.065 \pm 0.007 \pm 0.012$		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
$0.080 \pm 0.017 \pm 0.016$		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.059 ± 0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0715 ± 0.0035 OUR AVERAGE				
0.0724 $\pm 0.0011 \pm 0.0040$	410k	ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
0.074 $\pm 0.005 \pm 0.008$		EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
0.098 $\pm 0.021 \pm 0.024$		ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.058 $\pm 0.007 \pm 0.010$		NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
0.102 $\pm 0.018 \pm 0.021$		HAAS 84	CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.061 ± 0.014		KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.038 ± 0.004 OUR AVERAGE				
0.0375 $\pm 0.0012 \pm 0.0047$	198k	ARTUSO 05	CLEO	$e^+e^- \rightarrow \gamma X$
0.034 $\pm 0.005 \pm 0.006$		EDWARDS 99	CLE2	$\gamma(2S) \rightarrow \gamma\chi(1P)$
0.064 $\pm 0.014 \pm 0.016$		ALBRECHT 85E	ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$
0.036 $\pm 0.008 \pm 0.009$		NERNST 85	CBAL	$e^+e^- \rightarrow \gamma X$
0.044 $\pm 0.023 \pm 0.009$		HAAS 84	CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.035 ± 0.014		KLOPFEN... 83	CUSB	$e^+e^- \rightarrow \gamma X$

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<59	90	¹ ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.9	90	² ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$
¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.				
² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$.				

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<53	90	¹ ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$
¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.				

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<24.1	90	¹ ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma\pi^+\pi^-$
¹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.				

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.8	90	¹ ALBRECHT 89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$
¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.				

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$				Γ_{78}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.7 \times 10^{-5}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$				Γ_{79}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.0 \times 10^{-4}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$				Γ_{80}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<3.6 \times 10^{-6}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$				Γ_{81}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.5 \times 10^{-5}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$				Γ_{82}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.3 \times 10^{-5}$	90	1 WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
¹ WANG 11B reports $[\Gamma(\gamma(2S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 0.8 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.					
$\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$				Γ_{83}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.4 \times 10^{-6}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$				Γ_{84}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.8 \times 10^{-6}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$				Γ_{85}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.2 \times 10^{-6}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$				Γ_{86}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.3 \times 10^{-6}$	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$				Γ_{87}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.5^{+1.1}_{-0.9}$ OUR AVERAGE			Error includes scale factor of 1.2.		
$6.1^{+0.6+0.9}_{-0.7-0.6}$		29k	FULSUM	18	BELL $\gamma(2S) \rightarrow \gamma X$
$3.9 \pm 1.1^{+1.1}_{-0.9}$	$13 \pm 5k$		¹ AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90		LEES	11J BABR	$\gamma(2S) \rightarrow X\gamma$

< 8.4	90	¹ BONVICINI	10	CLEO	$\gamma(2S) \rightarrow \gamma X$
< 5.1	90	² ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b}(1S) = 10$ MeV.² Superseded by BONVICINI 10. **$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$** Γ_{88}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.7 \times 10^{-6}$	90	SANDILYA	13	$\gamma(2S) \rightarrow \gamma$ hadrons

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 4.9	90		SANDILYA	13	$\gamma(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. **• • •**46.2 $^{+29.7}_{-14.2} \pm 10.6$ 10 ¹ DOBBS 12 $\gamma(2S) \rightarrow \gamma$ hadrons¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration. **$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$** Γ_{90}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.95	95	ROSNER	07A	CLEO $e^+ e^- \rightarrow \gamma X$

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	¹ LEES	11H	BABR $\gamma(2S) \rightarrow \gamma$ hadrons

¹ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} . **$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$** Γ_{92}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	¹ AUBERT	09Z	BABR $e^+ e^- \rightarrow A^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from $0.26-8.3 \times 10^{-6}$.**— LEPTON FAMILY NUMBER (LF) VIOLATING MODES —** **$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$** Γ_{93}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	LEES	10B	BABR $e^+ e^- \rightarrow e^\pm \tau^\mp$

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	LEES	10B	BABR $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**<14.4 95 LOVE 08A CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

$\Upsilon(2S)$ Cross-Particle Branching Ratios

$$\mathbf{B}(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times \mathbf{B}(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.78±0.02±0.11	906k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

 $\Upsilon(2S)$ REFERENCES

GAO	23	PR D108 112015	B.S. Gao <i>et al.</i>	(BELLE Collab.)
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUELL...	88	HE $e^+ e^-$ Physics 412	W. Buchmuller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (errat.)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelfman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
NERNST	85	Translated from YAF 41 733.		
ARTAMONOV	84	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)

FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
