

b -baryon ADMIXTURE (Λ_b , Ξ_b , Ω_b)

b -baryon ADMIXTURE MEAN LIFE

Each measurement of the b -baryon mean life is an average over an admixture of various b baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different b -baryon mean life. More b -baryon flavor specific channels are not included in the measurement.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.218 ^{+0.130} _{-0.115} ± 0.042		¹ ABAZOV	07S D0	Repl. by ABAZOV 12U
1.22 ^{+0.22} _{-0.18} ± 0.04		¹ ABAZOV	05C D0	Repl. by ABAZOV 07S
1.16 ± 0.20 ± 0.08		² ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.19 ± 0.14 ± 0.07		³ ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.14 ± 0.08 ± 0.04		⁴ ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.11 ^{+0.19} _{-0.18} ± 0.05		⁵ ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.29 ^{+0.24} _{-0.22} ± 0.06		⁵ ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.20 ± 0.08 ± 0.06		⁶ BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.21 ± 0.11		⁵ BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.32 ± 0.15 ± 0.07		⁷ ABE	96M CDF	$p\bar{p}$ at 1.8 TeV
1.46 ^{+0.22} _{-0.21} ± 0.07		ABREU	96D DLPH	Repl. by ABREU 99W
1.10 ^{+0.19} _{-0.17} ± 0.09		⁵ ABREU	96D DLPH	$e^+ e^- \rightarrow Z$
1.16 ± 0.11 ± 0.06		⁵ AKERS	96 OPAL	$e^+ e^- \rightarrow Z$
1.27 ^{+0.35} _{-0.29} ± 0.09		ABREU	95S DLPH	Repl. by ABREU 99W
1.05 ^{+0.12} _{-0.11} ± 0.09	290	BUSKULIC	95L ALEP	Repl. by BARATE 98D
1.04 ^{+0.48} _{-0.38} ± 0.10	11	⁸ ABREU	93F DLPH	Excess $\Lambda\mu^-$, decay lengths
1.05 ^{+0.23} _{-0.20} ± 0.08	157	⁹ AKERS	93 OPAL	Excess $\Lambda\ell^-$, decay lengths
1.12 ^{+0.32} _{-0.29} ± 0.16	101	¹⁰ BUSKULIC	92I ALEP	Excess $\Lambda\ell^-$, impact parameters

¹ Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

² Measured using $\Lambda\ell^-$ decay length.

³ Measured using $p\ell^-$ decay length.

⁴ This ABREU 99W result is the combined result of the $\Lambda\ell^-$, $p\ell^-$, and excess $\Lambda\mu^-$ impact parameter measurements.

⁵ Measured using $\Lambda_c\ell^-$ and $\Lambda\ell^+\ell^-$.

⁶ Measured using the excess of $\Lambda\ell^-$, lepton impact parameter.

⁷ Measured using $\Lambda_c\ell^-$.

⁸ ABREU 93F superseded by ABREU 96D.

⁹ AKERS 93 superseded by AKERS 96.¹⁰ BUSKULIC 92I superseded by BUSKULIC 95L.

b -baryon ADMIXTURE DECAY MODES ($\Lambda_b, \Xi_b, \Omega_b$)

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates at the LHC, LEP, and Tevatron, branching ratios, and detection efficiencies. They scale with the b -baryon production fraction $B(b \rightarrow b\text{-baryon})$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note “Production and Decay of b -Flavored Hadrons.”

For inclusive branching fractions, e.g., $B \rightarrow D^\pm \text{anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Scale factor
$\Gamma_1 p \mu^- \bar{\nu} \text{anything}$	$(5.8^{+2.3}_{-2.0})\%$	
$\Gamma_2 p \ell \bar{\nu}_\ell \text{anything}$	$(5.6 \pm 1.2)\%$	
$\Gamma_3 p \text{anything}$	$(70 \pm 22)\%$	
$\Gamma_4 \Lambda \ell^- \bar{\nu}_\ell \text{anything}$	$(3.8 \pm 0.6)\%$	
$\Gamma_5 \Lambda \ell^+ \nu_\ell \text{anything}$	$(3.2 \pm 0.8)\%$	
$\Gamma_6 \Lambda \text{anything}$	$(39 \pm 7)\%$	
$\Gamma_7 \Xi^- \ell^- \bar{\nu}_\ell \text{anything}$	$(4.6 \pm 1.4) \times 10^{-3}$	1.2

b -baryon ADMIXTURE ($\Lambda_b, \Xi_b, \Omega_b$) BRANCHING RATIOS

$\Gamma(p \mu^- \bar{\nu} \text{anything})/\Gamma_{\text{total}}$	Γ_1/Γ
$5.8^{+2.2}_{-1.9} \pm 0.8$	

¹ ABREU 95S reports $[\Gamma(b\text{-baryon} \rightarrow p \mu^- \bar{\nu} \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $0.0049 \pm 0.0011^{+0.0015}_{-0.0011}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon})$ = $(8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment’s error and our second error is the systematic error from using our best value.

$\Gamma(p \ell \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$	Γ_2/Γ
$5.6 \pm 0.9 \pm 0.7$	

¹ BARATE 98V reports $[\Gamma(b\text{-baryon} \rightarrow p \ell \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $(4.72 \pm 0.66 \pm 0.44) \times 10^{-3}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon})$ = $(8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment’s error and our second error is the systematic error from using our best value.

$\Gamma(p\ell\bar{\nu}_\ell \text{anything})/\Gamma(p\text{anything})$	Γ_2/Γ_3		
VALUE (%)	DOCUMENT ID	TECN	COMMENT
$8.0 \pm 1.2 \pm 1.4$	BARATE 98V	ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(\Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_4/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(b \rightarrow b\text{-baryon})$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(b \rightarrow b\text{-baryon})$ as described in the note on “Production and Decay of b -Flavored Hadrons.”

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.6 OUR AVERAGE				
$3.9 \pm 0.5 \pm 0.5$		1 BARATE 98D	ALEP	$e^+ e^- \rightarrow Z$
$3.5 \pm 0.4 \pm 0.5$		2 AKERS 96	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$3.6 \pm 0.9 \pm 0.5$	262	3 ABREU 95S	DLPH	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$7.3 \pm 1.4 \pm 1.0$	290	4 BUSKULIC 95L	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	157	5 AKERS 93	OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$8.3 \pm 2.5 \pm 1.1$	101	6 BUSKULIC 92I	ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

¹ BARATE 98D reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00326 \pm 0.00016 \pm 0.00039$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using the excess of $\Lambda\ell^-$, lepton impact parameter.

² AKERS 96 reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00291 \pm 0.00023 \pm 0.00025$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABREU 95S reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0030 \pm 0.0006 \pm 0.0004$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BUSKULIC 95L reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0061 \pm 0.0006 \pm 0.0010$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AKERS 93 superseded by AKERS 96.

⁶ BUSKULIC 92I reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0070 \pm 0.0010 \pm 0.0018$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

$\Gamma(\Lambda\ell^+\nu_\ell \text{anything})/\Gamma(\Lambda \text{anything})$	Γ_5/Γ_6		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$8.0 \pm 1.2 \pm 0.8$	ABBIENDI 99L	OPAL	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
7.0 $\pm 1.2 \pm 0.7$	ACKERSTAFF 97N	OPAL	Repl. by ABBIENDI 99L

$\Gamma(\Lambda\text{anything})/\Gamma_{\text{total}}$		Γ_6/Γ	
VALUE (%)	DOCUMENT ID	TECN	COMMENT
39± 7 OUR AVERAGE			
42± 6±5	¹ ABBIENDI 99L	OPAL	$e^+ e^- \rightarrow Z$
$27^{+15}_{-9} \pm 3$	² ABREU 95C	DLPH	$e^+ e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
47± 7±6	³ ACKERSTAFF 97N	OPAL	Repl. by ABBIENDI 99L
¹ ABBIENDI 99L reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.035 \pm 0.0032 \pm 0.0035$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
² ABREU 95C reports $0.28^{+0.17}_{-0.12}$ from a measurement of $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ assuming $B(\bar{b} \rightarrow b\text{-baryon}) = 0.08 \pm 0.02$, which we rescale to our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
³ ACKERSTAFF 97N reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0393 \pm 0.0046 \pm 0.0037$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

$\Gamma(\Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}$		Γ_7/Γ	
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.6±1.4 OUR AVERAGE	Error includes scale factor of 1.2.		
3.6±1.2±0.5	¹ ABDALLAH 05C	DLPH	$e^+ e^- \rightarrow Z^0$
$6.4^{+1.6}_{-1.6} \pm 0.8$	² BUSKULIC 96T	ALEP	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
7.0±2.8±0.9	³ ABREU 95V	DLPH	Repl. by ABDALLAH 05C
¹ ABDALLAH 05C reports $[\Gamma(b\text{-baryon} \rightarrow \Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = (3.0 \pm 1.0 \pm 0.3) \times 10^{-4}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
² BUSKULIC 96T reports $[\Gamma(b\text{-baryon} \rightarrow \Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = (5.4 \pm 1.1 \pm 0.8) \times 10^{-4}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
³ ABREU 95V reports $[\Gamma(b\text{-baryon} \rightarrow \Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = (5.9 \pm 2.1 \pm 1.0) \times 10^{-4}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

***b*-baryon ADMIXTURE (Λ_b , Ξ_b , Ω_b) REFERENCES**

ABAZOV	12U	PR D85 112003	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07S	PRL 99 142001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05C	PRL 94 102001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	05C	EPJ C44 299	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABBIENDI	99L	EPJ C9 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	99W	EPJ C10 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98D	EPJ C2 197	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ABE	96M	PRL 77 1439	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96D	ZPHY C71 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	95C	PL B347 447	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	93F	PL B311 379	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	93	PL B316 435	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)