

$\Lambda(2100) 7/2^-$  $I(J^P) = 0(\frac{7}{2}^-)$  Status: \* \* \* \*

Most of the results published before 1973 are now obsolete and have been omitted. They may be found in our 1982 edition Physics Letters **111B** 1 (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and in invariant-mass distributions around 2100 MeV used to be listed in a separate entry immediately following. It may be found in our 1986 edition Physics Letters **170B** 1 (1986).

## $\Lambda(2100)$ POLE POSITION

### REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2040 ± 14</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2023	ZHANG	13A DPWA	Multichannel

### −2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>215 ± 29</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
239	ZHANG	13A DPWA	Multichannel

## $\Lambda(2100)$ POLE RESIDUE

The “normalized residue” is the residue divided by  $\Gamma_{pole}/2$ .

### Normalized residue in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
<b>0.28 ± 0.06</b>	<b>−40 ± 10</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

### Normalized residue in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma\pi$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
<b>0.09 ± 0.02</b>	<b>−35 ± 15</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

### Normalized residue in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma(1385)\pi$ , D-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
<b>0.04 ± 0.03</b>		SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

### Normalized residue in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma(1385)\pi$ , G-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
<b>0.06 ± 0.03</b>	<b>−45 ± 15</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

**Normalized residue in  $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892)$ ,  $S=3/2$ ,  $D$ -wave**

<u>MODULUS</u>	<u>PHASE (<math>^\circ</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.11±0.06</b>	<b>-30 ± 30</b>	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

 **$\Lambda(2100)$  MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2090 to 2110 (<math>\approx</math> 2100) OUR ESTIMATE</b>			
2090±15	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
2086±6	ZHANG 13A	DPWA	Multichannel
2104±10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2106±30	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2110±10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
2105±10	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$
2115±10	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2094	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
2094	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2110 or 2089	<sup>1</sup> NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 **$\Lambda(2100)$  WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>100 to 250 (<math>\approx</math> 200) OUR ESTIMATE</b>			
290±30	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
305±16	ZHANG 13A	DPWA	Multichannel
157±40	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
250±30	GOPAL 77	DPWA	$\bar{K}N$ multichannel
241±30	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$
152±15	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
98	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
250	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
244 or 302	<sup>1</sup> NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 **$\Lambda(2100)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $N\bar{K}$	25–35 %
$\Gamma_2$ $\Sigma\pi$	~ 5 %
$\Gamma_3$ $\Lambda\eta$	<3 %
$\Gamma_4$ $\Xi K$	<3 %
$\Gamma_5$ $\Lambda\omega$	<8 %
$\Gamma_6$ $\Sigma(1385)\pi$ , $D$ -wave	
$\Gamma_7$ $\Sigma(1385)\pi$ , $G$ -wave	(1.0±1.0) %
$\Gamma_8$ $N\bar{K}^*(892)$	10–20 %

$\Gamma_9$	$N\bar{K}^*(892)$ , $S=3/2$ , $D$ -wave	$(4.0 \pm 2.0) \%$
$\Gamma_{10}$	$N\bar{K}^*(892)$ , $S=1/2$ , $G$ -wave	
$\Gamma_{11}$	$N\bar{K}^*(892)$ , $S=3/2$ , $G$ -wave	

## $\Lambda(2100)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

### 0.25 to 0.35 ( $\approx 0.30$ ) OUR ESTIMATE

$0.24 \pm 0.05$	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
$0.23 \pm 0.01$	ZHANG	13A	DPWA	Multichannel
$0.34 \pm 0.03$	GOPAL	80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
$0.24 \pm 0.06$	DEBELLEFON	78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
$0.31 \pm 0.03$	HEMINGWAY	75	DPWA	$K^- p \rightarrow \bar{K}N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.29$	DECLAIS	77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
$0.30 \pm 0.03$	GOPAL	77	DPWA	See GOPAL 80

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$0.030 \pm 0.015$	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
-------------------	-----------	----	------	-------------------------

$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$				$\Gamma_6/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$< 0.01$	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
----------	-----------	----	------	-------------------------

$\Gamma(\Sigma(1385)\pi, G\text{-wave})/\Gamma_{\text{total}}$				$\Gamma_7/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$0.01 \pm 0.01$	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
-----------------	-----------	----	------	-------------------------

$\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$				$\Gamma_9/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$0.04 \pm 0.02$	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
-----------------	-----------	----	------	-------------------------

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma\pi$				$(\Gamma_1 \Gamma_2)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$+0.03 \pm 0.01$	ZHANG	13A	DPWA	Multichannel
$+0.12 \pm 0.04$	GOPAL	77	DPWA	$\bar{K}N$ multichannel
$+0.11 \pm 0.01$	KANE	74	DPWA	$K^- p \rightarrow \Sigma\pi$

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda\eta$				$(\Gamma_1 \Gamma_3)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

$-0.050 \pm 0.020$	RADER	73	MPWA	$K^- p \rightarrow \Lambda\eta$
--------------------	-------	----	------	---------------------------------

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Xi K$				$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.035 \pm 0.018$	LITCHFIELD	71	DPWA	$K^- p \rightarrow \Xi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.003	MULLER	69B	DPWA	$K^- p \rightarrow \Xi K$
0.05	TRIPP	67	RVUE	$K^- p \rightarrow \Xi K$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda \omega$				$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
-0.070	<sup>2</sup> BACCARI	77	DPWA	$GD_{37}$ wave
+0.011	<sup>2</sup> BACCARI	77	DPWA	$GG_{17}$ wave
+0.008	<sup>2</sup> BACCARI	77	DPWA	$GG_{37}$ wave
0.122 or 0.154	<sup>1</sup> NAKKASYAN	75	DPWA	$K^- p \rightarrow \Lambda \omega$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892), S=3/2, D\text{-wave}$				$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$+0.16 \pm 0.02$	ZHANG	13A	DPWA	Multichannel
$+0.21 \pm 0.04$	CAMERON	78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892), S=1/2, G\text{-wave}$				$(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$-0.03 \pm 0.02$	ZHANG	13A	DPWA	Multichannel
$-0.04 \pm 0.03$	<sup>3</sup> CAMERON	78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892), S=3/2, G\text{-wave}$				$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$+0.08 \pm 0.02$	ZHANG	13A	DPWA	Multichannel

### $\Lambda(2100)$ FOOTNOTES

<sup>1</sup> The NAKKASYAN 75 values are from the two best solutions found. Each has the  $\Lambda(2100)$  and one additional resonance ( $P_3$  or  $F_5$ ).

<sup>2</sup> Note that the three for BACCARI 77 entries are for three different waves.

<sup>3</sup> The published sign has been changed to be in accord with the baryon-first convention. The upper limit on the  $G_3$  wave is 0.03.

### $\Lambda(2100)$ REFERENCES

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
PDG	86	PL 170B 1	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
CAMERON	78B	NP B146 327	W. Cameron <i>et al.</i>	(RHEL, LOIC) IJP
DEBELLEFON	78	NC 42A 403	A. de Bellefon <i>et al.</i>	(CDEF, SACL) IJP
BACCARI	77	NC 41A 96	B. Baccari <i>et al.</i>	(SACL, CDEF) IJP
DECLAIS	77	CERN 77-16	Y. Declais <i>et al.</i>	(CAEN, CERN) IJP

GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
HEMINGWAY	75	NP B91 12	R.J. Hemingway <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
NAKKASYAN	75	NP B93 85	A. Nakkasyan	(CERN) IJP
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
RADER	73	NC 16A 178	R.K. Rader <i>et al.</i>	(SACL, HEID, CERN+)
LITCHFIELD	71	NP B30 125	P.J. Litchfield <i>et al.</i>	(RHEL, CDEF, SACL) IJP
MULLER	69B	Thesis UCRL 19372	R.A. Muller	(LRL)
TRIPP	67	NP B3 10	R.D. Tripp <i>et al.</i>	(LRL, SLAC, CERN+)

---