

CHARMED MESONS ($C = \pm 1$)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\overline{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^* 's

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1869.66 \pm 0.05$ MeV

Mean life $\tau = (1033 \pm 5) \times 10^{-15}$ s

$$c\tau = 309.8 \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.096 \pm 0.004 \text{ [a]}$$

$$\Gamma(c \rightarrow D^*(2010)^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_L^0 e^\pm \nu) = (-0.6 \pm 1.6)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K_L^0 K^\pm) \text{ in } D^\pm \rightarrow K_L^0 K^\pm = (-4.2 \pm 3.4) \times 10^{-2}$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.18 \pm 0.16)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (-0.3 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (-0.1 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \eta) \text{ in } D^\pm \rightarrow K_S^0 \pi^\pm \eta = (-0.9 \pm 3.1) \times 10^{-2}$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.0 \pm 1.2)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^- \pi^0) \text{ in } D^\pm \rightarrow K^\pm \pi^+ \pi^- \pi^0 = -0.04 \pm 0.06$$

$$A_{CP}(\pi^\pm \pi^0) = (0.4 \pm 1.3)\% \quad (S = 1.7)$$

$$A_{CP}(\pi^\pm \eta) = (0.3 \pm 0.5)\%$$

$$A_{CP}(\pi^\pm \pi^0 \eta) \text{ in } D^\pm \rightarrow \pi^\pm \pi^0 \eta = (-6 \pm 7) \times 10^{-2}$$

$$A_{CP}(\pi^\pm \eta \eta) \text{ in } D^\pm \rightarrow \pi^\pm \eta \eta = (8 \pm 9) \times 10^{-2}$$

$$A_{CP}(\pi^\pm \eta'(958)) = (0.41 \pm 0.23)\% \quad (S = 1.2)$$

$$A_{CP}(\overline{K}^0 / K^0 K^\pm) = (0.11 \pm 0.17)\%$$

$$A_{CP}(K_S^0 K^\pm) = (-0.01 \pm 0.07)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_S^0 K^\pm \pi^0 = (1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K_L^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_L^0 K^\pm \pi^0 = (-1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.37 \pm 0.29)\%$$

$$A_{CP}(K^\pm K^{*0}) = (-0.3 \pm 0.4)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.01 \pm 0.09)\% \quad (S = 1.8)$$

$$A_{CP}(K^\pm K_0^*(1430)^0) = (8^{+7}_{-6})\%$$

$$A_{CP}(K^\pm K_2^*(1430)^0) = (43^{+20}_{-26})\%$$

$$A_{CP}(K^\pm K_0^*(700)) = (-12^{+18}_{-13})\%$$

$$\begin{aligned}
A_{CP}(a_0(1450)^0 \pi^\pm) &= (-19^{+14}_{-16})\% \\
A_{CP}(\phi(1680)\pi^\pm) &= (-9 \pm 26)\% \\
A_{CP}(\pi^\pm 2\pi^0) \text{ in } D^\pm \rightarrow \pi^\pm 2\pi^0 &= (5.6 \pm 2.7)\% \\
A_{CP}(\pi^+ \pi^- \pi^\pm) &= (0.5 \pm 2.0)\% \\
A_{CP}(2\pi^\pm \pi^\mp \pi^0) \text{ in } D^\pm \rightarrow 2\pi^\pm \pi^\mp \pi^0 &= (0.3 \pm 2.0)\% \\
A_{CP}(2\pi^\pm \pi^\mp 2\pi^0) \text{ in } D^\pm \rightarrow 2\pi^\pm \pi^\mp 2\pi^0 &= (-4 \pm 4)\% \\
A_{CP}(\pi^+ \pi^- \pi^\pm \eta) \text{ in } D^\pm \rightarrow \pi^+ \pi^- \pi^\pm \eta &= (3 \pm 5) \times 10^{-2} \\
A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) &= (-4 \pm 7)\% \\
A_{CP}(K^\pm \pi^0) &= (-3 \pm 5)\% \\
A_{CP}(K^\pm \eta) \text{ in } D^\pm \rightarrow K^\pm \eta &= (-6 \pm 11) \times 10^{-2}
\end{aligned}$$

χ^2 tests of CP -violation (CPV)

Local CPV in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm = 78.1\%$

Local CPV in $D^\pm \rightarrow K^+ K^- \pi^\pm = 31\%$

Local CPV in $D^\pm \rightarrow K^+ K^- K^\pm = 31.6\%$

CP violating asymmetries of P -odd (T -odd) moments

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-3 \pm 8) \times 10^{-3} [b] \quad (S = 1.1)$$

$$A_{T\text{viol}}(K^+ K^- K_S^0 \pi^\pm) \text{ in } D^\pm \rightarrow K^+ K^- K_S^0 \pi^\pm = (-3.3 \pm 2.7)\%$$

D^+ form factors

$$f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell = 0.719 \pm 0.011 \quad (S = 1.6)$$

$$r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -2.13 \pm 0.14$$

$$r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell = -3 \pm 12 \quad (S = 1.5)$$

$$f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell = 0.1407 \pm 0.0025$$

$$r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -2.00 \pm 0.13$$

$$r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell = -4 \pm 5$$

$$f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta \ell^+ \nu_\ell (\ell = e \text{ or } \nu) = (8.4 \pm 0.4) \times 10^{-2}$$

$$r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e = -5.3 \pm 2.7 \quad (S = 1.9)$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.24 \pm 0.11$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e = 1.06 \pm 0.16$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 1.64 \pm 0.10 \quad (S = 1.2)$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e = 0.84 \pm 0.06$$

$$r_\nu \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.49 \pm 0.05 \quad (S = 2.1)$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.802 \pm 0.021$$

$$r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.0 \pm 0.4$$

$$\Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 1.13 \pm 0.08$$

$$\Gamma_+/ \Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell = 0.22 \pm 0.06 \quad (S = 1.6)$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 ± 0.30) %		—
μ^+ anything	(17.6 ± 3.2) %		—
K^- anything	(25.7 ± 1.4) %		—
K_S^0 anything	(33.1 ± 0.4) %		—
K^+ anything	(5.9 ± 0.8) %		—
$K^*(892)^-$ anything	(6 ± 5) %		—
$\bar{K}^*(892)^0$ anything	(23 ± 5) %		—
$K^*(892)^0$ anything	< 6.6 %	CL=90%	—
η anything	(6.3 ± 0.7) %		—
η' anything	(1.04 ± 0.18) %		—
ϕ anything	(1.12 ± 0.04) %		—
$\pi^+ \pi^+ \pi^-$ anything	(15.25 ± 0.20) %		—
Leptonic and semileptonic modes			
$e^+ \nu_e$	< 8.8 $\times 10^{-6}$	CL=90%	935
$\gamma e^+ \nu_e$	< 3.0 $\times 10^{-5}$	CL=90%	935
$\mu^+ \nu_\mu$	(3.74 ± 0.17) $\times 10^{-4}$		932
$\tau^+ \nu_\tau$	(1.20 ± 0.27) $\times 10^{-3}$		90
$\bar{K}^0 e^+ \nu_e$	(8.72 ± 0.09) %		869
$\bar{K}^0 \mu^+ \nu_\mu$	(8.76 ± 0.19) %		865
$K^- \pi^+ e^+ \nu_e$	(4.02 ± 0.18) %	S=3.2	864
$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.77 ± 0.17) %		722
$(K^- \pi^+) [0.8-1.0]\text{GeV} e^+ \nu_e$	(3.39 ± 0.09) %		864
$(K^- \pi^+)_{S-wave} e^+ \nu_e$	(2.28 ± 0.11) $\times 10^{-3}$		—
$\bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6 $\times 10^{-3}$	CL=90%	—
$\bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5 $\times 10^{-4}$	CL=90%	—
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 $\times 10^{-3}$	CL=90%	864
$\bar{K}^*(892)^0 e^+ \nu_e$	(5.40 ± 0.10) %	S=1.1	722
$K^- \pi^+ \mu^+ \nu_\mu$	(3.65 ± 0.34) %		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52 ± 0.10) %		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(1.9 ± 0.5) $\times 10^{-3}$		851

$\overline{K}^*(892)^0 \mu^+ \nu_\mu$	(5.27 \pm 0.15) %	717
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$ CL=90%	825
$\overline{K}_1(1270)^0 e^+ \nu_e, \overline{K}_1^0 \rightarrow K^- \pi^+ \pi^0$	(1.06 \pm 0.15) $\times 10^{-3}$	-
$\overline{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.3 $\times 10^{-4}$ CL=90%	380
$\overline{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$ CL=90%	105
$\pi^0 e^+ \nu_e$	(3.72 \pm 0.17) $\times 10^{-3}$ S=2.0	930
$\pi^0 \mu^+ \nu_\mu$	(3.50 \pm 0.15) $\times 10^{-3}$	927
$\eta e^+ \nu_e$	(1.11 \pm 0.07) $\times 10^{-3}$	855
$\eta \mu^+ \nu_\mu$	(1.04 \pm 0.11) $\times 10^{-3}$	851
$\pi^- \pi^+ e^+ \nu_e$	(2.49 \pm 0.11) $\times 10^{-3}$ S=1.2	924
$f_0(500)^0 e^+ \nu_e, f_0(500)^0 \rightarrow \pi^+ \pi^-$	(6.4 \pm 0.6) $\times 10^{-4}$	-
$\rho^0 e^+ \nu_e$	(1.90 \pm 0.10) $\times 10^{-3}$ S=1.2	774
$\rho^0 \mu^+ \nu_\mu$	(2.4 \pm 0.4) $\times 10^{-3}$	770
$\omega e^+ \nu_e$	(1.69 \pm 0.11) $\times 10^{-3}$	771
$\omega \mu^+ \nu_\mu$	(1.77 \pm 0.21) $\times 10^{-3}$	767
$\eta'(958) e^+ \nu_e$	(2.0 \pm 0.4) $\times 10^{-4}$	690
$a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$	(1.7 \pm 0.8) $\times 10^{-4}$	-
$b_1(1235)^0 e^+ \nu_e, b_1^0 \rightarrow \omega \pi^0$	< 1.75 $\times 10^{-4}$ CL=90%	-
$\phi e^+ \nu_e$	< 1.3 $\times 10^{-5}$ CL=90%	657
$D^0 e^+ \nu_e$	< 1.0 $\times 10^{-4}$ CL=90%	5

Hadronic modes with a \overline{K} or $\overline{K} \overline{K} \overline{K}$

$K_S^0 \pi^+$	(1.562 \pm 0.031) %	S=1.7	863
$K_L^0 \pi^+$	(1.46 \pm 0.05) %		863
$K^- 2\pi^+$	[c] (9.38 \pm 0.16) %	S=1.6	846
$(K^- \pi^+)_{S-\text{wave}} \pi^+$	(7.52 \pm 0.17) %		846
$\overline{K}_0^*(1430)^0 \pi^+, \overline{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] (1.25 \pm 0.06) %		382
$\overline{K}^*(892)^0 \pi^+, \overline{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.04 \pm 0.12) %		714
$\overline{K}^*(1410)^0 \pi^+, \overline{K}^{*0} \rightarrow K^- \pi^+$	not seen		381
$\overline{K}_2^*(1430)^0 \pi^+, \overline{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[d] (2.3 \pm 0.7) $\times 10^{-4}$		371
$\overline{K}^*(1680)^0 \pi^+, \overline{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] (2.2 \pm 1.1) $\times 10^{-4}$		58
$K^- (2\pi^+)_{I=2}$	(1.45 \pm 0.26) %		-
$K_S^0 \pi^+ \pi^0$	[c] (7.36 \pm 0.20) %		845
$K_S^0 \rho^+$	(6.14 \pm 0.35) %		677
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	(1.5 \pm 1.2) $\times 10^{-3}$		-

$\overline{K}^*(892)^0 \pi^+$, $\overline{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	$(2.64 \pm 0.32) \times 10^{-3}$	714
$\overline{K}_0^*(1430)^0 \pi^+$, $\overline{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(2.7 \pm 0.9) \times 10^{-3}$	—
$\overline{K}_0^*(1680)^0 \pi^+$, $\overline{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(10 \pm 7) \times 10^{-4}$	—
$\overline{\kappa}^0 \pi^+$, $\overline{\kappa}^0 \rightarrow K_S^0 \pi^0$	$(6 \pm 5) \times 10^{-3}$	—
$K_S^0 \pi^+ \pi^0$ nonresonant	$(3 \pm 4) \times 10^{-3}$	845
$K_S^0 \pi^+ \pi^0$ nonresonant and $\overline{\kappa}^0 \pi^+$	$(1.37 \pm 0.21) \%$	—
$(K_S^0 \pi^0)_{S\text{-wave}} \pi^+$	$(1.27 \pm 0.27) \%$	845
$K_S^0 \pi^+ \omega$	$(7.1 \pm 0.5) \times 10^{-3}$	606
$K_S^0 \pi^+ \eta$	$(1.31 \pm 0.05) \%$	722
$K_S^0 \pi^+ \eta'(958)$	$(1.90 \pm 0.21) \times 10^{-3}$	481
$K^- 2\pi^+ \pi^0$	[e] $(6.25 \pm 0.18) \%$	817
$K_S^0 2\pi^+ \pi^-$	[e] $(3.10 \pm 0.09) \%$	814
$K_S^0 \pi^+ 2\pi^0$	$(2.89 \pm 0.09) \%$	817
$K_S^0 a_1(1260)^+$, $a_1^+ \rightarrow \rho(770)^+ \pi^0$	$(8.7 \pm 1.6) \times 10^{-3}$	—
$K_S^0 a_1(1260)^+$, $a_1^+ \rightarrow f_0(500) \pi^+$, $f_0 \rightarrow \pi^0 \pi^0$	$(1.0 \pm 0.6) \times 10^{-3}$	—
$\overline{K}_1(1400)^0 \pi^+$, $\overline{K}_1^0 \rightarrow \overline{K}^*(892)^0 \pi^0$, $\overline{K}^* \rightarrow K_S^0 \pi^0$	$(2.3 \pm 0.4) \times 10^{-3}$	—
$\overline{K}^*(892)^0 \rho^+$, $\overline{K}^* \rightarrow K_S^0 \pi^0$	$(9.7 \pm 0.9) \times 10^{-3}$	—
$\overline{K}^*(892)^0 \pi^+ \pi^0$ non-resonant, $\overline{K}^* \rightarrow K_S^0 \pi^0$	$(2.6 \pm 0.7) \times 10^{-3}$	—
$K_S^0 \rho^+ \pi^0$ non-resonant	$(4.8 \pm 0.5) \times 10^{-3}$	—
$K^- 2\pi^+ \eta$	$(1.35 \pm 0.12) \times 10^{-3}$	657
$K_S^0 \pi^+ \pi^0 \eta$	$(1.22 \pm 0.25) \times 10^{-3}$	657
$K^- 3\pi^+ \pi^-$	[c] $(5.7 \pm 0.5) \times 10^{-3}$ S=1.1	772
$\overline{K}^*(892)^0 2\pi^+ \pi^-$, $\overline{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.2 \pm 0.4) \times 10^{-3}$	645
$\overline{K}^*(892)^0 \rho^0 \pi^+$, $\overline{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.3 \pm 0.4) \times 10^{-3}$	239
$\overline{K}^*(892)^0 a_1(1260)^+$	[f] $(9.3 \pm 1.9) \times 10^{-3}$	†
$K^- \rho^0 2\pi^+$	$(1.72 \pm 0.28) \times 10^{-3}$	524
$K^- 3\pi^+ \pi^-$ nonresonant	$(4.0 \pm 2.9) \times 10^{-4}$	772
$K_S^0 2\pi^+ \pi^- \pi^0$	$(1.53 \pm 0.08) \%$	773
$K_S^0 \pi^+ 3\pi^0$	$(5.5 \pm 0.5) \times 10^{-3}$	776

$K^- 2\pi^+ 2\pi^0$	$(4.95 \pm 0.32) \times 10^{-3}$	776
$K^+ 2K_S^0$	$(2.54 \pm 0.13) \times 10^{-3}$	545
$K^+ K^- K_S^0 \pi^+$	$(2.4 \pm 0.5) \times 10^{-4}$	436

Pionic modes

$\pi^+ \pi^0$	$(1.247 \pm 0.033) \times 10^{-3}$	925
$2\pi^+ \pi^-$	$(3.27 \pm 0.09) \times 10^{-3}$	909
$\rho^0 \pi^+$	$(8.4 \pm 0.8) \times 10^{-4}$	767
$\pi^+(\pi^+\pi^-)_{S\text{-wave}}$	$(2.01 \pm 0.06) \times 10^{-3}$	909
$\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$	$(1.38 \pm 0.10) \times 10^{-3}$	—
$f_0(980)\pi^+, f_0 \rightarrow \pi^+ \pi^-$	$(1.57 \pm 0.32) \times 10^{-4}$	669
$f_0(1370)\pi^+, f_0 \rightarrow \pi^+ \pi^-$	$(8 \pm 4) \times 10^{-5}$	—
$\omega \pi^+, \omega \rightarrow \pi^+ \pi^-$	$(3.4 \pm 0.5) \times 10^{-6}$	—
$f_2(1270)\pi^+, f_2 \rightarrow \pi^+ \pi^-$	$(4.58 \pm 0.28) \times 10^{-4}$	485
$\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	338
$\rho(1700)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	—
$f_0(1500)\pi^+, f_0 \rightarrow \pi^+ \pi^-$	$(1.1 \pm 0.4) \times 10^{-4}$	—
$f_0(1710)\pi^+, f_0 \rightarrow \pi^+ \pi^-$	$< 5 \times 10^{-5} \text{ CL}=95\%$	—
$f_0(1790)\pi^+, f_0 \rightarrow \pi^+ \pi^-$	$< 7 \times 10^{-5} \text{ CL}=95\%$	—
$(\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	$< 1.2 \times 10^{-4} \text{ CL}=95\%$	909
$2\pi^+ \pi^- \text{ nonresonant}$	$< 1.1 \times 10^{-4} \text{ CL}=95\%$	909
$\pi^+ 2\pi^0$	$(4.61 \pm 0.15) \times 10^{-3}$	910
$2\pi^+ \pi^- \pi^0$	$(1.165 \pm 0.030) \%$	883
$\pi^+ 3\pi^0$	$(4.17 \pm 0.26) \times 10^{-3}$	885
$\pi^+ 4\pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$	851
$2\pi^+ \pi^- 2\pi^0$	$(1.07 \pm 0.05) \%$	848
$3\pi^+ 2\pi^-$	$(1.66 \pm 0.16) \times 10^{-3} \text{ S}=1.1$	845
$2\pi^+ \pi^- 3\pi^0$	$(3.42 \pm 0.35) \times 10^{-3}$	803
$3\pi^+ 2\pi^- \pi^0$	$(2.34 \pm 0.27) \times 10^{-3}$	799
$\eta \pi^+$	$(3.77 \pm 0.09) \times 10^{-3}$	848
$\eta \pi^+ \pi^0$	$(2.05 \pm 0.35) \times 10^{-3} \text{ S}=2.2$	831
$\eta 2\pi^+ \pi^-$	$(3.41 \pm 0.20) \times 10^{-3}$	798
$\eta \pi^+ 2\pi^0$	$(3.20 \pm 0.33) \times 10^{-3}$	801
$\eta \pi^+ 3\pi^0$	$(2.9 \pm 0.5) \times 10^{-3}$	759
$\eta 2\pi^+ \pi^- \pi^0$	$(3.88 \pm 0.34) \times 10^{-3}$	755
$\eta \eta \pi^+$	$(2.96 \pm 0.26) \times 10^{-3}$	700
$\omega \pi^+$	$(2.8 \pm 0.6) \times 10^{-4}$	764
$\omega \pi^+ \pi^0$	$(3.9 \pm 0.9) \times 10^{-3}$	742
$\eta'(958)\pi^+$	$(4.97 \pm 0.19) \times 10^{-3}$	681
$\eta'(958)\pi^+ \pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$	654

Hadronic modes with a $K\bar{K}$ pair

$K_S^0 K^+$	$(3.04 \pm 0.09) \times 10^{-3} \text{ S}=2.2$	793
$K_L^0 K^+$	$(3.21 \pm 0.16) \times 10^{-3}$	793
$K_S^0 K^+ \pi^0$	$(5.07 \pm 0.30) \times 10^{-3}$	744

$K^*(892)^+ K_S^0, \quad K^{*+} \rightarrow$	$(2.89 \pm 0.30) \times 10^{-3}$	612
$\overline{K}^*(892)^0 K^+, \quad \overline{K}^{*0} \rightarrow$	$(5.2 \pm 1.4) \times 10^{-4}$	613
$K_S^0 \pi^0$		
$K_L^0 K^+ \pi^0$	$(5.24 \pm 0.31) \times 10^{-3}$	744
$K^+ K^- \pi^+$	[c] $(9.68 \pm 0.18) \times 10^{-3}$	744
$K^+ \overline{K}^*(892)^0,$	$(2.49 \pm 0.08) \times 10^{-3}$	613
$\overline{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K^+ \overline{K}_0^*(1430)^0,$	$(1.82 \pm 0.35) \times 10^{-3}$	-
$\overline{K}_0^*(1430)^0 \rightarrow K^- \pi^+$		
$K^+ \overline{K}_2^*(1430)^0, \quad \overline{K}_2^* \rightarrow$	$(1.6 \pm 1.2) \times 10^{-4}$	-
$K^- \pi^+$		
$K^+ \overline{K}_0^*(700), \quad \overline{K}_0^* \rightarrow K^- \pi^+$	$(6.8 \pm 3.5) \times 10^{-4}$	-
$a_0(1450)^0 \pi^+, \quad a_0^0 \rightarrow$	$(4.5 \pm 7.0) \times 10^{-4}$	-
$K^+ K^-$		
$\phi(1680) \pi^+, \quad \phi \rightarrow K^+ K^-$	$(4.9 \pm 4.0) \times 10^{-5}$	-
$\phi \pi^+, \quad \phi \rightarrow K^+ K^-$	$(2.69 \pm 0.07) \times 10^{-3}$	647
$\phi \pi^+$		
$K^+ K^- \pi^+ \pi^0$	$(5.70 \pm 0.14) \times 10^{-3}$	647
$K_S^0 K_S^0 \pi^+$	$(6.62 \pm 0.32) \times 10^{-3}$	682
$K_S^0 K_S^0 \pi^+ \pi^0$	$(2.70 \pm 0.13) \times 10^{-3}$	741
$K_S^0 K^+ \eta$	$(1.34 \pm 0.21) \times 10^{-3}$	679
$K^+ K_S^0 \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	516
$K^+ K_S^0 \pi^+ \pi^-$	$(1.89 \pm 0.13) \times 10^{-3}$	678
$K_S^0 K^+ \pi^0 \pi^0$	$(5.8 \pm 1.3) \times 10^{-4}$	683
$K_S^0 K^- 2\pi^+$	$(2.27 \pm 0.13) \times 10^{-3}$	678
$K^+ K^- 2\pi^+ \pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$	601

A few poorly measured branching fractions:

$\phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$	619
$\phi \rho^+$	$< 1.5 \quad \% \quad \text{CL}=90\%$	260
$K^+ K^- \pi^+ \pi^0 \text{ non-}\phi$	$(1.5 \pm 0.7) \%$	682

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(2.08 \pm 0.21) \times 10^{-4} \quad S=1.4$	864
$K^+ \eta$	$(1.25 \pm 0.16) \times 10^{-4} \quad S=1.1$	776
$K^+ \eta'(958)$	$(1.85 \pm 0.20) \times 10^{-4}$	571
$K^+ 2\pi^0$	$(2.1 \pm 0.4) \times 10^{-4}$	847
$K^*(892)^+ \pi^0$	$(3.4 \pm 1.4) \times 10^{-4}$	714
$K^+ \pi^+ \pi^-$	$(4.91 \pm 0.09) \times 10^{-4}$	846
$K^+ \rho^0$	$(1.9 \pm 0.5) \times 10^{-4}$	679
$K^+ \eta \pi^0$	$(2.1 \pm 0.5) \times 10^{-4}$	726

$K^*(892)^+ \eta$	(4.4 \pm 1.8) $\times 10^{-4}$	586
$K^*(892)^0 \pi^+$, $K^*(892)^0 \rightarrow K^+ \pi^-$	(2.3 \pm 0.4) $\times 10^{-4}$	714
$K^+ f_0(980)$, $f_0(980) \rightarrow \pi^+ \pi^-$	(4.4 \pm 2.6) $\times 10^{-5}$	—
$K_2^*(1430)^0 \pi^+$, $K_2^*(1430)^0 \rightarrow K^+ \pi^-$	(3.9 \pm 2.7) $\times 10^{-5}$	—
$K^+ \pi^-$ nonresonant	not seen	846
$K^+ \pi^+ \pi^- \pi^0$	(1.21 \pm 0.09) $\times 10^{-3}$	817
$K^+ \pi^+ \pi^- \pi^0$ nonresonant	(1.10 \pm 0.07) $\times 10^{-3}$	817
$K^+ \omega$	(5.7 \pm 2.5) $\times 10^{-5}$	675
$2K^+ K^-$	(6.14 \pm 0.11) $\times 10^{-5}$	550
$\phi(1020)^0 K^+$	< 2.1 $\times 10^{-5}$ CL=90%	—
$K^+ \phi(1020)$, $\phi \rightarrow K^+ K^-$	(4.4 \pm 0.6) $\times 10^{-6}$	—
$K^+(K^+ K^-)$ <i>S-wave</i>	(5.77 \pm 0.12) $\times 10^{-5}$	550

$\Delta C = 1$ weak neutral current (*C1*) modes, or Lepton Family number (*LF*), or Lepton number (*L*), or Baryon number (*B*) violating modes

$\pi^+ e^+ e^-$	<i>C1</i>	< 1.1 $\times 10^{-6}$ CL=90%	930
$\pi^+ \pi^0 e^+ e^-$		< 1.4 $\times 10^{-5}$ CL=90%	925
$\pi^+ \phi$, $\phi \rightarrow e^+ e^-$	[g]	(1.7 \pm 1.4) $\times 10^{-6}$	—
$\pi^+ \mu^+ \mu^-$	<i>C1</i>	< 6.7 $\times 10^{-8}$ CL=90%	918
$\pi^+ \phi$, $\phi \rightarrow \mu^+ \mu^-$	[g]	(1.8 \pm 0.8) $\times 10^{-6}$	—
$\rho^+ \mu^+ \mu^-$	<i>C1</i>	< 5.6 $\times 10^{-4}$ CL=90%	757
$K^+ e^+ e^-$	[h]	< 8.5 $\times 10^{-7}$ CL=90%	870
$K^+ \pi^0 e^+ e^-$		< 1.5 $\times 10^{-5}$ CL=90%	864
$K_S^0 \pi^+ e^+ e^-$		< 2.6 $\times 10^{-5}$ CL=90%	—
$K_S^0 K^+ e^+ e^-$		< 1.1 $\times 10^{-5}$ CL=90%	792
$K^+ \mu^+ \mu^-$	[h]	< 5.4 $\times 10^{-8}$ CL=90%	856
$\pi^+ e^+ \mu^-$	<i>LF</i>	< 2.1 $\times 10^{-7}$ CL=90%	927
$\pi^+ e^- \mu^+$	<i>LF</i>	< 2.2 $\times 10^{-7}$ CL=90%	927
$K^+ e^+ \mu^-$	<i>LF</i>	< 7.5 $\times 10^{-8}$ CL=90%	866
$K^+ e^- \mu^+$	<i>LF</i>	< 1.0 $\times 10^{-7}$ CL=90%	866
$\pi^- 2e^+$	<i>L</i>	< 5.3 $\times 10^{-7}$ CL=90%	930
$\pi^- 2\mu^+$	<i>L</i>	< 1.4 $\times 10^{-8}$ CL=90%	918
$\pi^- e^+ \mu^+$	<i>L</i>	< 1.3 $\times 10^{-7}$ CL=90%	927
$\rho^- 2\mu^+$	<i>L</i>	< 5.6 $\times 10^{-4}$ CL=90%	757
$K^- 2e^+$	<i>L</i>	< 9 $\times 10^{-7}$ CL=90%	870
$K_S^0 \pi^- 2e^+$		< 3.3 $\times 10^{-6}$ CL=90%	863
$K^- \pi^0 2e^+$		< 8.5 $\times 10^{-6}$ CL=90%	864
$K^- 2\mu^+$	<i>L</i>	< 1.0 $\times 10^{-5}$ CL=90%	856
$K^- e^+ \mu^+$	<i>L</i>	< 1.9 $\times 10^{-6}$ CL=90%	866

$K^*(892)^- 2\mu^+$	L	< 8.5	$\times 10^{-4}$	CL=90%	703
Λe^+	L, B	< 1.1	$\times 10^{-6}$	CL=90%	602
$\bar{\Lambda} e^+$	L, B	< 6.5	$\times 10^{-7}$	CL=90%	602
$\Sigma^0 e^+$	L, B	< 1.7	$\times 10^{-6}$	CL=90%	554
$\bar{\Sigma}^0 e^+$	L, B	< 1.3	$\times 10^{-6}$	CL=90%	554
$\bar{n} e^+$		< 1.43	$\times 10^{-5}$	CL=90%	699
$n e^+$		< 2.91	$\times 10^{-5}$	CL=90%	699

D⁰

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1864.84 \pm 0.05$ MeV $m_{D^\pm} - m_{D^0} = 4.822 \pm 0.015$ MeVMean life $\tau = (410.3 \pm 1.0) \times 10^{-15}$ s $c\tau = 123.01$ μ m**Mixing and related parameters**

$$|m_{D_1^0} - m_{D_2^0}| = (0.997 \pm 0.116) \times 10^{10} \text{ } \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.394 \pm 0.056) \times 10^{-2}$$

$$|\mathbf{q}/\mathbf{p}| = 0.995 \pm 0.016$$

$$\Delta\Gamma = (0.089 \pm 0.113) \times 10^{-3}$$

$$\phi^{K_S^0 \pi \pi} = 0.02^{+0.04}_{-0.05}$$

$$K^+ \pi^- \text{ relative strong phase: } \cos \delta = 0.990 \pm 0.025$$

$$K^- \pi^+ \pi^0 \text{ coherence factor } R_{K^- \pi^+ \pi^0} = 0.792 \pm 0.033$$

$$K^- \pi^+ \pi^0 \text{ average relative strong phase } \delta^{K^- \pi^+ \pi^0} = (198 \pm 10)^\circ$$

$$K^- \pi^- 2\pi^+ \text{ coherence factor } R_{K^- \pi^- 2\pi^+} = 0.52^{+0.10}_{-0.09}$$

$$K^- \pi^- 2\pi^+ \text{ average relative strong phase } \delta^{K^- \pi^- 2\pi^+} = (149^{+26}_{-16})^\circ \quad (S = 1.4)$$

$$D^0 \rightarrow K^- \pi^- 2\pi^+, R_{K^- \pi^- 2\pi^+} (y \cos \delta^{K^- \pi^- 2\pi^+} - x \sin \delta^{K^- \pi^- 2\pi^+}) = (-3.0 \pm 0.7) \times 10^{-3} \text{ TeV}^{-1}$$

$$K_S^0 K^+ \pi^- \text{ coherence factor } R_{K_S^0 K^+ \pi^-} = 0.70 \pm 0.08$$

$$K_S^0 K^+ \pi^- \text{ average relative strong phase } \delta^{K_S^0 K^+ \pi^-} = (0 \pm 16)^\circ$$

$$K^* K \text{ coherence factor } R_{K^* K} = 0.94 \pm 0.12$$

$$K^* K \text{ average relative strong phase } \delta^{K^* K} = (-17 \pm 18)^\circ$$

CP-even fractions (labeled by the D⁰ decay)

$$CP\text{-even fraction in } D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0 \text{ decays} = (23.6 \pm 0.9)\%$$

$$CP\text{-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^0 \text{ decays} = (97.3 \pm 1.7)\%$$

$$CP\text{-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- \text{ decays} = (74.6 \pm 1.6)\% \quad (S = 1.2)$$

$$CP\text{-even fraction in } D^0 \rightarrow \pi^+ \pi^- 2\pi^0 \text{ decays} = 0.68 \pm 0.08$$

CP -even fraction in $D^0 \rightarrow 2\pi^+ 2\pi^- \pi^0$ decays = 0.44 ± 0.10
 CP -even fraction in $D^0 \rightarrow \pi^+ \pi^- 3\pi^0$ decays = $0.52^{+0.34}_{-0.27}$
 CP -even fraction in $D^0 \rightarrow 2\pi^+ 2\pi^- 2\pi^0$ decays = 0.79 ± 0.26
 CP -even fraction in $D^0 \rightarrow K^+ K^- \pi^0$ decays = $(73 \pm 6)\%$
 CP -even fraction in $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ decays = $(74.1 \pm 3.0)\%$

CP -violation decay-rate asymmetries (labeled by the D^0 decay)

$A_{CP}(K^+ K^-) = (4 \pm 5) \times 10^{-4}$
 $A_{CP}(2K_S^0) = (-1.9 \pm 1.1)\% \quad (S = 1.1)$
 $A_{CP}(\pi^+ \pi^-) = (0.13 \pm 0.14)\%$
 $A_{CP}(\pi^0 \pi^0) = (0.0 \pm 0.6)\%$
 $A_{CP}(\rho \gamma) = (6 \pm 15) \times 10^{-2}$
 $A_{CP}(\phi \gamma) = (-9 \pm 7) \times 10^{-2}$
 $A_{CP}(\overline{K}^*(892)^0 \gamma) = (-0.3 \pm 2.0) \times 10^{-2}$
 $A_{CP}(\pi^+ \pi^- \pi^0) = (0.4 \pm 0.4)\%$
 $A_{CP}(\eta \pi^+ \pi^-)$ in D^0 , $\overline{D}^0 \rightarrow \eta \pi^+ \pi^- = (0.9 \pm 1.3) \times 10^{-2}$
 $A_{CP}(\rho(770)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (1.2 \pm 0.9)\% [i]$
 $A_{CP}(\rho(770)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-3.1 \pm 3.0)\% [i]$
 $A_{CP}(\rho(770)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (-1.0 \pm 1.7)\% [i]$
 $A_{CP}(\rho(1450)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 70)\% [i]$
 $A_{CP}(\rho(1450)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-20 \pm 40)\% [i]$
 $A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 9)\% [i]$
 $A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (-5 \pm 14)\% [i]$
 $A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (13 \pm 9)\% [i]$
 $A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (8 \pm 11)\% [i]$
 $A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 35)\% [i]$
 $A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (25 \pm 18)\% [i]$
 $A_{CP}(f_0(1500) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 18)\% [i]$
 $A_{CP}(f_0(1710) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 24)\% [i]$
 $A_{CP}(f_2(1270) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-4 \pm 6)\% [i]$
 $A_{CP}(\sigma(400) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 8)\% [i]$
 $A_{CP}(\text{nonresonant } \pi^+ \pi^- \pi^0) = (-13 \pm 23)\% [i]$
 $A_{CP}(\pi^+ \pi^- 2\pi^0)$ in D^0 , $\overline{D}^0 \rightarrow \pi^+ \pi^- 2\pi^0 = (-2.5 \pm 2.0)\%$
 $A_{CP}(a_1(1260)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) = (5 \pm 6)\%$
 $A_{CP}(a_1(1260)^- \pi^+ \rightarrow 2\pi^+ 2\pi^-) = (14 \pm 18)\%$
 $A_{CP}(\pi(1300)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) = (-2 \pm 15)\%$
 $A_{CP}(\pi(1300)^- \pi^+ \rightarrow 2\pi^+ 2\pi^-) = (-6 \pm 30)\%$
 $A_{CP}(a_1(1640)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) = (9 \pm 26)\%$
 $A_{CP}(\pi_2(1670)^+ \pi^- \rightarrow 2\pi^+ 2\pi^-) = (7 \pm 18)\%$
 $A_{CP}(\sigma f_0(1370) \rightarrow 2\pi^+ 2\pi^-) = (-15 \pm 19)\%$
 $A_{CP}(\sigma \rho(770)^0 \rightarrow 2\pi^+ 2\pi^-) = (3 \pm 27)\%$

$$\begin{aligned}
A_{CP}(2\rho(770)^0 \rightarrow 2\pi^+ 2\pi^-) &= (-6 \pm 6)\% \\
A_{CP}(2f_2(1270) \rightarrow 2\pi^+ 2\pi^-) &= (-28 \pm 24)\% \\
A_{CP}(\pi^+ \pi^- \pi^0 \eta) \text{ in } D^0, \overline{D}^0 \rightarrow \pi^+ \pi^- \pi^0 \eta &= (-6 \pm 6) \times 10^{-2} \\
A_{CP}(K^+ K^- \pi^0) &= (-1.0 \pm 1.7)\% \\
A_{CP}(K^*(892)^+ K^- \rightarrow K^+ K^- \pi^0) &= (-0.9 \pm 1.3)\% [i] \\
A_{CP}(K^*(1410)^+ K^- \rightarrow K^+ K^- \pi^0) &= (-21 \pm 24)\% [i] \\
A_{CP}((K^+ \pi^0)_{S-wave} K^- \rightarrow K^+ K^- \pi^0) &= (7 \pm 15)\% [i] \\
A_{CP}(\phi(1020) \pi^0 \rightarrow K^+ K^- \pi^0) &= (1.1 \pm 2.2)\% [i] \\
A_{CP}(f_0(980) \pi^0 \rightarrow K^+ K^- \pi^0) &= (-3 \pm 19)\% [i] \\
A_{CP}(a_0(980)^0 \pi^0 \rightarrow K^+ K^- \pi^0) &= (-5 \pm 16)\% [i] \\
A_{CP}(f'_2(1525) \pi^0 \rightarrow K^+ K^- \pi^0) &= (0 \pm 160)\% [i] \\
A_{CP}(K^*(892)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-5 \pm 4)\% [i] \\
A_{CP}(K^*(1410)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-17 \pm 29)\% [i] \\
A_{CP}((K^- \pi^0)_{S-wave} K^+ \rightarrow K^+ K^- \pi^0) &= (-10 \pm 40)\% [i] \\
A_{CP}(K^+ K^- \eta) \text{ in } D^0, \overline{D}^0 \rightarrow K^+ K^- \eta &= (-1.4 \pm 3.5) \times 10^{-2} \\
A_{CP}(\phi(1020) \eta \rightarrow K^+ K^- \eta) \text{ in } D^0, \overline{D}^0 \rightarrow \phi(1020) \eta &= (-2 \pm 4) \times 10^{-2} \\
A_{CP}(K_S^0 \pi^0) &= (-0.20 \pm 0.17)\% \\
A_{CP}(K_S^0 \eta) &= (0.5 \pm 0.5)\% \\
A_{CP}(K_S^0 \eta') &= (1.0 \pm 0.7)\% \\
A_{CP}(K_S^0 \phi) &= (-3 \pm 9)\% \\
A_{CP}(K^- \pi^+) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+ \pi^-) &= (-0.9 \pm 1.4)\% \\
A_{CP}(D_{CP(\pm 1)} \rightarrow K^\mp \pi^\pm) &= (13.1 \pm 1.0)\% \\
A_{CP}(K^- \pi^+ \pi^0) &= (0.1 \pm 0.5)\% \\
A_{CP}(K^+ \pi^- \pi^0) &= (0 \pm 5)\% \\
A_{CP}(K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.8)\% \\
A_{CP}(K^\mp \pi^\pm \eta) \text{ in } D^0, \overline{D}^0 \rightarrow K^\mp \pi^\pm \eta &= (-1.9 \pm 1.6) \times 10^{-2} \\
A_{CP}(K_S^0 \pi^0 \eta) \text{ in } D^0, \overline{D}^0 \rightarrow K_S^0 \pi^0 \eta &= (-3.9 \pm 3.3) \times 10^{-2} \\
A_{CP}(K^\mp \pi^\pm \pi^0 \eta) \text{ in } D^0, \overline{D}^0 \rightarrow K^\mp \pi^\pm \pi^0 \eta &= (-8 \pm 5) \times 10^{-2} \\
A_{CP}(K^*(892)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (0.4 \pm 0.5)\% \\
A_{CP}(K^*(892)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (1 \pm 6)\% \\
A_{CP}(\overline{K}^0 \rho^0 \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.5)\% \\
A_{CP}(\overline{K}^0 \omega \rightarrow K_S^0 \pi^+ \pi^-) &= (-13 \pm 7)\% \\
A_{CP}(\overline{K}^0 f_0(980) \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.4 \pm 2.7)\% \\
A_{CP}(\overline{K}^0 f_2(1270) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 5)\% \\
A_{CP}(\overline{K}^0 f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-) &= (-1 \pm 9)\% \\
A_{CP}(\overline{K}^0 \rho^0(1450) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 10)\% \\
A_{CP}(\overline{K}^0 f_0(600) \rightarrow K_S^0 \pi^+ \pi^-) &= (-3 \pm 5)\% \\
A_{CP}(K^*(1410)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (-2 \pm 9)\% \\
A_{CP}(K_0^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (4 \pm 4)\% \\
A_{CP}(K_0^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (12 \pm 15)\%
\end{aligned}$$

$$\begin{aligned}
A_{CP}(K_2^*(1430)^-\pi^+ \rightarrow K_S^0\pi^+\pi^-) &= (3 \pm 6)\% \\
A_{CP}(K_2^*(1430)^+\pi^- \rightarrow K_S^0\pi^+\pi^-) &= (-10 \pm 32)\% \\
A_{CP}(K^-\pi^+\pi^+\pi^-) &= (0.2 \pm 0.5)\% \\
A_{CP}(K^+\pi^-\pi^+\pi^-) &= (-2 \pm 4)\% \\
A_{CP}(K^+K^-\pi^+\pi^-) &= (1.3 \pm 1.7)\% \\
A_{CP}(2K_S^0\pi^+\pi^-) \text{ in } D^0, \overline{D}^0 \rightarrow 2K_S^0\pi^+\pi^- &= (-2.5 \pm 1.4) \times 10^{-2} \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow K^+K^-\pi^+\pi^-) &= (-2.3 \pm 1.7)\% \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow K^{*0}\pi^+K^-) &= (-1 \pm 10)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow \overline{K}^{*0}\pi^-K^+) &= (-10 \pm 32)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow K^+K^-\pi^+\pi^-) &= (1.7 \pm 3.5)\% \\
A_{CP}(K_1^*(1270)^+K^- \rightarrow \rho^0K^+K^-) &= (-7 \pm 17)\% \\
A_{CP}(K_1^*(1270)^-K^+ \rightarrow \rho^0K^-K^+) &= (10 \pm 13)\% \\
A_{CP}(K_1(1400)^+K^- \rightarrow K^+K^-\pi^+\pi^-) &= (-4.4 \pm 2.1)\% \\
A_{CP}(K^*(1410)^+K^- \rightarrow K^{*0}\pi^+K^-) &= (-20 \pm 17)\% \\
A_{CP}(K^*(1410)^-K^+ \rightarrow \overline{K}^{*0}\pi^-K^+) &= (-1 \pm 14)\% \\
A_{CP}(K^*(1680)^+K^- \rightarrow K^+K^-\pi^+\pi^-) &= (-17 \pm 29)\% \\
A_{CP}(K^{*0}\overline{K}^{*0}) \text{ in } D^0, \overline{D}^0 \rightarrow K^{*0}\overline{K}^{*0} &= (-5 \pm 14)\% \\
A_{CP}(K^{*0}\overline{K}^{*0} \text{ S-wave}) &= (-3.9 \pm 2.2)\% \\
A_{CP}(\phi\rho^0) \text{ in } D^0, \overline{D}^0 \rightarrow \phi\rho^0 &= (1 \pm 9)\% \\
A_{CP}(\phi\rho^0 \text{ S-wave}) &= (-3 \pm 5)\% \\
A_{CP}(\phi\rho^0 \text{ D-wave}) &= (-37 \pm 19)\% \\
A_{CP}(\phi(\pi^+\pi^-)_{S\text{-wave}}) &= (6 \pm 6)\% \\
A_{CP}(K^*(892)^0(K^-\pi^+)_{S\text{-wave}}) &= (-10 \pm 40)\% \\
A_{CP}(K^+K^-\pi^+\pi^- \text{ non-resonant}) &= (8 \pm 20)\% \\
A_{CP}((K^-\pi^+)_{P\text{-wave}}(K^+\pi^-)_{S\text{-wave}}) &= (3 \pm 11)\% \\
A_{CP}(K^+K^-\mu^+\mu^-) \text{ in } D^0, \overline{D}^0 \rightarrow K^+K^-\mu^+\mu^- &= (-2 \pm 6)\% \\
A_{CP}(\pi^+\pi^-\mu^+\mu^-) \text{ in } D^0, \overline{D}^0 \rightarrow \pi^+\pi^-\mu^+\mu^- &= (2.9 \pm 2.1)\%
\end{aligned}$$

CP-violation asymmetry difference

$$\Delta A_{CP} = A_{CP}(K^+K^-) - A_{CP}(\pi^+\pi^-) = (-0.154 \pm 0.029)\%$$

χ^2 tests of CP-violation (CPV) p-values

$$\begin{aligned}
\text{Local CPV in } D^0, \overline{D}^0 \rightarrow \pi^+\pi^-\pi^0 &= 10.6\% \\
\text{Local CPV in } D^0, \overline{D}^0 \rightarrow \pi^+\pi^-\pi^+\pi^- &= (0.6 \pm 0.2)\% \\
\text{Local CPV in } D^0, \overline{D}^0 \rightarrow K_S^0\pi^+\pi^- &= 96\% \\
\text{Local CPV in } D^0, \overline{D}^0 \rightarrow K^+K^-\pi^0 &= 16.6\% \\
\text{Local CPV in } D^0, \overline{D}^0 \rightarrow K^+K^-\pi^+\pi^- &= 9.1\%
\end{aligned}$$

T-violation decay-rate asymmetry

$$\begin{aligned}
A_T(K^+K^-\pi^+\pi^-) &= (2.9 \pm 2.2) \times 10^{-3} [b] \\
A_{T\text{viol}}(2K_S^0\pi^+\pi^-) \text{ in } D^0, \overline{D}^0 \rightarrow 2K_S^0\pi^+\pi^- &= (-1.9 \pm 1.4) \times 10^{-2} \\
A_{T\text{viol}}(K_S^0\pi^+\pi^-\pi^0) \text{ in } D^0, \overline{D}^0 \rightarrow K_S^0\pi^+\pi^-\pi^0 &= \\
&\quad (-0.3^{+1.4}_{-1.6}) \times 10^{-3}
\end{aligned}$$

***CPT*-violation decay-rate asymmetry**

$$A_{CPT}(K^\mp\pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$\begin{aligned} r_V &\equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^-\ell^+\nu_\ell = 1.46 \pm 0.07 \\ r_2 &\equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^-\ell^+\nu_\ell = 0.68 \pm 0.06 \\ f_+(0) \text{ in } D^0 \rightarrow K^-\ell^+\nu_\ell &= 0.736 \pm 0.004 \\ f_+(0)|V_{cs}| \text{ in } D^0 \rightarrow K^-\ell^+\nu_\ell &= 0.7166 \pm 0.0030 \\ r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^-\ell^+\nu_\ell &= -2.40 \pm 0.16 \\ r_2 \equiv a_2/a_0 \text{ in } D^0 \rightarrow K^-\ell^+\nu_\ell &= 5 \pm 4 \\ f_+(0) \text{ in } D^0 \rightarrow \pi^-\ell^+\nu_\ell &= 0.637 \pm 0.009 \\ f_+(0)|V_{cd}| \text{ in } D^0 \rightarrow \pi^-\ell^+\nu_\ell &= 0.1436 \pm 0.0026 \quad (S = 1.5) \\ r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^-\ell^+\nu_\ell &= -1.97 \pm 0.28 \quad (S = 1.4) \\ r_2 \equiv a_2/a_0 \text{ in } D^0 \rightarrow \pi^-\ell^+\nu_\ell &= -0.2 \pm 2.2 \quad (S = 1.7) \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ ρ Confidence level(MeV/c)
Topological modes		
0-prongs	[j] (15 ± 6) %	—
2-prongs	(71 ± 6) %	—
4-prongs	[k] (14.6 ± 0.5) %	—
6-prongs	[l] (6.5 ± 1.3) × 10 ⁻⁴	—
Inclusive modes		
e^+ anything	[n] (6.49 ± 0.11) %	—
μ^+ anything	(6.8 ± 0.6) %	—
K^- anything	(54.7 ± 2.8) %	S=1.3
K_S^0 anything	(20.75 ± 0.23) %	—
K^+ anything	(3.4 ± 0.4) %	—
$K^*(892)^-$ anything	(15 ± 9) %	—
$\bar{K}^*(892)^0$ anything	(9 ± 4) %	—
$K^*(892)^+$ anything	< 3.6 %	CL=90%
$K^*(892)^0$ anything	(2.8 ± 1.3) %	—
η anything	(9.5 ± 0.9) %	—
η' anything	(2.48 ± 0.27) %	—
ϕ anything	(1.08 ± 0.04) %	—
$\pi^+\pi^+\pi^-$ anything	(17.60 ± 0.25) %	—
invisibles	< 9.4 × 10 ⁻⁵	CL=90%

Semileptonic modes

$K^- e^+ \nu_e$	(3.549 \pm 0.026) %	S=1.2	867
$K^- \mu^+ \nu_\mu$	(3.41 \pm 0.04) %		864
$K^*(892)^- e^+ \nu_e$	(2.15 \pm 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$	(1.89 \pm 0.24) %		714
$K^- \pi^0 e^+ \nu_e$	(1.6 \pm 1.3) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	(1.44 \pm 0.04) %		860
$(\bar{K}^0 \pi^-)_{S-wave} e^+ \nu_e$	(7.9 \pm 1.7) $\times 10^{-4}$		860
$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 \pm 1.4) $\times 10^{-4}$		843
$K_1(1270)^- e^+ \nu_e$	(1.01 \pm 0.18) $\times 10^{-3}$		511
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.3 $\times 10^{-3}$	CL=90%	821
$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.5 $\times 10^{-3}$	CL=90%	692
$\pi^- e^+ \nu_e$	(2.91 \pm 0.04) $\times 10^{-3}$		927
$\pi^- \mu^+ \nu_\mu$	(2.67 \pm 0.12) $\times 10^{-3}$	S=1.3	924
$\pi^- \pi^0 e^+ \nu_e$	(1.45 \pm 0.07) $\times 10^{-3}$		922
$\rho^- e^+ \nu_e$	(1.50 \pm 0.12) $\times 10^{-3}$	S=1.9	771
$\rho^- \mu^+ \nu_\mu$	(1.35 \pm 0.13) $\times 10^{-3}$		767
$a(980)^- e^+ \nu_e, a^- \rightarrow \eta \pi^-$	(1.33 \pm 0.34) $\times 10^{-4}$		–
$b_1(1235)^- e^+ \nu_e, b_1^- \rightarrow \omega \pi^-$	< 1.12 $\times 10^{-4}$	CL=90%	–

Hadronic modes with one \bar{K}

$K^- \pi^+$	(3.947 \pm 0.030) %	S=1.2	861
$K_S^0 \pi^0$	(1.240 \pm 0.022) %		860
$K_L^0 \pi^0$	(9.76 \pm 0.32) $\times 10^{-3}$		860
$K_L^0 \eta$	(4.34 \pm 0.16) $\times 10^{-3}$		772
$K_L^0 \eta'$	(8.12 \pm 0.35) $\times 10^{-3}$	S=1.3	565
$K_L^0 \omega$	(1.16 \pm 0.04) %		670
$K_S^0 \pi^+ \pi^-$	[c] (2.80 \pm 0.18) %	S=1.1	842
$K_S^0 \rho^0$	(6.3 \pm 0.6) $\times 10^{-3}$		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	(2.0 \pm 0.6) $\times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S-wave}$	(3.3 \pm 0.8) $\times 10^{-3}$		842
$K_S^0 f_0(980), f_0 \rightarrow \pi^+ \pi^-$	(1.20 \pm 0.40) $\times 10^{-3}$		549
$K_S^0 f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	(2.8 \pm 0.9) $\times 10^{-3}$		†
$K_S^0 f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	(9 \pm 10) $\times 10^{-5}$		262
$K^*(892)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	(1.64 \pm 0.14) %		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow K_S^0 \pi^-$	(2.67 \pm 0.40) $\times 10^{-3}$		378

$K_2^*(1430)^-\pi^+$, $K_2^{*-} \rightarrow K_S^0\pi^-$	$(3.4 \pm 1.0) \times 10^{-4}$	367
$K^*(1680)^-\pi^+$, $K^{*-} \rightarrow K_S^0\pi^-$	$(4.4 \pm 3.5) \times 10^{-4}$	46
$K^*(892)^+\pi^-$, $K^{*+} \rightarrow K_S^0\pi^+$	[o] $(1.13 \pm 0.34) \times 10^{-4}$	711
$K_0^*(1430)^+\pi^-$, $K_0^{*+} \rightarrow K_S^0\pi^+$	[o] $< 1.4 \times 10^{-5}$ CL=95%	—
$K_2^*(1430)^+\pi^-$, $K_2^{*+} \rightarrow K_S^0\pi^+$	[o] $< 3.4 \times 10^{-5}$ CL=95%	—
$K_S^0\pi^+\pi^-$ nonresonant	$(2.5 \pm 1.6) \times 10^{-4}$	842
$K^-\pi^+\pi^0$	[c] $(14.4 \pm 0.6) \%$	S=2.2 844
$K^-\rho^+$	$(11.2 \pm 0.7) \%$	675
$K^-\rho(1700)^+$, $\rho^+ \rightarrow \pi^+\pi^0$	$(8.2 \pm 1.8) \times 10^{-3}$	†
$K^*(892)^-\pi^+$, $K^*(892)^- \rightarrow K^-\pi^0$	$(2.31 \pm 0.20) \%$	711
$\bar{K}^*(892)^0\pi^0$, $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(1.95 \pm 0.25) \%$	711
$K_0^*(1430)^-\pi^+$, $K_0^{*-} \rightarrow K^-\pi^0$	$(4.8 \pm 2.2) \times 10^{-3}$	378
$\bar{K}_0^*(1430)^0\pi^0$, $\bar{K}_0^{*0} \rightarrow K^-\pi^+$	$(5.9 \pm 1.6) \times 10^{-3}$	379
$K^*(1680)^-\pi^+$, $K^{*-} \rightarrow K^-\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$	46
$K^-\pi^+\pi^0$ nonresonant	$(1.15 \pm 0.20) \%$	844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2 843
$K_L^0\pi^0\pi^0$	$(1.26 \pm 0.06) \%$	843
$K_S^0(2\pi^0)_{S-wave}$	$(2.6 \pm 0.7) \times 10^{-3}$	—
$\bar{K}^*(892)^0\pi^0$, $\bar{K}^{*0} \rightarrow K_S^0\pi^0$	$(8.1 \pm 0.7) \times 10^{-3}$	711
$\bar{K}^*(1430)^0\pi^0$, $\bar{K}^{*0} \rightarrow K_S^0\pi^0$	$(4 \pm 23) \times 10^{-5}$	—
$\bar{K}^*(1680)^0\pi^0$, $\bar{K}^{*0} \rightarrow K_S^0\pi^0$	$(1.0 \pm 0.4) \times 10^{-3}$	—
$K_S^0 f_2(1270)$, $f_2 \rightarrow 2\pi^0$	$(2.3 \pm 1.1) \times 10^{-4}$	—
$2K_S^0$, one $K_S^0 \rightarrow 2\pi^0$	$(3.2 \pm 1.1) \times 10^{-4}$	—
$K_S^0 3\pi^0$	$(7.6 \pm 0.4) \times 10^{-3}$	815
$K^-2\pi^+\pi^-$	[c] $(8.22 \pm 0.14) \%$	813
$K^-\pi^+\rho^0$ total	$(6.87 \pm 0.31) \%$	609
$K^-\pi^+\rho^0$ 3-body	$(6.1 \pm 1.6) \times 10^{-3}$	609
$\bar{K}^*(892)^0\rho^0$, $\bar{K}^{*0} \rightarrow K^-\pi^+$	$(1.01 \pm 0.05) \%$	416

$\bar{K}^*(892)^0 \rho^0$ transverse, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(1.2 ± 0.4) %	417
$K^- a_1(1260)^+, a_1^+ \rightarrow \rho^0 \pi^+$	(4.32 ± 0.32) %	327
$K_1(1270)^- \pi^+, K_1^- \rightarrow K^- \pi^+ \pi^-$ total	(3.9 ± 0.4) × 10 ⁻³	—
$K_1(1270)^- \pi^+, K_1^- \rightarrow \bar{K}^*(892)^0 \pi^-, \bar{K}^{*0} \rightarrow K^- \pi^+$	(6.6 ± 2.3) × 10 ⁻⁴	484
$K^- 2\pi^+ \pi^-$ nonresonant	(1.81 ± 0.07) %	813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] (5.2 ± 0.6) %	813
$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	(1.17 ± 0.03) × 10 ⁻³	772
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	(9.9 ± 0.6) × 10 ⁻³	670
$K^- \pi^+ 2\pi^0$	(8.86 ± 0.23) %	815
$K^- \pi^+ 3\pi^0$	(9.5 ± 0.4) × 10 ⁻³	774
$K_S^0 \pi^+ \pi^- 2\pi^0$	(1.27 ± 0.06) %	773
$K^- 2\pi^+ \pi^- \pi^0$	(4.3 ± 0.4) %	771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	(1.3 ± 0.6) %	643
$\bar{K}^*(892)^0 \omega, \bar{K}^{*0} \rightarrow K^- \pi^+, \omega \rightarrow \pi^+ \pi^- \pi^0$	(6.5 ± 3.0) × 10 ⁻³	410
$K^- \pi^+ \omega$	(3.39 ± 0.10) %	605
$\bar{K}^*(892)^0 \omega$	(1.1 ± 0.5) %	410
$K_S^0 \pi^0 \omega$	(8.5 ± 0.6) × 10 ⁻³	605
$K_S^0 \eta \pi^0$	(1.01 ± 0.05) %	721
$K_S^0 a_0(980), a_0 \rightarrow \eta \pi^0$	(1.20 ± 0.28) %	—
$\bar{K}^*(892)^0 \eta, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$	(2.9 ± 0.7) × 10 ⁻³	—
$K^- \pi^+ \eta$	(1.88 ± 0.05) %	S=1.4 721
$K^*(892)^0 \eta, K^{*0} \rightarrow K^- \pi^+$	(8.9 ± 0.8) × 10 ⁻³	—
$a_0(980)^+ K^-, a_0^+ \rightarrow \eta \pi^+$	(7.4 ± 0.9) × 10 ⁻³	—
$K_2^*(1980)^- \pi^+, K_2^{*-} \rightarrow K^- \eta$	(2.2 ± 1.7) × 10 ⁻⁴	—
$K^- \pi^+ \pi^0 \eta$	(4.49 ± 0.27) × 10 ⁻³	656
$K_S^0 \pi^+ \pi^- \eta$	(2.80 ± 0.21) × 10 ⁻³	651
$K_S^0 2\pi^0 \eta$	(1.76 ± 0.26) × 10 ⁻³	656
$K_S^0 2\pi^+ 2\pi^-$	(2.66 ± 0.30) × 10 ⁻³	768
$K_S^0 \rho^0 \pi^+ \pi^-, \text{no } K^*(892)^-$	(1.1 ± 0.7) × 10 ⁻³	—
$K^*(892)^- 2\pi^+ \pi^-, K^*(892)^- \rightarrow K_S^0 \pi^-, \text{no } \rho^0$	(5 ± 7) × 10 ⁻⁴	642

$K^*(892)^-\rho^0\pi^+$	$(1.6 \pm 0.6) \times 10^{-3}$	230
$K^*(892)^-\rightarrow K_S^0\pi^-$		
$K_S^0 2\pi^+ 2\pi^-$ nonresonant	$< 1.2 \times 10^{-3}$	CL=90% 768
$K^- 3\pi^+ 2\pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	713

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. These nine modes below are all corrected for unseen decays of the resonances.

$K_S^0\eta$	$(5.09 \pm 0.13) \times 10^{-3}$	772
$K_S^0\omega$	$(1.11 \pm 0.06) \%$	670
$K_S^0\eta'(958)$	$(9.49 \pm 0.32) \times 10^{-3}$	565
$\bar{K}^*(892)^0\pi^+\pi^-\pi^0$	$(1.9 \pm 0.9) \%$	643
$\bar{K}^*(892)^0\eta$	$(1.41 \pm 0.12) \%$	583
$K^-\pi^+\eta'(958)$	$(6.43 \pm 0.34) \times 10^{-3}$	479
$K_S^0\eta'(958)\pi^0$	$(2.52 \pm 0.27) \times 10^{-3}$	479
$\bar{K}^*(892)^0\eta'(958)$	$< 1.0 \times 10^{-3}$	CL=90% 119

Hadronic modes with three K 's

$K_S^0 K^+ K^-$	$(4.42 \pm 0.32) \times 10^{-3}$	544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	$(2.9 \pm 0.4) \times 10^{-3}$	—
$K^- a_0(980)^+, a_0^+ \rightarrow$	$(5.9 \pm 1.8) \times 10^{-4}$	—
$K^+ K_S^0$		
$K^+ a_0(980)^-, a_0^- \rightarrow$	$< 1.1 \times 10^{-4}$	CL=95% —
$K^- K_S^0$		
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	$< 9 \times 10^{-5}$	CL=95% —
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	$(2.03 \pm 0.15) \times 10^{-3}$	520
$K_L^0 \phi$	$(4.14 \pm 0.23) \times 10^{-3}$	521
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	$(1.7 \pm 1.1) \times 10^{-4}$	—
$3K_S^0$	$(7.5 \pm 0.7) \times 10^{-4}$	S=1.4 539
$K^+ 2K^- \pi^+$	$(2.25 \pm 0.32) \times 10^{-4}$	434
$K^+ K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$	$(4.5 \pm 1.8) \times 10^{-5}$	†
$K^- \pi^+$		
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.7) \times 10^{-5}$	422
$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-,$	$(1.08 \pm 0.21) \times 10^{-4}$	†
$\bar{K}^{*0} \rightarrow K^- \pi^+$		
$K^+ 2K^- \pi^+$ nonresonant	$(3.4 \pm 1.5) \times 10^{-5}$	434
$2K_S^0 K^\pm \pi^\mp$	$(5.9 \pm 1.3) \times 10^{-4}$	427

Pionic modes

$\pi^+ \pi^-$	$(1.454 \pm 0.024) \times 10^{-3}$	S=1.4	922
$2\pi^0$	$(8.26 \pm 0.25) \times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	$(1.49 \pm 0.07) \%$	S=2.3	907
$\rho^+ \pi^-$	$(1.01 \pm 0.05) \%$		764
$\rho^0 \pi^0$	$(3.86 \pm 0.24) \times 10^{-3}$		764

$\rho^- \pi^+$	$(5.15 \pm 0.26) \times 10^{-3}$	764
$\rho(1450)^+ \pi^-$, $\rho^+ \rightarrow \pi^+ \pi^0$	$(1.6 \pm 2.1) \times 10^{-5}$	—
$\rho(1450)^0 \pi^0$, $\rho^0 \rightarrow \pi^+ \pi^-$	$(4.5 \pm 2.0) \times 10^{-5}$	—
$\rho(1450)^- \pi^+$, $\rho^- \rightarrow \pi^- \pi^0$	$(2.7 \pm 0.4) \times 10^{-4}$	—
$\rho(1700)^+ \pi^-$, $\rho^+ \rightarrow \pi^+ \pi^0$	$(6.1 \pm 1.5) \times 10^{-4}$	—
$\rho(1700)^0 \pi^0$, $\rho^0 \rightarrow \pi^+ \pi^-$	$(7.4 \pm 1.8) \times 10^{-4}$	—
$\rho(1700)^- \pi^+$, $\rho^- \rightarrow \pi^- \pi^0$	$(4.8 \pm 1.1) \times 10^{-4}$	—
$f_0(980) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	$(3.7 \pm 0.9) \times 10^{-5}$	—
$f_0(500) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	$(1.22 \pm 0.22) \times 10^{-4}$	—
$f_0(1370) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	$(5.5 \pm 2.1) \times 10^{-5}$	—
$f_0(1500) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	$(5.8 \pm 1.6) \times 10^{-5}$	—
$f_0(1710) \pi^0$, $f_0 \rightarrow \pi^+ \pi^-$	$(4.6 \pm 1.6) \times 10^{-5}$	—
$f_2(1270) \pi^0$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.97 \pm 0.21) \times 10^{-4}$	—
$\pi^+ \pi^- \pi^0$ nonresonant	$(1.3 \pm 0.4) \times 10^{-4}$	907
$3\pi^0$	$(2.0 \pm 0.5) \times 10^{-4}$	908
$2\pi^+ 2\pi^-$	$(7.56 \pm 0.20) \times 10^{-3}$	880
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow 2\pi^+ \pi^-$ total	$(4.53 \pm 0.31) \times 10^{-3}$	—
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+ S\text{-wave}$	$(3.13 \pm 0.21) \times 10^{-3}$	—
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+ D\text{-wave}$	$(1.9 \pm 0.5) \times 10^{-4}$	—
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	$(6.4 \pm 0.7) \times 10^{-4}$	—
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \rho^0 \pi^- S\text{-wave}$	$(2.3 \pm 0.9) \times 10^{-4}$	—
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \sigma \pi^-$	$(6.0 \pm 3.4) \times 10^{-5}$	—
$\pi(1300)^+ \pi^-$, $\pi(1300)^+ \rightarrow \sigma \pi^+$	$(5.1 \pm 2.7) \times 10^{-4}$	—
$\pi(1300)^- \pi^+$, $\pi(1300)^- \rightarrow \sigma \pi^-$	$(2.3 \pm 2.2) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \rho^0 \pi^+ D\text{-wave}$	$(3.2 \pm 1.6) \times 10^{-4}$	—
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	$(1.8 \pm 1.4) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow f_2(1270)^0 \pi^+$	$(2.0 \pm 0.9) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow \sigma \pi^-$	$(2.6 \pm 1.0) \times 10^{-4}$	—
$2\rho^0$ total	$(1.85 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.3 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.8 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.27 \pm 0.10) \times 10^{-3}$	—
$2\rho(770)^0$, $S\text{-wave}$	$(1.8 \pm 1.3) \times 10^{-4}$	—

$2\rho(770)^0$, <i>P</i> -wave	(5.3 \pm 1.3) $\times 10^{-4}$	–
$2\rho(770)^0$, <i>D</i> -wave	(6.2 \pm 3.0) $\times 10^{-4}$	–
Resonant $(\pi^+\pi^-)\pi^+\pi^-$	(1.51 \pm 0.12) $\times 10^{-3}$	–
3-body total		
$\sigma\pi^+\pi^-$	(6.2 \pm 0.9) $\times 10^{-4}$	–
$\sigma\rho(770)^0$	(5.0 \pm 2.5) $\times 10^{-4}$	–
$f_0(980)\pi^+\pi^-$, $f_0 \rightarrow \pi^+\pi^-$	(1.8 \pm 0.5) $\times 10^{-4}$	–
$f_2(1270)\pi^+\pi^-$, $f_2 \rightarrow \pi^+\pi^-$	(3.7 \pm 0.6) $\times 10^{-4}$	–
$2f_2(1270)$, $f_2 \rightarrow \pi^+\pi^-$	(1.6 \pm 1.8) $\times 10^{-4}$	–
$f_0(1370)\sigma$, $f_0 \rightarrow \pi^+\pi^-$	(1.6 \pm 0.5) $\times 10^{-3}$	–
$\pi^+\pi^-2\pi^0$	(1.002 \pm 0.031) %	882
$4\pi^0$	(7.6 \pm 1.1) $\times 10^{-4}$	883
$\eta\pi^0$	[q] (6.3 \pm 0.6) $\times 10^{-4}$	S=1.1 846
$\omega\pi^0$	[q] (1.17 \pm 0.35) $\times 10^{-4}$	761
$\omega\eta$	(1.98 \pm 0.18) $\times 10^{-3}$	S=1.1 648
$2\pi^+2\pi^-\pi^0$	(3.46 \pm 0.21) $\times 10^{-3}$	844
$\pi^+\pi^-3\pi^0$	(1.53 \pm 0.21) $\times 10^{-3}$	847
$2\pi^+2\pi^-2\pi^0$	(4.8 \pm 0.4) $\times 10^{-3}$	798
$\eta\pi^+\pi^-$	[q] (1.16 \pm 0.07) $\times 10^{-3}$	827
$\omega\pi^+\pi^-$	[q] (1.33 \pm 0.20) $\times 10^{-3}$	738
$\omega\pi^0\pi^0$	< 1.10 $\times 10^{-3}$	CL=90% 740
$\eta2\pi^0$	(3.8 \pm 1.3) $\times 10^{-4}$	829
$\pi^+\pi^-\pi^0\eta$	(3.23 \pm 0.22) $\times 10^{-3}$	797
$\eta3\pi^0$	(2.36 \pm 0.28) $\times 10^{-3}$	799
$\eta2\pi^+2\pi^-$	(6.0 \pm 1.2) $\times 10^{-4}$	751
$3\pi^+3\pi^-$	(4.3 \pm 1.2) $\times 10^{-4}$	795
$\eta'(958)\pi^0$	(9.2 \pm 1.0) $\times 10^{-4}$	678
$\eta'(958)\pi^+\pi^-$	(4.5 \pm 1.7) $\times 10^{-4}$	650
2η	(2.11 \pm 0.19) $\times 10^{-3}$	S=2.2 754
$2\eta\pi^0$	(7.3 \pm 2.2) $\times 10^{-4}$	699
$2\eta\pi^+\pi^-$	(8.5 \pm 1.4) $\times 10^{-4}$	623
3η	< 1.3 $\times 10^{-4}$	CL=90% 421
$\eta\eta'(958)$	(1.01 \pm 0.19) $\times 10^{-3}$	537

Hadronic modes with a $K\bar{K}$ pair

K^+K^-	(4.08 \pm 0.06) $\times 10^{-3}$	S=1.6 791
$2K_S^0$	(1.41 \pm 0.05) $\times 10^{-4}$	S=1.1 789
$K_S^0 K^- \pi^+$	(3.3 \pm 0.5) $\times 10^{-3}$	S=1.1 739
$\bar{K}^*(892)^0 K_S^0$, $\bar{K}^{*0} \rightarrow K^- \pi^+$	(8.2 \pm 1.6) $\times 10^{-5}$	608
$K^*(892)^+ K^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	(1.89 \pm 0.30) $\times 10^{-3}$	–

$\bar{K}^*(1410)^0 K_S^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(1.3 \pm 1.9) \times 10^{-4}$	-
$K^*(1410)^+ K^-, K^{*+} \rightarrow K_S^0 \pi^+$	$(3.2 \pm 1.9) \times 10^{-4}$	-
$(K^- \pi^+)_{S-wave} K_S^0$	$(6.0 \pm 2.9) \times 10^{-4}$	739
$(K_S^0 \pi^+)_{S-wave} K^-$	$(3.9 \pm 1.0) \times 10^{-4}$	739
$a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-$	$(1.3 \pm 1.4) \times 10^{-4}$	-
$a_0(1450)^- \pi^+, a_0^- \rightarrow K_S^0 K^-$	$(2.5 \pm 2.0) \times 10^{-5}$	-
$a_2(1320)^- \pi^+, a_2^- \rightarrow K_S^0 K^-$	$(5 \pm 5) \times 10^{-6}$	-
$\rho(1450)^- \pi^+, \rho^- \rightarrow K_S^0 K^-$	$(4.6 \pm 2.5) \times 10^{-5}$	-
$K_S^0 K^+ \pi^-$	$(2.17 \pm 0.35) \times 10^{-3}$	S=1.1 739
$K^*(892)^0 K_S^0, K^{*0} \rightarrow K^+ \pi^-$	$(1.12 \pm 0.21) \times 10^{-4}$	608
$K^*(892)^- K^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(6.2 \pm 1.1) \times 10^{-4}$	-
$K^*(1410)^0 K_S^0, K^{*0} \rightarrow K^+ \pi^+$	$(5 \pm 8) \times 10^{-5}$	-
$K^*(1410)^- K^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(2.6 \pm 2.0) \times 10^{-4}$	-
$(K^+ \pi^-)_{S-wave} K_S^0$	$(3.7 \pm 1.9) \times 10^{-4}$	739
$(K_S^0 \pi^-)_{S-wave} K^+$	$(1.4 \pm 0.6) \times 10^{-4}$	739
$a_0(980)^+ \pi^-, a_0^+ \rightarrow K_S^0 K^+$	$(6 \pm 4) \times 10^{-4}$	-
$a_0(1450)^+ \pi^-, a_0^+ \rightarrow K_S^0 K^+$	$(3.2 \pm 2.5) \times 10^{-5}$	-
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow K_S^0 K^+$	$(1.1 \pm 0.6) \times 10^{-5}$	-
$K^+ K^- \pi^0$	$(3.42 \pm 0.15) \times 10^{-3}$	743
$K^*(892)^+ K^-, K^*(892)^+ \rightarrow K^+ \pi^0$	$(1.52 \pm 0.08) \times 10^{-3}$	-
$K^*(892)^- K^+, K^*(892)^- \rightarrow K^- \pi^0$	$(5.4 \pm 0.4) \times 10^{-4}$	-
$(K^+ \pi^0)_{S-wave} K^-$	$(2.43 \pm 0.18) \times 10^{-3}$	743
$(K^- \pi^0)_{S-wave} K^+$	$(1.3 \pm 0.5) \times 10^{-4}$	743
$f_0(980) \pi^0, f_0 \rightarrow K^+ K^-$	$(3.6 \pm 0.6) \times 10^{-4}$	-
$\phi \pi^0, \phi \rightarrow K^+ K^-$	$(6.6 \pm 0.4) \times 10^{-4}$	-
$2K_S^0 \pi^0$	$< 1.45 \times 10^{-4}$	CL=90% 740
$K^+ K^- \eta$	$(5.9 \pm 1.9) \times 10^{-5}$	514
$\phi(1020) \eta$	$(1.84 \pm 0.12) \times 10^{-4}$	489
$K^+ K^- \eta$ nonresonant	$(9.9 \pm 0.9) \times 10^{-5}$	514
$2K_S^0 \eta$	$(1.3 \pm 0.6) \times 10^{-4}$	508
$K^+ K^- \pi^0 \pi^0$	$(6.9 \pm 0.8) \times 10^{-4}$	681

$K^+ K^- \pi^+ \pi^-$	$(2.47 \pm 0.11) \times 10^{-3}$	677
$\phi(\pi^+ \pi^-)_{S-wave}, \phi \rightarrow K^+ K^-$	$(10 \pm 5) \times 10^{-5}$	614
$(\phi\rho^0)_{S-wave}, \phi \rightarrow K^+ K^-$	$(6.9 \pm 0.6) \times 10^{-4}$	250
$(\phi\rho^0)_{P-wave}, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.9) \times 10^{-5}$	—
$(\phi\rho^0)_{D-wave}, \phi \rightarrow K^+ K^-$	$(4.2 \pm 1.4) \times 10^{-5}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{S-wave}, K^{*0} \rightarrow K^\pm \pi^\mp$	$(2.24 \pm 0.13) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{P-wave}, K^* \rightarrow K^\pm \pi^\mp$	$(1.20 \pm 0.08) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{D-wave}, K^* \rightarrow K^\pm \pi^\mp$	$(4.7 \pm 0.4) \times 10^{-5}$	—
$K^*(892)^0 (K^- \pi^+)_{S-wave}$	$(1.4 \pm 0.6) \times 10^{-4}$	—
3-body, $K^{*0} \rightarrow K^+ \pi^-$		
$K_1(1270)^+ K^-, K_1^+ \rightarrow K^{*0} \pi^+$	$(1.4 \pm 0.9) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(1.5 \pm 0.5) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow \rho^0 K^+$	$(2.2 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-, K_1^+ \rightarrow \omega(782) K^+, \omega \rightarrow \pi^+ \pi^-$	$(1.5 \pm 1.2) \times 10^{-5}$	—
$K_1(1270)^- K^+, K_1^- \rightarrow \rho^0 K^-$	$(1.3 \pm 0.4) \times 10^{-4}$	—
$K_1(1400)^+ K^-, K_1^+ \rightarrow K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(4.6 \pm 0.4) \times 10^{-4}$	—
$K^*(1410)^- K^+, K^{*-} \rightarrow \bar{K}^{*0} \pi^-$	$(7.0 \pm 1.1) \times 10^{-5}$	—
$K_1(1680)^+ K^-, K_1^+ \rightarrow K^{*0} \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(8.9 \pm 3.2) \times 10^{-5}$	—
$K^+ K^- \pi^+ \pi^-$ non-resonant	$(2.7 \pm 0.6) \times 10^{-4}$	—
$2K_S^0 \pi^+ \pi^-$	$(5.3 \pm 0.9) \times 10^{-4}$	673
$K_S^0 K^- \pi^+ \pi^0$	$(1.32 \pm 0.16) \times 10^{-3}$	677
$K_S^0 K^+ \pi^- \pi^0$	$(6.5 \pm 0.7) \times 10^{-4}$	677
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.4 \times 10^{-4}$	CL=90% 595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi \pi^0$	$(1.17 \pm 0.04) \times 10^{-3}$	645
$\phi \eta$	$(1.8 \pm 0.5) \times 10^{-4}$	489
$\phi \omega$	$(6.5 \pm 1.0) \times 10^{-4}$	238

Radiative modes					
$\rho^0 \gamma$		$(1.82 \pm 0.32) \times 10^{-5}$			771
$\omega \gamma$		$< 2.4 \times 10^{-4}$	CL=90%	768	
$\phi \gamma$		$(2.81 \pm 0.19) \times 10^{-5}$			654
$\overline{K}^*(892)^0 \gamma$		$(4.1 \pm 0.7) \times 10^{-4}$			719
Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes					
$K^+ \ell^- \bar{\nu}_\ell$ via \overline{D}^0	[r] < 2.2	$\times 10^{-5}$	CL=90%	—	
K^+ or $K^*(892)^+ e^- \bar{\nu}_e$ via \overline{D}^0	< 6	$\times 10^{-5}$	CL=90%	—	
$K^+ \pi^-$	DC	$(1.50 \pm 0.07) \times 10^{-4}$	S=3.0	861	
$K^+ \pi^-$ via DCS		$(1.363 \pm 0.025) \times 10^{-4}$		—	
$K^+ \pi^-$ via \overline{D}^0		$< 1.6 \times 10^{-5}$	CL=95%	861	
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \overline{D}^0$		$< 1.8 \times 10^{-4}$	CL=95%	—	
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$	DC	$(1.13 \pm 0.60) \times 10^{-4}$		711	
$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow K_S^0 \pi^+$		< 1.4 $\times 10^{-5}$		—	
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow K_S^0 \pi^+$		< 3.4 $\times 10^{-5}$		—	
$K^+ \pi^- \pi^0$	DC	$(3.06 \pm 0.16) \times 10^{-4}$	S=1.4	844	
$K^+ \pi^- \pi^0$ via \overline{D}^0		$(7.6 \pm 0.5) \times 10^{-4}$		—	
$K^+ \pi^- 2\pi^0$		$< 3.6 \times 10^{-4}$	CL=90%	815	
$K^+ \pi^+ 2\pi^-$ via DCS		$(2.49 \pm 0.07) \times 10^{-4}$		—	
$K^+ \pi^+ 2\pi^-$	DC	$(2.65 \pm 0.06) \times 10^{-4}$		813	
$K^+ \pi^+ 2\pi^-$ via \overline{D}^0		$(7.9 \pm 3.0) \times 10^{-6}$		812	
μ^- anything via \overline{D}^0		$< 4 \times 10^{-4}$	CL=90%	—	
$\Delta C = 1$ weak neutral current (C1) modes, Lepton Family number (LF) violating modes, Lepton (L) or Baryon (B) number violating modes					
$\gamma \gamma$	C1	< 8.5 $\times 10^{-7}$	CL=90%	932	
$e^+ e^-$	C1	< 7.9 $\times 10^{-8}$	CL=90%	932	
$\mu^+ \mu^-$	C1	< 3.1 $\times 10^{-9}$	CL=90%	926	
$\pi^0 e^+ e^-$	C1	< 4 $\times 10^{-6}$	CL=90%	928	
$\pi^0 \mu^+ \mu^-$	C1	< 1.8 $\times 10^{-4}$	CL=90%	915	
$\pi^0 \nu \bar{\nu}$		< 2.1 $\times 10^{-4}$	CL=90%	928	
$\eta e^+ e^-$	C1	< 3 $\times 10^{-6}$	CL=90%	852	
$\eta \mu^+ \mu^-$	C1	< 5.3 $\times 10^{-4}$	CL=90%	838	
$\pi^+ \pi^- e^+ e^-$	C1	< 7 $\times 10^{-6}$	CL=90%	922	
$\rho^0 e^+ e^-$	C1	< 1.0 $\times 10^{-4}$	CL=90%	771	
$\pi^+ \pi^- \mu^+ \mu^-$	C1	$(9.6 \pm 1.2) \times 10^{-7}$		894	
$\pi^+ \pi^- \mu^+ \mu^-$ (non-res)		< 5.5 $\times 10^{-7}$	CL=90%	—	

$\rho^0 \mu^+ \mu^-$	$C1$	< 2.2	$\times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	$C1$	< 6	$\times 10^{-6}$	CL=90%	768
$\omega \mu^+ \mu^-$	$C1$	< 8.3	$\times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	$C1$	< 1.1	$\times 10^{-5}$	CL=90%	791
$\phi e^+ e^-$	$C1$	< 5.2	$\times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	$C1$	$(1.54 \pm 0.32) \times 10^{-7}$			710
$K^- K^+ \mu^+ \mu^-$ (non-res)		< 3.3	$\times 10^{-5}$	CL=90%	—
$\phi \mu^+ \mu^-$	$C1$	< 3.1	$\times 10^{-5}$	CL=90%	631
$\bar{K}^0 e^+ e^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90%	866
$\bar{K}^0 \mu^+ \mu^-$		$[h] < 2.6$	$\times 10^{-4}$	CL=90%	852
$K^- \pi^+ e^+ e^-$, $675 < m_{ee} < 875$ MeV		$(4.0 \pm 0.5) \times 10^{-6}$			—
$K^- \pi^+ e^+ e^-$, $1.005 < m_{ee} < 1.035$ GeV		< 5	$\times 10^{-7}$	CL=90%	—
$\bar{K}^*(892)^0 e^+ e^-$		$[h] < 4.7$	$\times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	$C1$	< 3.59	$\times 10^{-4}$	CL=90%	829
$K^- \pi^+ \mu^+ \mu^-$, $675 < m_{\mu\mu} < 875$ MeV		$(4.2 \pm 0.4) \times 10^{-6}$			—
$\bar{K}^*(892)^0 \mu^+ \mu^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	$C1$	< 8.1	$\times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	LF	$[s] < 1.3$	$\times 10^{-8}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	LF	$[s] < 8.0$	$\times 10^{-7}$	CL=90%	924
$\eta e^\pm \mu^\mp$	LF	$[s] < 2.25$	$\times 10^{-6}$	CL=90%	848
$\pi^+ \pi^- e^\pm \mu^\mp$	LF	$[s] < 1.71$	$\times 10^{-6}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	LF	$[s] < 5.0$	$\times 10^{-7}$	CL=90%	767
$\omega e^\pm \mu^\mp$	LF	$[s] < 1.71$	$\times 10^{-6}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	LF	$[s] < 1.00$	$\times 10^{-6}$	CL=90%	754
$\phi e^\pm \mu^\mp$	LF	$[s] < 5.1$	$\times 10^{-7}$	CL=90%	648
$\bar{K}^0 e^\pm \mu^\mp$	LF	$[s] < 1.74$	$\times 10^{-6}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	LF	$[s] < 1.90$	$\times 10^{-6}$	CL=90%	848
$\bar{K}^*(892)^0 e^\pm \mu^\mp$	LF	$[s] < 1.25$	$\times 10^{-6}$	CL=90%	714
$2\pi^- 2e^+$	L	< 9.1	$\times 10^{-7}$	CL=90%	922
$2\pi^- 2\mu^+$	L	< 1.52	$\times 10^{-6}$	CL=90%	894
$K^- \pi^- 2e^+$	L	< 5.0	$\times 10^{-7}$	CL=90%	861
$K^- \pi^- 2\mu^+$	L	< 5.3	$\times 10^{-7}$	CL=90%	829
$2K^- 2e^+$	L	< 3.4	$\times 10^{-7}$	CL=90%	791
$2K^- 2\mu^+$	L	< 1.0	$\times 10^{-7}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+$	L	< 3.06	$\times 10^{-6}$	CL=90%	911
$K^- \pi^- e^+ \mu^+$	L	< 2.10	$\times 10^{-6}$	CL=90%	848
$2K^- e^+ \mu^+$	L	< 5.8	$\times 10^{-7}$	CL=90%	754
$p e^-$	L, B	< 2.2	$\times 10^{-6}$	CL=90%	696
$\bar{p} e^+$	L, B	< 1.2	$\times 10^{-6}$	CL=90%	696

$D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 2006.85 \pm 0.05$ MeV ($S = 1.1$)

$m_{D^{*0}} - m_{D^0} = 142.014 \pm 0.030$ MeV ($S = 1.5$)

Full width $\Gamma < 2.1$ MeV, CL = 90%

$\overline{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$D^0\pi^0$	(64.7 \pm 0.9) %		43
$D^0\gamma$	(35.3 \pm 0.9) %		137
$D^0e^+e^-$	(3.91 \pm 0.33) $\times 10^{-3}$		137
$\mu^+\mu^-$	< 2.5 $\times 10^{-8}$	90%	998
e^+e^-	< 1.7 $\times 10^{-6}$	90%	1003

 $D^*(2010)^{\pm}$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 2010.26 \pm 0.05$ MeV

$m_{D^*(2010)^+} - m_{D^+} = 140.603 \pm 0.015$ MeV

$m_{D^*(2010)^+} - m_{D^0} = 145.4258 \pm 0.0017$ MeV

Full width $\Gamma = 83.4 \pm 1.8$ keV

$D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^{\pm}$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0\pi^+$	(67.7 \pm 0.5) %	39
$D^+\pi^0$	(30.7 \pm 0.5) %	38
$D^+\gamma$	(1.6 \pm 0.4) %	136

 $D_0^*(2300)$

$$I(J^P) = \frac{1}{2}(0^+)$$

was $D_0^*(2400)$

Mass $m = 2343 \pm 10$ MeV ($S = 1.5$)

Full width $\Gamma = 229 \pm 16$ MeV

$D_0^*(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D\pi^{\pm}$	seen	411

$D_1(2420)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2422.1 \pm 0.6$ MeV ($S = 1.7$)

$m_{D_1(2420)^0} - m_{D^{*+}} = 411.8 \pm 0.6$ MeV ($S = 1.7$)

$m_{D_1(2420)^{\pm}} - m_{D_1(2420)^0} = 4 \pm 4$ MeV

Full width $\Gamma = 31.3 \pm 1.9$ MeV ($S = 2.8$)

$\overline{D}_1(2420)$ modes are charge conjugates of modes below.

 $D_1(2420)$ DECAY MODESFraction (Γ_i/Γ) p (MeV/c) $D^*(2007)^0 \pi$

seen

359

 $D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2412 \pm 9$ MeV

Full width $\Gamma = 314 \pm 29$ MeV

 $D_1(2430)^0$ DECAY MODESFraction (Γ_i/Γ) p (MeV/c) $D^*(2010)^+ \pi^-$

seen

345

 $D_2^*(2460)$

$$I(J^P) = \frac{1}{2}(2^+)$$

Mass $m = 2461.1 \pm 0.8$ MeV ($S = 6.3$)

$m_{D_2^*(2460)^0} - m_{D^+} = 591.5 \pm 0.8$ MeV ($S = 6.0$)

$m_{D_2^*(2460)^0} - m_{D^{*+}} = 450.9 \pm 0.8$ MeV ($S = 6.0$)

$m_{D_2^*(2460)^{\pm}} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7$ MeV

Full width $\Gamma = 47.3 \pm 0.8$ MeV ($S = 1.5$)

$\overline{D}_2^*(2460)$ modes are charge conjugates of modes below.

 $D_2^*(2460)$ DECAY MODESFraction (Γ_i/Γ) p (MeV/c) $D \pi^-$

seen

509

 $D^*(2010) \pi^-$

seen

389

 $D_3^*(2750)$

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass $m = 2763.1 \pm 3.2$ MeV ($S = 2.1$)

Full width $\Gamma = 66 \pm 5$ MeV

D₃[*](2750) DECAY MODES	Fraction (Γ_i/Γ)	<i>p</i> (MeV/c)
$D\pi$	seen	743
$D^+\pi^-$	seen	739
$D^0\pi^\pm$	seen	743
$D^*\pi$	seen	639
$D^{*+}\pi^-$	seen	639

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^-2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] See the listings under "D $\rightarrow K\pi\pi\pi$ partial wave analyses" and our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+\ell^+\ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 Review, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^-2\pi^+\pi^-$, $K^-2\pi^+\pi^-\pi^0$, $\bar{K}^02\pi^+2\pi^-$, $K^+2K^-\pi^+$, $2\pi^+2\pi^-$, $2\pi^+2\pi^-\pi^0$, $K^+K^-\pi^+\pi^-$, and $K^+K^-\pi^+\pi^-\pi^0$, branching fractions.
- [l] This is the sum of our $K^-3\pi^+2\pi^-$ and $3\pi^+3\pi^-$ branching fractions.
- [n] The branching fractions for the $K^-e^+\nu_e$, $K^*(892)^-e^+\nu_e$, $\pi^-e^+\nu_e$, and $\rho^-e^+\nu_e$ modes add up to 6.17 ± 0.17 %.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] Submodes of the $D^0 \rightarrow K_S^0\pi^+\pi^-\pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new

for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.

- [q] This branching fraction includes all the decay modes of the resonance in the final state.
- [r] This limit assumes the average of $B(D^0 \rightarrow K^- e^+ \nu_e)$ and $B(D^0 \rightarrow K^- \mu^+ \nu_\mu)$ for the $B(D^0 \rightarrow K^- \ell^+ \nu_\ell)$ value.
- [s] The value is for the sum of the charge states or particle/antiparticle states indicated.