

$\gamma(1S)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\gamma(1S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9460.40±0.09±0.04	¹ SHAMOV 23	RVUE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9460.11±0.11±0.07	² SHAMOV 23	RVUE	$e^+ e^- \rightarrow$ hadrons
9460.51±0.09±0.05	^{3,4} ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
9460.60±0.09±0.05	^{5,6} BARU 92B	MD1	$e^+ e^- \rightarrow$ hadrons
9460.59±0.12	BARU 86	MD1	$e^+ e^- \rightarrow$ hadrons
9460.6 ±0.4	^{6,7} ARTAMONOV 84	MD1	$e^+ e^- \rightarrow$ hadrons
9459.97±0.11±0.07	⁸ MACKAY 84	CUSB	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.			
2 Obtained by reanalysing CUSB data (MACKAY 84), but not authored by the CUSB collaboration.			
3 Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
4 Superseded by SHAMOV 23.			
5 Supersedes BARU 86.			
6 Superseded by ARTAMONOV 00.			
7 Value includes data of ARTAMONOV 82.			
8 Reanalysed by SHAMOV 23.			

 $\gamma(1S)$ WIDTH

VALUE (keV)	DOCUMENT ID
54.02±1.25 OUR EVALUATION	See the Note on "Width Determinations of the γ States"

 $\gamma(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \tau^+ \tau^-$	(2.60 ± 0.10) %	
$\Gamma_2 e^+ e^-$	(2.39 ± 0.08) %	
$\Gamma_3 \mu^+ \mu^-$	(2.48 ± 0.04) %	

Hadronic decays

$\Gamma_4 ggg$	(81.7 ± 0.7) %	
$\Gamma_5 \gamma gg$	(2.2 ± 0.6) %	
$\Gamma_6 \eta'(958)$ anything	(2.94 ± 0.24) %	
$\Gamma_7 J/\psi(1S)$ anything	(5.4 ± 0.4) $\times 10^{-4}$	S=1.4
$\Gamma_8 J/\psi(1S) \eta_c$	< 2.2 $\times 10^{-6}$	CL=90%
$\Gamma_9 J/\psi(1S) \chi_{c0}$	< 3.4 $\times 10^{-6}$	CL=90%
$\Gamma_{10} J/\psi(1S) \chi_{c1}$	(3.9 ± 1.2) $\times 10^{-6}$	
$\Gamma_{11} J/\psi(1S) \chi_{c2}$	< 1.4 $\times 10^{-6}$	CL=90%

Γ_{12}	$J/\psi(1S)\eta_c(2S)$	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{13}	$J/\psi(1S)X(3940)$	< 5.4	$\times 10^{-6}$	CL=90%
Γ_{14}	$J/\psi(1S)X(4160)$	< 5.4	$\times 10^{-6}$	CL=90%
Γ_{15}	$X(4350)$ anything, $X \rightarrow J/\psi(1S)\phi$	< 8.1	$\times 10^{-6}$	CL=90%
Γ_{16}	$T_{c\bar{c}1}(3900)^{\pm}$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^{\pm}$	< 1.3	$\times 10^{-5}$	CL=90%
Γ_{17}	$T_{c\bar{c}1}(4200)^{\pm}$ anything, $Z_c \rightarrow J/\psi(1S)\pi^{\pm}$	< 6.0	$\times 10^{-5}$	CL=90%
Γ_{18}	$T_{c\bar{c}1}(4430)^{\pm}$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^{\pm}$	< 4.9	$\times 10^{-5}$	CL=90%
Γ_{19}	X_{cs}^{\pm} anything, $X \rightarrow J/\psi K^{\pm}$	< 5.7	$\times 10^{-6}$	CL=90%
Γ_{20}	$\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)\pi^{+}\pi^{-}$	< 3.8	$\times 10^{-5}$	CL=90%
Γ_{21}	$\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)K^{+}K^{-}$	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{22}	$\chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow J/\psi(1S)\phi$	< 5.2	$\times 10^{-6}$	CL=90%
Γ_{23}	χ_{c0} anything	< 4	$\times 10^{-3}$	CL=90%
Γ_{24}	χ_{c1} anything	(1.90 \pm 0.35)	$\times 10^{-4}$	
Γ_{25}	$\chi_{c1}(1P)X_{tetra}$	< 3.78	$\times 10^{-5}$	CL=90%
Γ_{26}	χ_{c2} anything	(2.8 \pm 0.8)	$\times 10^{-4}$	
Γ_{27}	$\psi(2S)$ anything	(1.23 \pm 0.20)	$\times 10^{-4}$	
Γ_{28}	$\psi(2S)\eta_c$	< 3.6	$\times 10^{-6}$	CL=90%
Γ_{29}	$\psi(2S)\chi_{c0}$	< 6.5	$\times 10^{-6}$	CL=90%
Γ_{30}	$\psi(2S)\chi_{c1}$	< 4.5	$\times 10^{-6}$	CL=90%
Γ_{31}	$\psi(2S)\chi_{c2}$	< 2.1	$\times 10^{-6}$	CL=90%
Γ_{32}	$\psi(2S)\eta_c(2S)$	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\psi(2S)X(3940)$	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{34}	$\psi(2S)X(4160)$	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{35}	$\psi(4230)$ anything, $\psi \rightarrow \psi(2S)\pi^{+}\pi^{-}$	< 7.9	$\times 10^{-5}$	CL=90%
Γ_{36}	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^{+}\pi^{-}$	< 5.2	$\times 10^{-5}$	CL=90%
Γ_{37}	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^{+}\pi^{-}$	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{38}	$T_{c\bar{c}}(4050)^{\pm}$ anything, $X \rightarrow \psi(2S)\pi^{\pm}$	< 8.8	$\times 10^{-5}$	CL=90%
Γ_{39}	$T_{c\bar{c}1}(4430)^{\pm}$ anything, $T_{c\bar{c}1} \rightarrow \psi(2S)\pi^{\pm}$	< 6.7	$\times 10^{-5}$	CL=90%
Γ_{40}	$\chi_{c1}(3872)$ anything	< 2.7	$\times 10^{-4}$	CL=90%
Γ_{41}	$T_{c\bar{c}1}(4200)^{+}T_{c\bar{c}1}(4200)^{-}$	< 2.23	$\times 10^{-5}$	CL=90%
Γ_{42}	$T_{c\bar{c}1}(3900)^{\pm}T_{c\bar{c}1}(4200)^{\mp}$	< 8.1	$\times 10^{-6}$	CL=90%
Γ_{43}	$T_{c\bar{c}1}(3900)^{+}T_{c\bar{c}1}(3900)^{-}$	< 1.8	$\times 10^{-6}$	CL=90%

Γ_{44}	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	< 1.58	$\times 10^{-5}$	CL=90%
Γ_{45}	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	< 2.66	$\times 10^{-5}$	CL=90%
Γ_{46}	$T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$	< 4.42	$\times 10^{-5}$	CL=90%
Γ_{47}	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%
Γ_{48}	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$	< 2.33	$\times 10^{-5}$	CL=90%
Γ_{49}	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$	< 4.55	$\times 10^{-5}$	CL=90%
Γ_{50}	$\rho\pi$	< 3.68	$\times 10^{-6}$	CL=90%
Γ_{51}	$\omega\pi^0$	< 3.90	$\times 10^{-6}$	CL=90%
Γ_{52}	$\pi^+\pi^-$	< 5	$\times 10^{-4}$	CL=90%
Γ_{53}	K^+K^-	< 5	$\times 10^{-4}$	CL=90%
Γ_{54}	$p\bar{p}$	< 5	$\times 10^{-4}$	CL=90%
Γ_{55}	$\pi^+\pi^-\pi^0$	(2.1 \pm 0.8) $\times 10^{-6}$		
Γ_{56}	ϕK^+K^-	(2.4 \pm 0.5) $\times 10^{-6}$		
Γ_{57}	$\omega\pi^+\pi^-$	(4.5 \pm 1.0) $\times 10^{-6}$		
Γ_{58}	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(4.4 \pm 0.8) $\times 10^{-6}$		
Γ_{59}	$\phi f_2'(1525)$	< 1.63	$\times 10^{-6}$	CL=90%
Γ_{60}	$\omega f_2(1270)$	< 1.79	$\times 10^{-6}$	CL=90%
Γ_{61}	$\rho(770) a_2(1320)$	< 2.24	$\times 10^{-6}$	CL=90%
Γ_{62}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	(3.0 \pm 0.8) $\times 10^{-6}$		
Γ_{63}	$K_1(1270)^\pm K^\mp$	< 2.41	$\times 10^{-6}$	CL=90%
Γ_{64}	$K_1(1400)^\pm K^\mp$	(1.0 \pm 0.4) $\times 10^{-6}$		
Γ_{65}	$b_1(1235)^\pm \pi^\mp$	< 1.25	$\times 10^{-6}$	CL=90%
Γ_{66}	$\pi^+\pi^-\pi^0\pi^0$	(1.28 \pm 0.30) $\times 10^{-5}$		
Γ_{67}	$K_S^0 K^+ \pi^- + \text{c.c.}$	(1.6 \pm 0.4) $\times 10^{-6}$		
Γ_{68}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	(2.9 \pm 0.9) $\times 10^{-6}$		
Γ_{69}	$K^*(892)^- K^+ + \text{c.c.}$	< 1.11	$\times 10^{-6}$	CL=90%
Γ_{70}	$f_1(1285)$ anything	(4.6 \pm 3.1) $\times 10^{-3}$		
Γ_{71}	$D^*(2010)^\pm$ anything	(2.52 \pm 0.20) %		
Γ_{72}	$f_1(1285) X_{\text{tetra}}$	< 6.24	$\times 10^{-5}$	CL=90%
Γ_{73}	2H anything	(2.85 \pm 0.25) $\times 10^{-5}$		
Γ_{74}	Sum of 100 exclusive modes	(1.200 \pm 0.017) %		

Radiative decays

Γ_{75}	$\gamma\pi^+\pi^-$	(6.3 \pm 1.8) $\times 10^{-5}$		
Γ_{76}	$\gamma\pi^0\pi^0$	(1.7 \pm 0.7) $\times 10^{-5}$		
Γ_{77}	$\gamma\pi\pi$ (S-wave)	(4.6 \pm 0.7) $\times 10^{-5}$		
Γ_{78}	$\gamma\pi^0\eta$	< 2.4	$\times 10^{-6}$	CL=90%
Γ_{79}	$\gamma K^+ K^-$	[a] (1.14 \pm 0.13) $\times 10^{-5}$		
Γ_{80}	$\gamma p\bar{p}$	[b] < 6	$\times 10^{-6}$	CL=90%
Γ_{81}	$\gamma 2h^+ 2h^-$	(7.0 \pm 1.5) $\times 10^{-4}$		
Γ_{82}	$\gamma 3h^+ 3h^-$	(5.4 \pm 2.0) $\times 10^{-4}$		
Γ_{83}	$\gamma 4h^+ 4h^-$	(7.4 \pm 3.5) $\times 10^{-4}$		
Γ_{84}	$\gamma\pi^+\pi^- K^+ K^-$	(2.9 \pm 0.9) $\times 10^{-4}$		
Γ_{85}	$\gamma 2\pi^+ 2\pi^-$	(2.5 \pm 0.9) $\times 10^{-4}$		

Γ_{86}	$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
Γ_{87}	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
Γ_{88}	$\gamma \pi^+ \pi^- p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
Γ_{89}	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	
Γ_{90}	$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
Γ_{91}	$\gamma \eta'(958)$	$< 1.9 \times 10^{-6}$	CL=90%
Γ_{92}	$\gamma \eta$	$< 1.0 \times 10^{-6}$	CL=90%
Γ_{93}	$\gamma f_0(980)$	$< 3 \times 10^{-5}$	CL=90%
Γ_{94}	$\gamma f'_2(1525)$	$(2.9 \pm 0.6) \times 10^{-5}$	
Γ_{95}	$\gamma f_2(1270)$	$(1.01 \pm 0.06) \times 10^{-4}$	
Γ_{96}	$\gamma \eta(1405)$	$< 8.2 \times 10^{-5}$	CL=90%
Γ_{97}	$\gamma f_0(1500)$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{98}	$\gamma f_0(1500) \rightarrow \gamma K^+ K^-$	$(1.0 \pm 0.4) \times 10^{-5}$	
Γ_{99}	$\gamma f_0(1710)$	$< 2.6 \times 10^{-4}$	CL=90%
Γ_{100}	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	$(1.01 \pm 0.32) \times 10^{-5}$	
Γ_{101}	$\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-$	$(5.3 \pm 2.0) \times 10^{-6}$	
Γ_{102}	$\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0$	$< 1.4 \times 10^{-6}$	CL=90%
Γ_{103}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$< 1.8 \times 10^{-6}$	CL=90%
Γ_{104}	$\gamma f_4(2050)$	$< 5.3 \times 10^{-5}$	CL=90%
Γ_{105}	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2 \times 10^{-4}$	CL=90%
Γ_{106}	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 8 \times 10^{-7}$	CL=90%
Γ_{107}	$\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-$	$< 6 \times 10^{-7}$	CL=90%
Γ_{108}	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$< 1.1 \times 10^{-6}$	CL=90%
Γ_{109}	$\gamma \eta(2225) \rightarrow \gamma \phi \phi$	$< 3 \times 10^{-3}$	CL=90%
Γ_{110}	$\gamma \eta_c(1S)$	$< 2.9 \times 10^{-5}$	CL=90%
Γ_{111}	$\gamma \eta_c(2S)$	$< 4 \times 10^{-4}$	CL=90%
Γ_{112}	$\gamma \chi_{c0}$	$< 6.6 \times 10^{-5}$	CL=90%
Γ_{113}	$\gamma \chi_{c1}$	$(4.7 \pm 2.4) \times 10^{-5}$	
Γ_{114}	$\gamma \chi_{c2}$	$< 7.6 \times 10^{-6}$	CL=90%
Γ_{115}	$\gamma \chi_{c1}(3872)$	$< 5 \times 10^{-5}$	CL=90%
Γ_{116}	$\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	$< 2.8 \times 10^{-6}$	CL=90%
Γ_{117}	$\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi$	$< 3.0 \times 10^{-6}$	CL=90%
Γ_{118}	$\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi$	$< 2.2 \times 10^{-6}$	CL=90%
Γ_{119}	$\gamma X\bar{X} (m_X < 3.1 \text{ GeV})$	$[c] < 1 \times 10^{-3}$	CL=90%
Γ_{120}	$\gamma X\bar{X} (m_X < 4.5 \text{ GeV})$	$[d] < 2.4 \times 10^{-4}$	CL=90%
Γ_{121}	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	$[e] < 1.78 \times 10^{-4}$	CL=95%
Γ_{122}	γA^0	$[f]$	
Γ_{123}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[g] < 9 \times 10^{-6}$	CL=90%
Γ_{124}	$\gamma A^0 \rightarrow \gamma \tau^+ \tau^-$	$[a] < 1.30 \times 10^{-4}$	CL=90%
Γ_{125}	$\gamma A^0 \rightarrow \gamma g g$	$[h] < 1 \%$	CL=90%
Γ_{126}	$\gamma A^0 \rightarrow \gamma s\bar{s}$	$[h] < 1 \times 10^{-3}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{127}	$e^\pm \mu^\mp$	<i>LF</i>	< 3.9	$\times 10^{-7}$	CL=90%
Γ_{128}	$\mu^\pm \tau^\mp$	<i>LF</i>	< 2.7	$\times 10^{-6}$	CL=90%
Γ_{129}	$e^\pm \tau^\mp$	<i>LF</i>	< 2.7	$\times 10^{-6}$	CL=90%
Γ_{130}	$\gamma e^\pm \mu^\mp$	<i>LF</i>	< 4.2	$\times 10^{-7}$	CL=90%
Γ_{131}	$\gamma \mu^\pm \tau^\mp$	<i>LF</i>	< 6.1	$\times 10^{-6}$	CL=90%
Γ_{132}	$\gamma e^\pm \tau^\mp$	<i>LF</i>	< 6.5	$\times 10^{-6}$	CL=90%

Other decays

Γ_{133}	invisible	< 3.0	$\times 10^{-4}$	CL=90%
Γ_{134}	hadrons	(96 \pm 4)	%	

[a] $2m_\tau < M(\tau^+ \tau^-) < 9.2$ GeV

[b] 2 GeV $< m_{K^+ K^-} < 3$ GeV

[c] $X \bar{X}$ = vectors with $m < 3.1$ GeV

[d] X and \bar{X} = zero spin with $m < 4.5$ GeV

[e] 1.5 GeV $< m_X < 5.0$ GeV

[f] A^0 = scalar with $m < 8.0$ GeV

[g] 201 MeV $< M(\mu^+ \mu^-) < 3565$ MeV

[h] 0.5 GeV $< m_X < 9.0$ GeV, where m_X is the invariant mass of the hadronic final state.

 $\Gamma(1S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_3 \Gamma_2 / \Gamma$
<u>VALUE (eV)</u> $31.2 \pm 1.6 \pm 1.7$	<u>DOCUMENT ID</u> KOBEL 92 <u>TECN</u> <u>COMMENT</u> $e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_{134} \Gamma_2 / \Gamma$
<u>VALUE (keV)</u> 1.240 ± 0.016 OUR AVERAGE	<u>DOCUMENT ID</u> 1 ROSNER 06 CLEO 9.5 $e^+ e^- \rightarrow$ hadrons 1 BARU 92B MD1 $e^+ e^- \rightarrow$ hadrons 1 JAKUBOWSKI 88 CBAL $e^+ e^- \rightarrow$ hadrons 2 GILES 84B CLEO $e^+ e^- \rightarrow$ hadrons 2 ALBRECHT 82 DASP $e^+ e^- \rightarrow$ hadrons 2 NICZYPORUK 82 LENA $e^+ e^- \rightarrow$ hadrons 2 BOCK 80 CNTR $e^+ e^- \rightarrow$ hadrons 3 BERGER 79 PLUT $e^+ e^- \rightarrow$ hadrons

¹ Radiative corrections evaluated following KURAEV 85.

² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

³ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS $\Gamma(e^+ e^-)$

VALUE (keV)

 1.340 ± 0.018 OUR EVALUATION Γ_2

DOCUMENT ID

 $\Upsilon(1S)$ BRANCHING RATIOS $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_1/Γ VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT

 2.60 ± 0.10 OUR AVERAGE

2.53 $\pm 0.13 \pm 0.04$	60k	¹ BESSON	07	CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+ \tau^-$
2.61 $\pm 0.12^{+0.09}_{-0.13}$	25k	CINABRO	94B	CLE2	$e^+ e^- \rightarrow \tau^+ \tau^-$
2.7 ± 0.4 ± 0.2		² ALBRECHT	85C	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \tau^+ \tau^-$
3.4 ± 0.4 ± 0.4		GILES	83	CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(1S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+ \mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_2/Γ VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT

 2.39 ± 0.08 OUR AVERAGE

2.40 $\pm 0.01 \pm 0.12$	191k	PATRA	22	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
2.29 $\pm 0.08 \pm 0.11$		ALEXANDER	98	CLE2	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
2.42 $\pm 0.14 \pm 0.14$	307	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
2.8 ± 0.3 ± 0.2	826	BESSON	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
5.1 ± 3.0		BERGER	80C	PLUT	$e^+ e^- \rightarrow e^+ e^-$

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_3/Γ VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT

 2.48 ± 0.04 OUR AVERAGE

2.46 $\pm 0.01 \pm 0.11$	246k	PATRA	22	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
2.49 $\pm 0.02 \pm 0.07$	345k	ADAMS	05	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.49 $\pm 0.08 \pm 0.13$		ALEXANDER	98	CLE2	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
2.12 $\pm 0.20 \pm 0.10$		¹ BARU	92	MD1	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.31 $\pm 0.12 \pm 0.10$		¹ KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.52 $\pm 0.07 \pm 0.07$		CHEN	89B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.61 $\pm 0.09 \pm 0.11$		KAARSBERG	89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.30 $\pm 0.25 \pm 0.13$	86	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
2.9 ± 0.3 ± 0.2	864	BESSON	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
2.7 ± 0.3 ± 0.3		ANDREWS	83	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
3.2 ± 1.3 ± 0.3		ALBRECHT	82	DASP	$e^+ e^- \rightarrow \mu^+ \mu^-$

3.8 ± 1.5 ± 0.2	NICZYPORUK 82	LENA	$e^+ e^- \rightarrow \mu^+ \mu^-$
1.4 ± 3.4 -1.4	BOCK 80	CNTR	$e^+ e^- \rightarrow \mu^+ \mu^-$
2.2 ± 2.0	BERGER 79	PLUT	$e^+ e^- \rightarrow \mu^+ \mu^-$

¹ Taking into account interference between the resonance and continuum.

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_3
1.008 ± 0.023 OUR AVERAGE					
1.005 ± 0.013 ± 0.022	0.7M	¹ DEL-AMO-SA..10c	BABR	$\gamma(3S) \rightarrow \pi^+ \pi^- \gamma(1S)$	
1.02 ± 0.02 ± 0.05	60k	BESSON 07	CLEO	$e^+ e^- \rightarrow \gamma(1S)$	

¹ Allows any number of extra photons with total energy < 500 MeV.

$\Gamma(ggg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
81.7 ± 0.7	20M	¹ BESSON	06A	CLEO	$\gamma(1S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSON 06A and PDG 08 values of $B(\mu^+ \mu^-) = (2.48 \pm 0.05)\%$ and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
2.20 ± 0.60	400k	¹ BESSON	06A	CLEO	$\gamma(1S) \rightarrow \gamma + \text{hadrons}$

¹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma gg)/\Gamma(ggg)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_4
2.70 $\pm 0.01 \pm 0.27$	20M	BESSON	06A	CLEO	$\gamma(1S) \rightarrow (\gamma +) \text{hadrons}$

$\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
0.0294 ± 0.0024 OUR AVERAGE				
0.030 ± 0.002 ± 0.002	AQUINES 06A	CLE3	$\gamma(1S) \rightarrow \eta' \text{ anything}$	
0.028 ± 0.004 ± 0.002	ARTUSO 03	CLE2	$\gamma(1S) \rightarrow \eta' \text{ anything}$	

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
5.4 ± 0.4 OUR FIT					Error includes scale factor of 1.4.	
5.4 ± 0.4 OUR AVERAGE					Error includes scale factor of 1.5.	
5.25 ± 0.13 ± 0.25		3k	SHEN 16	BELL	$e^+ e^- \rightarrow J/\psi X$	
6.4 ± 0.4 ± 0.6		730	BRIERE 04	CLEO	$e^+ e^- \rightarrow J/\psi X$	
11 ± 4 ± 2			¹ FULTON 89	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<6.8		90	ALBRECHT 92J	ARG	$e^+ e^- \rightarrow e^+ e^- X, \mu^+ \mu^- X$	
<17		90	MASCHMANN 90	CBAL	$e^+ e^- \rightarrow \text{hadrons}$	
<200		90	NICZYPORUK 83	LENA		

¹ Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (6.9 \pm 0.9)\%$.

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$

VALUE	CL%
$<2.2 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_8/Γ

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$

VALUE	CL%
$<3.4 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_9/Γ

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS
$3.90 \pm 1.21 \pm 0.23$	20

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_{10}/Γ

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$

VALUE	CL%
$<1.4 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_{11}/Γ

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$

VALUE	CL%
$<2.2 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_{12}/Γ

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$

VALUE	CL%
$<5.4 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_{13}/Γ

$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$

VALUE	CL%
$<5.4 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
YANG	BELL	$e^+ e^- \rightarrow J/\psi X$

Γ_{14}/Γ

$\Gamma(X(4350) \text{ anything}, X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$

VALUE	CL%
$<8.1 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
SHEN	BELL	$\gamma(1S) \rightarrow J/\psi K^+ K^- X$

Γ_{15}/Γ

$\Gamma(T_{c\bar{c}1}(3900)^{\pm} \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$

VALUE	CL%
$<1.3 \times 10^{-5}$	90

DOCUMENT ID	TECN	COMMENT
SHEN	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$

Γ_{16}/Γ

$\Gamma(T_{c\bar{c}1}(4200)^{\pm} \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$

VALUE	CL%
$<6.0 \times 10^{-5}$	90

DOCUMENT ID	TECN	COMMENT
SHEN	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$

Γ_{17}/Γ

$\Gamma(T_{c\bar{c}1}(4430)^{\pm} \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$

VALUE	CL%
$<4.9 \times 10^{-5}$	90

DOCUMENT ID	TECN	COMMENT
SHEN	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$

Γ_{18}/Γ

$\Gamma(X_{cs}^{\pm} \text{ anything}, X \rightarrow J/\psi K^{\pm})/\Gamma_{\text{total}}$

VALUE	CL%
$<5.7 \times 10^{-6}$	90

DOCUMENT ID	TECN	COMMENT
SHEN	BELL	$\gamma(1S) \rightarrow J/\psi K^- X$

Γ_{19}/Γ

$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.8 \times 10^{-5}$	90	SHEN	16	BELL $\gamma(1S) \rightarrow J/\psi\pi^+\pi^- X$

 $\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.5 \times 10^{-6}$	90	SHEN	16	BELL $\gamma(1S) \rightarrow J/\psi K^+K^- X$

 $\Gamma(\chi_{c1}(4140) \text{ anything}, \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL $\gamma(1S) \rightarrow J/\psi K^+K^- X$

 $\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{23}/Γ_7

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7.4	90	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.90 ± 0.35 OUR FIT				
$1.90 \pm 0.43 \pm 0.14$	215	JIA	17	BELL $\gamma(1S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{24}/Γ_7

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.35 ± 0.07 OUR FIT				
$0.35 \pm 0.08 \pm 0.06$	52 ± 12	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$

 $\Gamma(\chi_{c1}(1P)X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<37.8 \times 10^{-6}$	90	1 JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 37.8×10^{-6} .

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{26}/Γ_7

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.52 \pm 0.12 \pm 0.09$	47 ± 11	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.23 \pm 0.17 \pm 0.11$	215	SHEN	16	BELL $e^+e^- \rightarrow \psi(2S) X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{27}/Γ_7

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.41 \pm 0.11 \pm 0.08$	42 ± 11	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi\pi^+\pi^- X$

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.6 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S) X$

$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$	Γ_{29}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.5 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$	Γ_{30}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.5 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$	Γ_{31}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.1 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$	Γ_{32}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.2 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$	Γ_{33}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.9 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$	Γ_{34}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.9 \times 10^{-6}$	90	YANG	14	$e^+ e^- \rightarrow \psi(2S)X$
$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{35}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.9 \times 10^{-5}$	90	SHEN	16	$\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$
$\Gamma(\psi(4360) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{36}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.2 \times 10^{-5}$	90	SHEN	16	$\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$
$\Gamma(\psi(4660) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{37}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.2 \times 10^{-5}$	90	SHEN	16	$\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$
$\Gamma(T_{c\bar{c}}(4050)^{\pm} \text{ anything}, X \rightarrow \psi(2S)\pi^{\pm})/\Gamma_{\text{total}}$	Γ_{38}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.8 \times 10^{-5}$	90	SHEN	16	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$
$\Gamma(T_{c\bar{c}1}(4430)^{\pm} \text{ anything}, T_{c\bar{c}1} \rightarrow \psi(2S)\pi^{\pm})/\Gamma_{\text{total}}$	Γ_{39}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.7 \times 10^{-5}$	90	SHEN	16	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$
$\Gamma(\chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}$	Γ_{40}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.7 \times 10^{-4}$	90	1 SHEN	16	$\gamma(1S) \rightarrow J/\psi\pi^+\pi^- X$

¹ SHEN 16 reports $[\Gamma(\gamma(1S) \rightarrow \chi_{c1}(3872)\text{anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 9.5 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.

$\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<22.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$.

$\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.1 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$.

$\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<15.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.6 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<44.2 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

$\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<23.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<45.5 \times 10^{-6}$	90	¹ JIA	18	BELL $\gamma(1S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_{50}/Γ
<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.68	90	SHEN	13	BELL	$\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<1 \times 10^3$	90	BLINOV	90	MD1	$\gamma(1S) \rightarrow \rho^0 \pi^0$
$<2 \times 10^2$	90	FULTON	90B		$\gamma(1S) \rightarrow \rho^0 \pi^0$
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA	$\gamma(1S) \rightarrow \rho^0 \pi^0$
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{51}/Γ
<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.90	90	SHEN	13	BELL	$\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{52}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<5	90	BARU	92	MD1	$\gamma(1S) \rightarrow \pi^+ \pi^-$
$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$					Γ_{53}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<5	90	BARU	92	MD1	$\gamma(1S) \rightarrow K^+ K^-$
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$					Γ_{54}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<5	90	¹ BARU	96	MD1	$\gamma(1S) \rightarrow p\bar{p}$
$\bullet \bullet \bullet$ Supersedes BARU 92 in this node.					
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{55}/Γ
<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.14±0.72±0.34		26 ± 9	SHEN	13	$\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<18.4	90		ANASTASSOV	99	$e^+ e^- \rightarrow \text{hadrons}$
$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$					Γ_{56}/Γ
<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.36±0.37±0.29	56	SHEN	12A	BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$
$\Gamma(\omega\pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{57}/Γ
<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.46±0.67±0.72	64	SHEN	12A	BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{58}/Γ
<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.42±0.50±0.58	173	SHEN	12A	BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$					Γ_{59}/Γ
<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.63	90	SHEN	12A	BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$				Γ_{60}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.79	90	SHEN	12A	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$				Γ_{61}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.24	90	SHEN	12A	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{62}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.02 ± 0.68 ± 0.34	42	SHEN	12A	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$				Γ_{63}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.41	90	SHEN	12A	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$				Γ_{64}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.02 ± 0.35 ± 0.22	24	SHEN	12A	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$				Γ_{65}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.25	90	SHEN	12A	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$				Γ_{66}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.8 ± 2.0 ± 2.3	143 ± 22	SHEN	13	$\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{67}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
1.59 ± 0.33 ± 0.18		37 ± 8	SHEN	13
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.4	90	¹ DOBBS	12A	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{68}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.92 ± 0.85 ± 0.37	16 ± 5	SHEN	13	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$
$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{69}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.11	90	SHEN	13	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$
$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$				Γ_{70}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.6 ± 2.8 ± 1.3	3.1k	JIA	17A	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(D^*(2010)^{\pm} \text{ anything})/\Gamma_{\text{total}}$ Γ_{71}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$25.2 \pm 1.3 \pm 1.5$	$\approx 2k$	1	AUBERT	10C	BABR $\gamma(2S) \rightarrow \pi^+ \pi^- \gamma(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19 90 2 ALBRECHT 92J ARG $e^+ e^- \rightarrow D^0 \pi^\pm X$

¹ For $x_p > 0.1$.

² For $x_p > 0.2$.

 $\Gamma(f_1(1285)X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<62.4 \times 10^{-6}$	90	1 JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.6×10^{-6} to 62.4×10^{-6} .

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.85 ± 0.25 OUR AVERAGE				
$2.81 \pm 0.49^{+0.20}_{-0.24}$		LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$
$2.86 \pm 0.19 \pm 0.21$	455	ASNER	07	CLEO $e^+ e^- \rightarrow \overline{2H} X$

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>COMMENT</u>
1.200 ± 0.017	1,2 DOBBS	12A $\gamma(1S) \rightarrow \text{hadrons}$

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(ggg, \gamma gg \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.36 \pm 0.23 \pm 0.25$	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$

 $\Gamma(\gamma\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.3 \pm 1.2 \pm 1.3$	1 ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

¹ For $m_{\pi\pi} > 1$ GeV.

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.7 \pm 0.6 \pm 0.3$	1 ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

¹ For $m_{\pi\pi} > 1$ GeV.

 $\Gamma(\gamma\pi\pi(\text{S-wave}))/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.63 \pm 0.56 \pm 0.48$	LEES	18A	BABR $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$

$\Gamma(\gamma\pi^0\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{78}/Γ
<2.4	90	¹ BESSON	07A	CLEO $e^+ e^- \rightarrow \gamma(1S)$	

¹ BESSON 07A obtained this limit for $0.7 < m_{\pi^0\eta} < 3$ GeV.

 $\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}}$

($2 < m_{K^+ K^-} < 3$ GeV)

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{79}/Γ
1.14±0.08±0.10	90	ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma K^+ K^-$	

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$

($2 < m_{p\bar{p}} < 3$ GeV)

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{80}/Γ
<0.6	90	ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma p\bar{p}$	

 $\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
7.0±1.1±1.0	80 ± 12

 $\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
5.4±1.5±1.3	39 ± 11

 $\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
7.4±2.5±2.5	36 ± 12

 $\Gamma(\gamma\pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
2.9±0.7±0.6	29 ± 8

 $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
2.5±0.7±0.5	26 ± 7

 $\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
2.5±0.9±0.8	17 ± 5

 $\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
2.4±0.9±0.8	18 ± 7

 $\Gamma(\gamma\pi^+\pi^- p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
1.5±0.5±0.3	22 ± 6

 Γ_{78}/Γ Γ_{79}/Γ Γ_{80}/Γ Γ_{81}/Γ Γ_{82}/Γ Γ_{83}/Γ Γ_{84}/Γ Γ_{85}/Γ Γ_{86}/Γ Γ_{87}/Γ Γ_{88}/Γ

$\Gamma(\gamma 2\pi^+ 2\pi^- p\bar{p})/\Gamma_{\text{total}}$ Γ_{89}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.4 \pm 0.4 \pm 0.4$	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.2 ± 0.2	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{91}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.9	90	ATHAR	07A CLEO	$\gamma(1S) \rightarrow \gamma\eta' \rightarrow \gamma\pi^+\pi^-\eta, \gamma\rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	RICHICHI	01B CLE2	$\gamma(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{92}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.0	90	ATHAR	07A CLEO	$\gamma(1S) \rightarrow \gamma\eta \rightarrow \gamma\gamma\gamma, \gamma\pi^+\pi^-\pi^0, \gamma 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<21	90	MASEK	02 CLEO	$\gamma(1S) \rightarrow \gamma\eta$

 $\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$ Γ_{93}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	¹ ATHAR	06 CLE3	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$.

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{94}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.9 ± 0.6 OUR AVERAGE					
2.13 ± 0.28 ± 0.72			¹ LEES	18A BABR	$\gamma(1S) \rightarrow \gamma K^+ K^-$
4.0 ± 1.4 ± 0.1		17	² BESSON	11 CLEO	$\gamma(1S) \rightarrow K_S^0 K_S^0$
3.7 ± 0.9 ± 0.8			ATHAR	06 CLE3	$\gamma(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

<14 90 ³ FULTON 90B CLEO $\gamma(1S) \rightarrow \gamma K^+ K^-$

<19.4 90 ³ ALBRECHT 89 ARG $\gamma(1S) \rightarrow \gamma K^+ K^-$

¹ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$ and $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$.

² BESSON 11 reports $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\gamma(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})]$ assuming $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$, which we rescale to our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ and $B(f'_2(1525) \rightarrow K\bar{K}) = 4 B(f'_2(1525) \rightarrow K_S^0 K_S^0)$.

³ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$	Γ_{95}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.1 ± 0.6 OUR AVERAGE				
$10.15 \pm 0.59^{+0.54}_{-0.43}$		¹ LEES	18A BABR	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
$10.5 \pm 1.6^{+1.9}_{-1.8}$		² BESSON	07A CLE3	$\gamma(1S) \rightarrow \gamma\pi^0\pi^0$
$10.2 \pm 0.8 \pm 0.7$		ATHAR	06 CLE3	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
$8.1 \pm 2.3^{+2.9}_{-2.7}$		³ ANASTASSOV	99 CLE2	$e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	³ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
<13	90	³ ALBRECHT	89 ARG	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
<81	90	SCHMITT	88 CBAL	$\gamma(1S) \rightarrow \gamma X$
¹ Using $B(f_2(1270) \rightarrow \pi^0\pi^0) = 1/3 B(f_2(1270) \rightarrow \pi\pi)$ and $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9})\%$.				
² Using $B(f_2(1270) \rightarrow \pi^0\pi^0) = B(f_2(1270) \rightarrow \pi\pi)/3$ and $B(f_2(1270) \rightarrow \pi\pi) = (84.7^{+2.5}_{-1.2})\%$.				
³ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.				

$\Gamma(\gamma\eta(1405))/\Gamma_{\text{total}}$	Γ_{96}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.2	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K^\pm\pi^\mp K_S^0$

¹ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm\pi^\mp K_S^0$.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$	Γ_{97}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5	90	¹ BESSON	07A CLEO	$e^+e^- \rightarrow \gamma(1S) \rightarrow \gamma\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.1	90	² BESSON	07A CLEO	$e^+e^- \rightarrow \gamma(1S) \rightarrow \gamma\eta\eta$
¹ Using $B(f_0(1500) \rightarrow \pi^0\pi^0) = B(f_0(1500) \rightarrow \pi\pi)/3$ and $B(f_0(1500) \rightarrow \pi\pi) = (0.349 \pm 0.023)\%$.				
² Calculated by us using $B(f_0(1500) \rightarrow \eta\eta) = (5.1 \pm 0.9)\%$.				

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+K^-)/\Gamma_{\text{total}}$	Γ_{98}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.04 $\pm 0.14 \pm 0.33$	¹ LEES	18A BABR	$e^+e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+K^-$

¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K\bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$.

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$	Γ_{99}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.6	90	¹ ALBRECHT	89 ARG	$\gamma(1S) \rightarrow \gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.3	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K^+K^-$
< 19	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K_S^0\bar{K}_S^0$
< 8	90	² ALBRECHT	89 ARG	$\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
< 24	90	³ SCHMITT	88 CBAL	$\gamma(1S) \rightarrow \gamma X$

¹ Assuming $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$.

² Assuming $B(f_0(1710) \rightarrow \pi\pi) = 0.04$.

³ Assuming $B(f_0(1710) \rightarrow \eta\eta) = 0.18$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.01 ± 0.26 ± 0.18	1	LEES	18A	BABR $e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	ATHAR	06	CLEO $e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+ K^-$
¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K\bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$ assuming $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$.				

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.53 ± 0.17 ± 0.11	1 LEES	18A	BABR $\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$
¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi\pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$ assuming $B(\pi^0\pi^0) = 1/3 B(\pi\pi)$.			

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \pi^0 \pi^0$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta\eta)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \eta\eta$

$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.3	90	1 ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$

¹ Assuming $B(f_4(2050) \rightarrow \pi\pi) = 0.17$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0002	90	BARU	89	MD1 $\gamma(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 8	90	ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 160	90	MASEK	02	CLEO $\gamma(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B	CLEO $\gamma(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89	ARG $\gamma(1S) \rightarrow \gamma K^+ K^-$
<2000	90	BARU	89	MD1 $\gamma(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<120 90 MASEK 02 CLEO $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR	06	$\gamma(1S) \rightarrow \gamma p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<160 90 MASEK 02 CLEO $\gamma(1S) \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma\eta(2225) \rightarrow \gamma\phi\phi)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89	$\gamma(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9×10^{-5}	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.7×10^{-5} 90 SHEN 10A BELL $\gamma(1S) \rightarrow \gamma X$

¹ Using $\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-$ decays.

$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 4×10^{-4}	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$

¹ Using $\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-$ decays.

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.6×10^{-5}	90	¹ KATRENKO	20	BELL $\gamma(1S) \rightarrow \gamma + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.5×10^{-4} 90 SHEN 10A BELL $\gamma(1S) \rightarrow \gamma X$

¹ Using $\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-$ decays.

$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ Γ_{113}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7^{+2.4+0.4}_{-1.8-0.5}$	5	¹ KATRENKO	20	BELL	$\gamma(1S) \rightarrow \gamma + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3 90 SHEN 10A BELL $\gamma(1S) \rightarrow \gamma X$

¹ Using $\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-$ decays.

$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7.6×10^{-6}	90	SHEN	10A	BELL $\gamma(1S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.3×10^{-5} 90 ¹KATRENKO 20 BELL $\gamma(1S) \rightarrow \gamma + \text{hadrons}$

¹ Using $\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-$ decays.

$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$					Γ_{115}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5 \times 10^{-5}$	90	¹ SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$
¹ SHEN 10A reports $[\Gamma(\gamma(1S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.					

$\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$					Γ_{116}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.8 \times 10^{-6}$	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$

$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$					Γ_{117}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.0	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$

$\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{118}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.2	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$

$\Gamma(\gamma X \bar{X} (m_X < 3.1 \text{ GeV})) / \Gamma_{\text{total}}$					Γ_{119}/Γ
$(X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV})$					
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1	90	¹ BAlest	95	CLEO	$e^+ e^- \rightarrow \gamma + X \bar{X}$

¹ For a noninteracting vector X with mass < 3.1 GeV.

$\Gamma(\gamma X \bar{X} (m_X < 4.5 \text{ GeV})) / \Gamma_{\text{total}}$					Γ_{120}/Γ
$(X \bar{X} = \text{zero spin with } m < 4.5 \text{ GeV})$					
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<24	90	¹ DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X \bar{X}$	

¹ For a noninteracting scalar X with mass $m < 4.5$ GeV.

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}) / \Gamma_{\text{total}}$					Γ_{121}/Γ
$(1.5 \text{ GeV} < m_X < 5.0 \text{ GeV})$					
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.78	95	ROSNER	07A	CLEO	$e^+ e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0) / \Gamma_{\text{total}}$					Γ_{122}/Γ
$(A^0 = \text{scalar with } m < 8.0 \text{ GeV})$					
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4.5 \times 10^{-6}$	90	¹ DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X$	
$<3 \times 10^{-5}$	90	² BAlest	95	CLEO	$e^+ e^- \rightarrow \gamma + X$
$<5.6 \times 10^{-5}$	90	² ANTREASYAN	90c	CBAL	$e^+ e^- \rightarrow \gamma + X$

¹ For a non-interacting scalar or pseudoscalar, A^0 , with mass $m_{A^0} < 8.0$ GeV. 90% CL upper limits range from 1.9×10^{-6} to 4.5×10^{-6} .

² For any non-interacting long-lived particle with mass < 7.2 GeV.

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ
(201 < $M(\mu^+ \mu^-)$ < 3565 MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 9	90	1 LOVE	08 CLEO	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 16	90	2 JIA	22 BELL	$\gamma(2S) \rightarrow \gamma \mu^+ \mu^- \pi^+ \pi^-$
< 9.7	90	3 LEES	13C BABR	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with $201 < M(\mu^+ \mu^-) < 3565$ MeV, excluding J/ψ . Measured 90% CL limits as a function of $M(\mu^+ \mu^-)$ range from $1-9 \times 10^{-6}$.

² For a narrow scalar or pseudoscalar, A^0 , with $0.22 < M(A^0) < 9.2$ GeV, resulting in 90% CL upper limits ranging from 3.1×10^{-7} at $M(A^0) = 0.22$ GeV to 1.6×10^{-5} at $M(A^0) = 9.2$ GeV.

³ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 0.212–9.2 GeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} are in the range $0.28-9.7 \times 10^{-6}$.

 $\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{124}/Γ
(2 m_τ < $M(\tau^+ \tau^-)$ < 9.2 GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 130	90	1 LEES	13R BABR	$\gamma(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 150	90	2 JIA	22 BELL	$\gamma(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
< 50	90	3 LOVE	08 CLEO	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 9.2$ GeV, resulting in 90% CL upper limits of 0.9×10^{-5} at $M(A^0) = 2m_\tau$, $\approx 1.5 \times 10^{-5}$ at $M(A^0) = 7.5$ GeV, and 13×10^{-5} at $M(A^0) = 9.2$ GeV.

² For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 9.2$ GeV, resulting in 90% CL upper limits ranging from 3.8×10^{-6} at $M(A^0) = 2m_\tau$ to 1.5×10^{-4} at $M(A^0) = 9.2$ GeV.

³ For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 7.5$ GeV, resulting in 90% CL limits ranging from 1×10^{-5} at $M(A^0) = 2m_\tau$ to 5×10^{-5} at $M(A^0) = 7.5$ GeV.

 $\Gamma(\gamma A^0 \rightarrow \gamma gg)/\Gamma_{\text{total}}$ Γ_{125}/Γ
(0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1 \times 10^{-2}$	90	1 LEES	13L BABR	$\gamma(1S) \rightarrow \gamma X$

¹ For a narrow, CP -odd pseudoscalar, A^0 , searched for in 26 hadronic decay modes with invariant mass $0.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$. Measured 90% CL limits as a function of m_{A^0} range from 10^{-6} to 10^{-2} .

 $\Gamma(\gamma A^0 \rightarrow \gamma s\bar{s})/\Gamma_{\text{total}}$ Γ_{126}/Γ
(0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1 \times 10^{-3}$	90	1 LEES	13L BABR	$\gamma(1S) \rightarrow \gamma X$

¹ For a narrow, CP -odd pseudoscalar, A^0 , searched for in 14 hadronic decay modes with invariant mass $1.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$. Measured 90% CL limits as a function of m_{A^0} range from 10^{-5} to 10^{-3} .

LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$				Γ_{127}/Γ
<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.9	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \mu^\mp$
$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{128}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.7 \times 10^{-6}$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^\pm \tau^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.0 \times 10^{-6}$	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$
$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{129}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.7	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \tau^\mp$
$\Gamma(\gamma e^\pm \mu^\mp)/\Gamma_{\text{total}}$				Γ_{130}/Γ
<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.2	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \mu^\mp$
$\Gamma(\gamma \mu^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{131}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.1	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma \mu^\pm \tau^\mp$
$\Gamma(\gamma e^\pm \tau^\mp)/\Gamma_{\text{total}}$				Γ_{132}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.5	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \tau^\mp$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$				Γ_{133}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.0	90	AUBERT	09AX BABR	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<39	90	RUBIN	07	CLEO $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
<25	90	TAJIMA	07	BELL $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

 $\Upsilon(1S)$ REFERENCES

SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
JIA	22	PRL 128 081804	S. Jia <i>et al.</i>	(BELLE Collab.)
PATRA	22	JHEP 2205 095	S. Patra <i>et al.</i>	(BELLE Collab.)
KATRENKO	20	PRL 124 122001	P. Katreko <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)

LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
BESSON	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
BESSON	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
ANTREASYAN	90C	PL B251 204	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueler, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, RÖCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)

BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)
