


$f'_2(1525)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f'_2(1525)$ MASS**VALUE (MeV)DOCUMENT ID

1517.3 \pm 2.4 OUR AVERAGE Includes data from the 6 datablocks that follow this one.
Error includes scale factor of 2.8. See the ideogram below.

 $f'_2(1525)$ MASS (MeV)**PRODUCED BY PION BEAM**VALUE (MeV)EVTSDOCUMENT IDTECNCOMMENT

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

1521 \pm 13	TIKHOMIROV 03	SPEC	40.0 π^- C \rightarrow	$K_S^0 K_S^0 K_L^0 X$
1547 $^{+10}_{-2}$	¹ LONGACRE 86	MPS	22 π^- p \rightarrow	$K_S^0 K_S^0 n$

1496^{+9}_{-8}	² CHABAUD	81	ASPK	$6 \pi^- p \rightarrow K^+ K^- n$
1497^{+8}_{-9}	CHABAUD	81	ASPK	$18.4 \pi^- p \rightarrow K^+ K^- n$
1492 ± 29	GORLICH	80	ASPK	$17 \pi^- p$ polarized $\rightarrow K^+ K^- n$
1502 ± 25	³ CORDEN	79	OMEG	$12\text{--}15 \pi^- p \rightarrow \pi^+ \pi^- n$
1480	14	66	HBC	$6.0 \pi^- p \rightarrow K_S^0 K_S^0 n$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

³ From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.


PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1518.0 \pm 2.7 OUR AVERAGE Includes data from the datablock that follows this one.
Error includes scale factor of 2.9. See the ideogram below.

1526.8 ± 4.3	ASTON	88D	LASS	$11 K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1504 ± 12	BOLONKIN	86	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 Y$
1529 ± 3	ARMSTRONG	83B	OMEG	$18.5 K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	81B	HBC	$4.2 K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	81	HBC	$8.25 K^- p \rightarrow \Lambda K\bar{K}$
1522 ± 6	123	77	HBC	$4.15 K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	77	OMEG	$10 K^- p \rightarrow K^+ K^-(\Lambda, \Sigma)$
1527 ± 3	120	76C	ASPK	$13 K^- p \rightarrow K^+ K^-(\Lambda, \Sigma)$
1519 ± 7	100	72B	HBC	$3.9, 4.6 K^- p \rightarrow K\bar{K}(\Lambda, \Sigma)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1514 ± 8	61	BINON	07	GAMS $32.5 K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
1513 ± 10	¹ BARKOV	99	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 Y$

¹ Systematic errors not estimated.

PRODUCED BY K^\pm BEAM (MeV)**PRODUCED IN e^+e^- ANNIHILATION AND PARTICLE DECAYS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

1514 ± 4 OUR AVERAGE Error includes scale factor of 3.6. See the ideogram below.

1515.4 ± 2.5	$^{+3.2}_{-7.6}$	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1522.2 ± 2.8	$^{+5.3}_{-2.0}$		AAIJ	13AN LHCb	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
1513 ± 5	$^{+4}_{-10}$	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1525.3 ± 1.2	$^{+3.7}_{-1.4}$		UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
1521 ± 5			ABLIKIM	05 BES2	$J/\psi \rightarrow \phi K^+ K^-$
1518 ± 1	± 3		ABE	04 BELL	$10.6 \frac{e^+ e^-}{e^+ e^- K^+ K^-} \rightarrow$
1519 ± 2	$^{+15}_{-5}$		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
1523 ± 6		331	² ACCIARRI	01H L3	$91, 183-209 \frac{e^+ e^-}{e^+ e^- K_S^0 K_S^0} \rightarrow$
1535 ± 5	± 4		ABREU	96C DLPH	$Z^0 \rightarrow K^+ K^- + X$
1516 ± 5	$^{+9}_{-15}$		BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$

1531.6 \pm 10.0	AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
1496 \pm 2	3 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$
1525 \pm 10 \pm 10	BALTRUSAIT...87	..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
1518 \pm 3	4 KLEMPPT	22	RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0,$ $\gamma K_S^0 K_S^0$
1503 \pm 11	5 RODAS	22	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
1532 \pm 3 \pm 6	6,7 DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
1557 \pm 9 \pm 3	6,7 DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
1526 \pm 7	8 LEES	14H	BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
1523 \pm 5	9 SCHEGELSKY	06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1515 \pm 5	10 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$

¹ From partial wave analysis including all possible combinations of 0⁺⁺, 2⁺⁺, and 4⁺⁺ resonances.

² Supersedes ACCIARRI 95J.

³ From an analysis including interference with $f_0(1710)$.

⁴ Fit of the tensor partial waves from BES3 in the multipole basis.

⁵ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABELIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABELIKIM 18AA).


⁶ Using CLEO-c data but not authored by the CLEO Collaboration.

⁷ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 73$ MeV.

⁸ From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.

⁹ From analysis of L3 data at 91 and 183–209 GeV.

¹⁰ From an analysis ignoring interference with $f_0(1710)$.



PRODUCED IN $\bar{p}p$ ANNIHILATION

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.			

1512 \pm 4 OUR AVERAGE

1513 \pm 4	AMSLER	06	CBAR $0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1508 \pm 9	¹ AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
1495.0 \pm 1.1 \pm 8.1	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
1530 \pm 12	³ ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$

¹ T-matrix pole.² T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).³ 4-poles, 5-channel K matrix fit.**CENTRAL PRODUCTION**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.			

1515 \pm 15	BARBERIS	99	OMEG $450 pp \rightarrow p_s p_f K^+ K^-$
---------------------------------	----------	----	---

PRODUCED IN $e p$ COLLISIONS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
The data in this block is included in the average printed for a previous datablock.				

1512 \pm 3 $^{+1.4}_{-0.5}$
--

¹ CHEKANOV	08	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
-----------------------	----	--------------------------------------

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$


1537 $^{+9}_{-8}$	84	² CHEKANOV	04	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
-------------------	----	-----------------------	----	--------------------------------------

¹ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.² Systematic errors not estimated. **$f'_2(1525)$ WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
--------------------	--------------------

72 $^{+7}_{-6}$ OUR FIT

84.4 \pm 2.7 OUR AVERAGE Includes data from the 6 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below.



$f'_2(1525)$ WIDTH (MeV)

PRODUCED BY PION BEAM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

• • • We do not use the following data for averages, fits, limits, etc. • • •

102±42	TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
108 ⁺⁵ ₋₂	¹ LONGACRE	86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
69 ⁺²² ₋₁₆	² CHABAUD	81	ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
137 ⁺²³ ₋₂₁	CHABAUD	81	ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
150 ⁺⁸³ ₋₅₀	GORLICH	80	ASPK	17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$

165 ± 42	³ CORDEN	79	OMEG	12–15 $\pi^- p \rightarrow \pi^+ \pi^- n$
92^{+39}_{-22}	⁴ POLYCHRO...	79	STR...	$7 \pi^- p \rightarrow n K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

³ From an amplitude analysis where the $f'_2(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.

⁴ From a fit to the D with $f_2(1270)-f'_2(1525)$ interference. Mass fixed at 1516 MeV.

PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

82 \pm 6 OUR AVERAGE

90 ± 12	ASTON	88D	LASS	$11 K^- p \rightarrow K_S^0 K_S^0 \Lambda$
73 ± 18	BOLONKIN	86	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 Y$
83 ± 15	ARMSTRONG	83B	OMEG	$18.5 K^- p \rightarrow K^- K^+ \Lambda$
85 ± 16	650	AGUILAR....	81B	HBC $4.2 K^- p \rightarrow \Lambda K^+ K^-$
80^{+14}_{-11}	572	ALHARRAN	81	HBC $8.25 K^- p \rightarrow \Lambda K\bar{K}$
72 ± 25	166	EVANGELIS...	77	OMEG $10 K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69 ± 22	100	AGUILAR....	72B	HBC $3.9, 4.6 K^- p \rightarrow K\bar{K} (\Lambda, \Sigma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

92^{+25}_{-16}	61	BINON	07	GAMS $32.5 K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
75 ± 20		¹ BARKOV	99	SPEC $40 K^- p \rightarrow K_S^0 K_S^0 Y$
62^{+19}_{-14}	123	BARREIRO	77	HBC $4.15 K^- p \rightarrow \Lambda K_S^0 K_S^0$
61 ± 8	120	BRANDENB...	76C	ASPK $13 K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$

¹ Systematic errors not estimated.

PRODUCED IN $e^+ e^-$ ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

86 \pm 4 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.

$64.0 \pm 4.3^{+2.0}_{-6.1}$	126K	ABLIKIM	23M	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
$84 \pm 6^{+10}_{-5}$		AAIJ	13AN	LHCb $\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
$75^{+12}_{-10} \pm 16_{-8}$	5.5k	¹ ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
$82.9^{+2.1+3.3}_{-2.2-2.0}$		UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
77 ± 15		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi K^+ K^-$
$82 \pm 2 \pm 3$		ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
$75 \pm 4^{+15}_{-5}$		BAI	03G	BES $J/\psi \rightarrow \gamma K\bar{K}$
100 ± 15	331	² ACCIARRI	01H	L3 $91, 183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$

60	± 20	± 19	ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$
60	± 23	$+13$ -20	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
103	± 30		AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
100	± 3		³ FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$
85	± 35		BALTRUSAIT...87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

78	± 6		⁴ KLEMPPT	22	RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0,$ $\gamma K_S^0 K_S^0$
84	± 15		⁵ RODAS	22	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi,$ $KK)$
37	± 12	29	⁶ LEES	14H	BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
104	± 10	870	⁷ SCHEGELSKY	06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
62	± 10		⁸ FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$

¹ From partial wave analysis including all possible combinations of 0++, 2++, and 4++ resonances.

² Supersedes ACCIARRI 95J.

³ From an analysis including interference with $f_0(1710)$.


⁴ Fit of the tensor partial waves from BES3 in the multipole basis.

⁵ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).

⁶ From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.

⁷ From analysis of L3 data at 91 and 183–209 GeV.

⁸ From an analysis ignoring interference with $f_0(1710)$.



PRODUCED IN $\bar{p}p$ ANNIHILATION

<i>VALUE (MeV)</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
The data in this block is included in the average printed for a previous datablock.			

77 ± 5 OUR AVERAGE

76 ± 6	AMSLER	06	CBAR 0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
79 ± 8	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
104.8 ± 0.9 ± 9.8	ALBRECHT	20	RVUE 0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
128 ± 20	ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$

¹ T-matrix pole.² T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).³ K-matrix, 4-poles, 5-channel fit.**CENTRAL PRODUCTION**

<i>VALUE (MeV)</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
The data in this block is included in the average printed for a previous datablock.			

70±25	BARBERIS	99	OMEG 450 $p p \rightarrow p_S p_f K^+ K^-$
--------------	----------	----	--

PRODUCED IN $e p$ COLLISIONS

<i>VALUE (MeV)</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
The data in this block is included in the average printed for a previous datablock.				

83± 9⁺⁵₋₄¹ CHEKANOV 08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

50⁺³⁴₋₂₂ 84 ² CHEKANOV 04 ZEUS $e p \rightarrow K_S^0 K_S^0 X$ ¹ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.² Systematic errors not estimated. **$f'_2(1525)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	(88.8 ± 2.2) %
Γ_2 $\eta\eta$	(10.3 ± 2.2) %
Γ_3 $\pi\pi$	(8.2 ± 1.5) × 10 ⁻³
Γ_4 $K\bar{K}^*(892) + \text{c.c.}$	
Γ_5 $\pi K\bar{K}$	
Γ_6 $\pi\pi\eta$	
Γ_7 $\pi^+\pi^+\pi^-\pi^-$	
Γ_8 $\gamma\gamma$	(1.12 ± 0.15) × 10 ⁻⁶

CONSTRAINED FIT INFORMATION

An overall fit to 2 partial widths, a combination of partial widths obtained from integrated cross sections, and 3 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 14.2$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-5	-1		
x_8	0	0	1	
Γ	-28	28	-1	-62
	x_1	x_2	x_3	x_8

	Mode	Rate (MeV)
Γ_1	$K\bar{K}$	64^{+6}_{-5}
Γ_2	$\eta\eta$	7.4 ± 1.9
Γ_3	$\pi\pi$	0.59 ± 0.12
Γ_8	$\gamma\gamma$	$(8.1 \pm 0.9) \times 10^{-5}$

$f'_2(1525)$ PARTIAL WIDTHS

$\Gamma(K\bar{K})$	Γ_1
<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

64^{+6}_{-5} OUR FIT

63^{+6}_{-5} ¹ LONGACRE 86 MPS $22\pi^- p \rightarrow K_S^0 K_S^0 n$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

$\Gamma(\eta\eta)$	Γ_2
<u>VALUE (MeV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

7.4 ± 1.9 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0 ± 0.8	870	¹ SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
24^{+3}_{-1}		² LONGACRE 86 MPS $22\pi^- p \rightarrow K_S^0 K_S^0 n$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f'_2(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

$\Gamma(\pi\pi)$					Γ_3
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.59±0.12 OUR FIT					
1.4 ±0.5		¹ LONGACRE 86	MPS	$22 \pi^- p \rightarrow K_S^0 K_S^0 n$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.2 ±0.2	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f'_2(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.

$\Gamma(\gamma\gamma)$					Γ_8
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.081±0.009 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.13 ±0.03	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f'_2(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.746±0.002 ^{+0.166} _{-0.162}		¹ ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$	

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

$f'_2(1525) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_8/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.072 ±0.007 OUR FIT					
0.072 ±0.007 OUR AVERAGE					
0.048 ±0.007	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
0.0564±0.0048±0.0116	ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.076 ±0.006	1 ACCIARRI	01H	L3	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.067 ±0.008	² ALBRECHT	90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.11 ±0.03	BEHREND	89C	CELL	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.10 ±0.04	BERGER	88	PLUT	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.12 ±0.07	² AIHARA	86B	TPC	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.11 ±0.02	² ALTHOFF	83	TASS	$e^+ e^- \rightarrow e^+ e^- K\bar{K}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0314±0.0050±0.0077	³ ALBRECHT	90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	

¹ Supersedes ACCIARRI 95J. From analysis of L3 data at 91 and 183–209 GeV,² Using an incoherent background.³ Using a coherent background.

$f'_2(1525)$ BRANCHING RATIOS

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.059±0.003±0.026	¹ ALBRECHT 20 RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$	
seen	UEHARA 10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$	
0.10 ±0.03	² PROKOSHKIN 91 GAM4	300 $\pi^-p \rightarrow \pi^-p\eta\eta$	

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).² Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$

Γ_2/Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.116±0.028 OUR FIT					
0.115±0.028 OUR AVERAGE					
0.119±0.015±0.036	61	¹ BINON	07 GAMS	32.5 $K^-p \rightarrow \eta\eta(\Lambda/\Sigma^0)$	
0.11 ±0.04		² PROKOSHKIN 91	GAM4 300	$\pi^-p \rightarrow \pi^-p\eta\eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.14	90	BARBERIS	00E	450 $p\bar{p} \rightarrow p_f\eta\eta p_s$	
< 0.50		BARNES	67 HBC	4.6,5.0 K^-p	

¹ Using the compilation of the cross sections for $f'_2(1525)$ production in K^-p collisions from ASTON 88D.² Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
0.82±0.16 OUR FIT				
0.75±0.16 OUR AVERAGE				
0.7 ±0.2		COSTA 80	OMEG 10	$\pi^-p \rightarrow K^+K^-n$
2.7 ^{+7.1} _{-1.3}		¹ GORLICH 80	ASPK 17,18	π^-p
0.75±0.25		1,2 MARTIN 79	RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.4 ±1.5 ±1.0		³ ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
< 6	95	AGUILAR-...	81B HBC	4.2 $K^-p \rightarrow \Lambda K^+K^-$
19 ±3		CORDEN 79	OMEG 12–15	$\pi^-p \rightarrow \pi^+\pi^-n$
< 4.5	95	BARREIRO 77	HBC 4.15	$K^-p \rightarrow \Lambda K_S^0 K_S^0$
1.2 ±0.4		¹ PAWLICKI 77	SPEC 6	$\pi N \rightarrow K^+K^-N$
< 6.3	90	BRANDENB... 76C	ASPK 13	$K^-p \rightarrow K^+K^-(\Lambda, \Sigma)$
< 0.86		¹ BEUSCH 75B	OSPK 8.9	$\pi^-p \rightarrow K^0\bar{K}^0n$

¹ Assuming that the $f'_2(1525)$ is produced by an one-pion exchange production mechanism.

² MARTIN 79 uses the PAWLICKI 77 data with different input value of the $f'_2(1525) \rightarrow K\bar{K}$ branching ratio.

³ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0092 ± 0.0018 OUR FIT			
0.075 ± 0.035	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$

$[\Gamma(K\bar{K}^*(892)+c.c.) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.35	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$
<0.4	67	AMMAR	67	HBC

$\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.41	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$
<0.3	67	AMMAR	67	HBC

$\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.32	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$

Γ_3/Γ_1

Γ_6/Γ_1

Γ_7/Γ_1

$f'_2(1525)$ REFERENCES

ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
KLEMPT	22	PL B830 137171	E. Klempert <i>et al.</i>	(BONN)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
BINON	07	PAN 70 1713	F. Binon <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 70 1758.		
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)

BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARKOV	99	JETPL 70 248	B.P. Barkov <i>et al.</i>	
		Translated from ZETFP 70 242.		
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
ACCIARRI	95J	PL B363 118	M. Acciarri <i>et al.</i>	(L3 Collab.)
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)
		Translated from DANS 316 900.		
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BERGER	88	ZPHY C37 329	C. Berger <i>et al.</i>	(PLUTO Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAZO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAZO, CLER, FRAS+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
BOLONKIN	86	SJNP 43 776	B.V. Bolonkin <i>et al.</i>	(ITEP) JP
		Translated from YAF 43 1211.		
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i>	(TASSO Collab.)
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
AGUILAR-...	81B	ZPHY C8 313	M. Aguilar-Benitez <i>et al.</i>	(CERN, CDEF+)
ALHARRAN	81	NP B191 26	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
BARREIRO	77	NP B121 237	F. Barreiro <i>et al.</i>	(CERN, AMST, NIJM+)
EVANGELIS...	77	NP B127 384	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
PAWLIKCI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJP
BRANDENB...	76C	NP B104 413	G.W. Brandenburg <i>et al.</i>	(SLAC)
BEUSCH	75B	PL 60B 101	W. Beusch <i>et al.</i>	(CERN, ETH)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
AMMAR	67	PRL 19 1071	R. Ammar <i>et al.</i>	(NWES, ANL) JP
BARNES	67	PRL 19 964	V.E. Barnes <i>et al.</i>	(BNL, SYRA) IJPC
CRENNELL	66	PRL 16 1025	D.J. Crennell <i>et al.</i>	(BNL) I
