Average Hadron Multiplicities in Hadronic e^+e^- Annihilation Events

Table 36.1: Average hadron multiplicity per e^+e^- annihilation event at $\sqrt{s} \approx 10$, 29–35, and 91 GeV. The rates given include decay products from resonances with $c\tau < 10$ cm, and include charge conjugated states. (Updated July 1995 by O. Biebel.)

Particle	$\sqrt{s} \approx 10$ GeV		$\sqrt{s} = 29$–35 GeV		$\sqrt{s} = 91$ GeV	
Pseudoscalar mesons:						
π^+	6.6	$\pm$ 0.2	10.3	$\pm$ 0.4	17.1	$\pm$ 0.4
π^0	3.2	$\pm$ 0.3	5.83	$\pm$ 0.28	9.18	$\pm$ 0.73
K^+	0.90	$\pm$ 0.04	1.48	$\pm$ 0.09	2.39	$\pm$ 0.12
K^0	0.91	$\pm$ 0.05	1.48	$\pm$ 0.07	2.01	$\pm$ 0.04
η	0.20	$\pm$ 0.04	0.61	$\pm$ 0.07	0.95	$\pm$ 0.11
$\eta\prime(958)$	0.03	$\pm$ 0.01	0.26	$\pm$ 0.10	0.17	$\pm$ 0.06
D^+	0.16	$\pm$ 0.03	0.17	$\pm$ 0.03	0.20	$\pm$ 0.03
D^0	0.37	$\pm$ 0.06	0.45	$\pm$ 0.07	0.40	$\pm$ 0.06
D_s^+	0.13	$\pm$ 0.02	0.45	$\pm$ 0.20 [a]		—
B^+, B_d^0		—		—	0.165	$\pm$ 0.026 [b]
B_s^0		—		—	0.057	$\pm$ 0.013 [b]
Scalar mesons:						
$f_0(980)$	0.024	$\pm$ 0.006	0.05	$\pm$ 0.02 [c]	0.14	$\pm$ 0.06 [d]
Vector mesons:						
$\rho(770)^0$	0.35	$\pm$ 0.04	0.81	$\pm$ 0.08	1.21	$\pm$ 0.16
$\omega(782)$	0.30	$\pm$ 0.08		—		—
$K^*(892)^+$	0.27	$\pm$ 0.03	0.64	$\pm$ 0.05	0.715	$\pm$ 0.059
$K^*(892)^0$	0.29	$\pm$ 0.03	0.56	$\pm$ 0.06	0.742	$\pm$ 0.042
$\phi(1020)$	0.044	$\pm$ 0.003	0.085	$\pm$ 0.011	0.100	$\pm$ 0.008
$D^*(2010)^+$	0.22	$\pm$ 0.04	0.43	$\pm$ 0.07	0.180	$\pm$ 0.013
$D^*(2007)^0$	0.23	$\pm$ 0.06	0.27	$\pm$ 0.11		—
B^* [e]		—		—	0.288	$\pm$ 0.026
$J/\psi(1S)$		—		—	0.0054	$\pm$ 0.0005 [f]
$\psi(2S)$		—		—	0.0023	$\pm$ 0.0011 [f]
Pseudovector mesons:						
$\chi_{c1}(1P)$		—		—	0.0087	$\pm$ 0.0028 [f]
Tensor mesons:						
$f_2(1270)$	0.09	$\pm$ 0.02	0.14	$\pm$ 0.04	0.31	$\pm$ 0.12
$K_2^*(1430)^+$		—	0.09	$\pm$ 0.03		—
$K_2^*(1430)^0$		—	0.12	$\pm$ 0.06	0.19	$\pm$ 0.07 [g]
B^{**} [h]		—		—	0.12	$\pm$ 0.24
Baryons:						
p	0.253	$\pm$ 0.016	0.640	$\pm$ 0.050	0.964	$\pm$ 0.102
Λ	0.080	$\pm$ 0.007	0.205	$\pm$ 0.010	0.368	$\pm$ 0.014
Σ^0	0.023	$\pm$ 0.008		—		—
$\Sigma^\pm$		—		—	0.170	$\pm$ 0.063
Ξ^-	0.0059	$\pm$ 0.0007	0.0176	$\pm$ 0.0027	0.0227	$\pm$ 0.0018
$\Delta(1232)^{++}$	0.040	$\pm$ 0.010		—	0.022	$\pm$ 0.06
$\Sigma(1385)^-$	0.006	$\pm$ 0.002	0.017	$\pm$ 0.004		—
$\Sigma(1385)^+$	0.005	$\pm$ 0.001	0.017	$\pm$ 0.004		—
$\Sigma(1385)^\pm$	0.0106	$\pm$ 0.0020	0.033	$\pm$ 0.008	0.0380	$\pm$ 0.0062
$\Xi(1530)^0$	0.0015	$\pm$ 0.0006		—	0.0048	$\pm$ 0.0005
Ω^-	0.0007	$\pm$ 0.0004	0.014	$\pm$ 0.007	0.0050	$\pm$ 0.0015
Λ_c^+	0.100	$\pm$ 0.030 [i]	0.110	$\pm$ 0.050		—
Λ_b^0		—		—	0.031	$\pm$ 0.016
$\Sigma_c^{++}, \Sigma_c^0$	0.014	$\pm$ 0.007		—		—

All average multiplicites are per hadronic e^+e^- annihilation event.

(a) B($D_s \to \eta\pi,\ \eta'\pi$) has been used (RPP 1994).

(b) The Standard Model B($Z \to b\overline{b}$) = 0.217 was used.

(c) $x_p = p/p_{\rm beam} > 0.1$ only.

(d) Extrapolation to the unobserved region using the shape predicted by JETSET.

(e) Any charge state (*i.e.*, B_d^*, B_u^*, or B_s^*).

(f) B($Z \to$ hadrons) = 0.699 has been used (RPP 1994).

(g) $x_E = E[K_2^*(1430)0]/E_{\rm beam} < 0.3$ only.

(h) Any charge state (*i.e.*, B_d^{**}, B_u^{**}, or B_s^{**}).

(i) The value was taken from the cross section of the $\Lambda_c^+ \to p\pi K$, assuming the branching fraction to be (3.2$\pm$0.7)% (RPP 1992).

References:

RPP92: Phys. Rev. **D45** (1992) and references therein

RPP94: Phys. Rev. **D50**, 1173 (1994) and references therein

A. De Angelis, J. Phys. **G19**, 1233 (1993) and references therein

R. Marshall, Rept. on Prog. in Phys. **52**, 1329 (1989)

ALEPH: D. Buskulic *et al.*, Phys. Lett. **B295**, 396 (1992)

ALEPH: D. Buskulic *et al.*, Z. Phys. **C64**, 361 (1994)

ARGUS: A. Albrecht *et al.*, Z. Phys. **C61**, 1 (1994)

ARGUS: H. Albrecht *et al.*, Z. Phys. **C58**, 199 (1993)

ARGUS: H. Albrecht *et al.*, Z. Phys. **C54**, 1 (1992)

ARGUS: H. Albrecht *et al.*, Z. Phys. **C46**, 15 (1990)

ARGUS: H. Albrecht *et al.*, Z. Phys. **C44**, 547 (1989)

ARGUS: H. Albrecht *et al.*, Phys. Lett. **B230**, 169 (1989)

CELLO: H.J. Behrend *et al.*, Z. Phys. **C47**, 1 (1990)

CELLO: H.J. Behrend *et al.*, Z. Phys. **C46**, 397 (1990)

CLEO: D. Bortoletto *et al.*, Phys. Rev. **D37**, 1719 (1988)

Crystal Ball: Ch. Bieler *et al.*, Z. Phys. **C49**, 225 (1991)

DELPHI: P. Abreu *et al.*, Nucl. Phys. **B**, CERN PPE/95-28

DELPHI: P. Abreu *et al.*, Z. Phys. **C**, CERN PPE/95-39

DELPHI: P. Abreu *et al.*, Z. Phys. **C**, CERN PPE/95-53

DELPHI: P. Abreu *et al.*, Phys. Lett. **B345**, 598 (1995)

DELPHI: P. Abreu *et al.*, Z. Phys. **C65**, 587 (1995)

DELPHI: P. Abreu *et al.*, Phys. Lett. **B341**, 109 (1994)

DELPHI: P. Abreu *et al.*, Z. Phys. **C61**, 407 (1994)

DELPHI: P. Abreu *et al.*, Z. Phys. **C59**, 533 (1993)

DELPHI: P. Abreu *et al.*, Z. Phys. **C57**, 181 (1993)

HRS: M. Derrick *et al.*, Phys. Rev. **D35**, 2639 (1987)

HRS: S. Abachi *et al.*, Phys. Rev. Lett. **57**, 1990 (1986)

JADE: D.D. Pietzl *et al.*, Z. Phys. **C46**, 1 (1990)

JADE: W. Bartel *et al.*, Z. Phys. **C20**, 187 (1983)

L3: M. Acciarri *et al.*, Phys. Lett. **B345**, 589 (1995)

L3: M. Acciarri *et al.*, Phys. Lett. **B328**, 223 (1994)

L3: O. Adriani *et al.*, Phys. Lett. **B317**, 467 (1993)

MARK II: G. Wormser *et al.*, Phys. Rev. Lett. **61**, 1057 (1988)

MARK II: H. Schellman *et al.*, Phys. Rev. **D31**, 3013 (1985)

OPAL: R. Akers *et al.*, Z. Phys. **C**, CERN PPE/95-12

OPAL: R. Akers *et al.*, Z. Phys. **C**, CERN PPE/95-24

OPAL: R. Akers *et al.*, Z. Phys. **C**, CERN PPE/95-27

OPAL: G. Alexander *et al.*, Phys. Lett. **B**, CERN PPE/95-99

OPAL: R. Akers *et al.*, Z. Phys. **C**, CERN PPE/94-217

OPAL: R. Akers *et al.*, Z. Phys. **C63**, 181 (1994)

OPAL: P.D. Acton *et al.*, Z. Phys. **C56**, 521 (1992)

OPAL: P.D. Acton *et al.*, Phys. Lett. **B291**, 503 (1992)

OPAL: G. Alexander *et al.*, Phys. Lett. **B266**, 485 (1991)

PLUTO: Ch. Berger *et al.*, Phys. Lett. **104B**, 79 (1981)

TASSO: H. Aihara *et al.*, Z. Phys. **C27**, 27 (1985)

TPC: H. Aihara *et al.*, Phys. Rev. Lett. **53**, 2378 (1984)