W

$J = 1$

W MASS

OUR FIT uses the W and Z mass, mass difference, and mass ratio measurements.

Because of the high current interest, we mention the following preliminary results here, but do not average them or include them in the Listings or Tables. Performing an overall fit (assuming the Standard Model) of published and unpublished (CDF and DØ) collider results ($M_W = 80.33 \pm 0.15$ GeV), of the νN results ($1 - M_W^2/M_Z^2 = 0.2257 \pm 0.0047$) and of published and unpublished LEP and SLD preliminary electroweak results (as of end of March 1996), the W mass is fitted to be $M_W = (80.350 \pm 0.042^{+0.016}_{-0.025})$ GeV (the second errors correspond to varying the Higgs mass in the interval 60–1000 GeV).

VALUE (GeV)	EVTS	DOCUMENT ID		TECN	COMMENT
80.33 ± 0.15 OUR FIT					
80.32 ± 0.19 OUR AVERAGE					Error includes scale factor of 1.2.
80.410 ± 0.180	8986	[1] ABE	95P	CDF	$E_{cm}^{p\overline{p}} = 1800$ GeV
79.91 ± 0.39	1722	[2] ABE	90G	CDF	$E_{cm}^{p\overline{p}} = 1800$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
80.84 ± 0.22 ±0.83	2065	[3] ALITTI	92B	UA2	See W/Z ratio below
80.79 ± 0.31 ±0.84		[4] ALITTI	90B	UA2	$E_{cm}^{p\overline{p}} = 546,630$ GeV
80.0 ± 3.3 ±2.4	22	[5] ABE	89I	CDF	$E_{cm}^{p\overline{p}} = 1800$ GeV
82.7 ± 1.0 ±2.7	149	[6] ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}} = 546,630$ GeV
$81.8 \pm {}^{6.0}_{5.3}$ ±2.6	46	[7] ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}} = 546,630$ GeV
89 ± 3 ±6	32	[8] ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}} = 546,630$ GeV
80.2 ± 0.6 ±1.4	251	[9] ANSARI	87	UA2	Repl. by ALITTI 90B
81.2 ± 1.0 ±1.4	119	[9] APPEL	86	UA2	Repl. by ANSARI 87
$83.5 \pm {}^{1.1}_{1.0}$ ±2.7	86	[10] ARNISON	86	UA1	Repl. by ALBAJAR 89
$81. \pm {}^{6.}_{7.}$	14	[11] ARNISON	84D	UA1	Repl. by ALBAJAR 89
83.1 ± 1.9 ±1.3	37	BAGNAIA	84	UA2	Repl. by ALITTI 90B
81. ± 5.	6	ARNISON	83	UA1	Repl. by ARNISON 83D
80.9 ± 2.9	27	ARNISON	83D	UA1	Repl. by ARNISON 86
81.0 ± 2.8		BAGNAIA	83	UA2	Repl. by BAGNAIA 84
$80. \pm {}^{10.}_{6.}$	4	BANNER	83B	UA2	Repl. by ALITTI 90B

[1] ABE 95P use 3268 $W \to \mu\nu_\mu$ events to find $M = 80.310 \pm 0.205 \pm 0.130$ GeV and 5718 $W \to e\nu_e$ events to find $M = 80.490 \pm 0.145 \pm 0.175$ GeV. The result given here combines these while accounting for correlated uncertainties.

[2] ABE 90G result from $W \to e\nu$ is $79.91 \pm 0.35 \pm 0.24 \pm 0.19$(scale) GeV and from $W \to \mu\nu$ is $79.90 \pm 0.53 \pm 0.32 \pm 0.08$(scale) GeV.

[3] ALITTI 92B result has two contributions to the systematic error (± 0.83); one (± 0.81) cancels in m_W/m_Z and one (± 0.17) is noncancelling. These were added in quadrature. We choose the ALITTI 92B value without using the LEP m_Z value, because we perform our own combined fit.

[4] There are two contributions to the systematic error (± 0.84): one (± 0.81) which cancels in m_W/m_Z and one (± 0.21) which is non-cancelling. These were added in quadrature.

[5] ABE 89I systematic error dominated by the uncertainty in the absolute energy scale.

[6] ALBAJAR 89 result is from a total sample of 299 $W \to e\nu$ events.

[7] ALBAJAR 89 result is from a total sample of 67 $W \to \mu\nu$ events.

[8] ALBAJAR 89 result is from $W \to \tau\nu$ events.

[9] There are two contributions to the systematic error (± 1.4): one (± 1.3) which cancels in m_W/m_Z and one (± 0.5) which is non-cancelling. These were added in quadrature.

[10] This is enhanced subsample of 172 total events.

[11] Using $W^\pm \to \mu^\pm\nu$.

W/Z MASS RATIO

The fit uses the W and Z mass, mass difference, and mass ratio measurements.

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.8810±0.0016 OUR FIT					
0.8813±0.0036±0.0019	156	[12] ALITTI	92B	UA2	$E_{\rm cm}^{p\overline{p}}$= 630 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.8831±0.0048±0.0026		[12] ALITTI	90B	UA2	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV

[12] Scale error cancels in this ratio.

$m_Z - m_W$

The fit uses the W and Z mass, mass difference, and mass ratio measurements.

VALUE (GeV)	DOCUMENT ID		TECN	COMMENT
10.85±0.15 OUR FIT				
10.4 ±1.4 ±0.8	ALBAJAR	89	UA1	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.3 ±1.3 ±0.9	ANSARI	87	UA2	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV

$m_{W^+} - m_{W^-}$

Test of CPT invariance.

VALUE (GeV)	EVTS	DOCUMENT ID		TECN	COMMENT
−0.19±0.58	1722	ABE	90G	CDF	$E_{\rm cm}^{p\overline{p}}$= 1800 GeV

W WIDTH

The CDF and DØ widths labelled "extracted value" are obtained by measuring $R = [\sigma(W)/\sigma(Z)]\,[\Gamma(W \to e\nu_e)]/(\mathrm{B}(Z \to ee)\Gamma(W))$ where the bracketed quantities can be calculated with plausible reliability. $\Gamma(W)$ is then extracted by using a value of $\mathrm{B}(Z \to ee)$ measured at LEP. The UA1 and UA2 widths used $R = [\sigma(W)/\sigma(Z)]\,[\Gamma(W \to e\nu_e)/\Gamma(Z \to ee)]\,\Gamma(Z)/\Gamma(W)$ and the measured value of $\Gamma(Z)$. The Standard Model prediction is 2.067 ± 0.021 (ROSNER 94).

VALUE (GeV)	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
2.07 ±0.06 OUR AVERAGE						
2.044±0.093		13k	[13] ABACHI	95D	D0	Extracted value
2.11 ±0.28 ±0.16		58	[14] ABE	95C	CDF	Direct meas.
2.064±0.060±0.059			[15] ABE	95W	CDF	Extracted value
$2.10\ ^{+0.14}_{-0.13}$ ±0.09		3559	[16] ALITTI	92	UA2	Extracted value
$2.18\ ^{+0.26}_{-0.24}$ ±0.04			[17] ALBAJAR	91	UA1	Extracted value
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.16 ±0.17			[18] ABE	92I	CDF	Repl. by ABE 95W
2.12 ±0.20			[19] ABE	90	CDF	Repl. by ABE 92I
2.30 ±0.19 ±0.06			[20] ALITTI	90C	UA2	Extracted value
<5.4	90	149	[21] ALBAJAR	89	UA1	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV
$2.8\ ^{+1.4}_{-1.5}$ ±1.3		149	[21] ALBAJAR	89	UA1	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV
<7	90	251	ANSARI	87	UA2	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV
<7	90	119	APPEL	86	UA2	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV
<6.5	90	86	[22] ARNISON	86	UA1	Repl. by ALBAJAR 89
<7	90	27	ARNISON	83D	UA1	Repl. by ARNISON 86

[13] ABACHI 95D measured $R = 10.90 \pm 0.49$ and used the measured value $\mathrm{B}(Z \to \ell\ell) = (3.367 \pm 0.006)\%$ from LEP.

[14] ABE 95C use the tail of the transverse mass distribution of $W \to e\nu_e$ decays.

[15] ABE 95W measured $R = 10.90 \pm 0.32 \pm 0.29$. They use m_W=80.23 ± 0.18 GeV, $\sigma(W)/\sigma(Z) = 3.35 \pm 0.03$, $\Gamma(W \to e\nu) = 225.9 \pm 0.9$ MeV, $\Gamma(Z \to e^+e^-) = 83.98 \pm 0.18$ MeV, and $\Gamma(Z) = 2.4969 \pm 0.0038$ GeV.

[16] ALITTI 92 measured $R = 10.4^{+0.7}_{-0.6} \pm 0.3$. The values of $\sigma(Z)$ and $\sigma(W)$ come from $O(\alpha_s^2)$ calculations using $m_W = 80.14 \pm 0.27$ GeV, and $m_Z = 91.175 \pm 0.021$ GeV along with the corresponding value of $\sin^2\theta_W = 0.2274$. They use $\sigma(W)/\sigma(Z) = 3.26 \pm 0.07 \pm 0.05$ and $\Gamma(Z) = 2.487 \pm 0.010$ GeV.

[17] ALBAJAR 91 measured $R = 9.5^{+1.1}_{-1.0}$ (stat. + syst.). $\sigma(W)/\sigma(Z)$ is calculated in QCD at the parton level using $m_W = 80.18 \pm 0.28$ GeV and $m_Z = 91.172 \pm 0.031$ GeV along with $\sin^2\theta_W = 0.2322 \pm 0.0014$. They use $\sigma(W)/\sigma(Z) = 3.23 \pm 0.05$ and $\Gamma(Z) = 2.498 \pm 0.020$ GeV.

[18] ABE 92I report $1216 \pm 38^{+27}_{-31}$ $W \to \mu\nu$ and $106 \pm 10^{+0.2}_{-1}$ $Z \to \mu^+\mu^-$ events which are combined with 2426 $W \to e\nu$ events of ABE 91C to derive the ratio σ_W B($W \to \ell\nu$)/σ_Z B($Z \to \ell^+\ell^-$)= $10.0 \pm 0.6 \pm 0.4$. Finally the value of $\Gamma(Z)$ measured by LEP 92 is used to extract $\Gamma(W)$.

[19] ABE 90 extract $\Gamma(W) = 2.19 \pm 0.20$ by using the value $\Gamma(Z) = 2.57 \pm 0.07$ GeV. However, in ABE 91C they update their analysis with a new LEP value $\Gamma(Z) = 2.496 \pm 0.016$; the value $\Gamma(W) = 2.12 \pm 0.20$ above reflects this update. They measured $R = 10.2 \pm 0.8 \pm 0.4$, assumed $\sin^2\theta_W = 0.229 \pm 0.007$, and took predicted values $\sigma(W)/\sigma(Z) = 3.23 \pm 0.03$ and $\Gamma(W \to e\nu)/\Gamma(Z \to ee) = 2.70 \pm 0.02$. This yields $\Gamma(W)/\Gamma(Z) = 0.85 \pm 0.08$. The quoted error for $\Gamma(W)$ includes systematic uncertainties. $E_{\rm cm}^{p\overline{p}} = 1800$ GeV.

[20] ALITTI 90C used the same technique as described for ABE 90. They measured $R = 9.38^{+0.82}_{-0.72} \pm 0.25$, obtained $\Gamma(W)/\Gamma(Z) = 0.902 \pm 0.074 \pm 0.024$. Using $\Gamma(Z) = 2.546 \pm 0.032$ GeV, they obtained the $\Gamma(W)$ value quoted above and the limits $\Gamma(W) < 2.56$ (2.64) GeV at the 90% (95%) CL. $E_{\rm cm}^{p\overline{p}} = 546,630$ GeV.

[21] ALBAJAR 89 result is from a total sample of 299 $W \to e\nu$ events.

[22] If systematic error is neglected, result is $2.7^{+1.4}_{-1.5}$ GeV. This is enhanced subsample of 172 total events.

W ANOMALOUS MAGNETIC MOMENT ($\Delta\kappa$)

The full magnetic moment is given by $\mu_W = e(1+\kappa+\lambda)/2m_W$. In the Standard Model, at tree level, $\kappa = 1$ and $\lambda = 0$. Some papers have defined $\Delta\kappa = 1-\kappa$ and assume that $\lambda = 0$. Note that the electric quadrupole moment is given by $-e(\kappa-\lambda)/m_W^2$. A description of the parameterization of these moments and additional references can be found in HAGIWARA 87 and BAUR 88. The parameter Λ appearing in the theoretical limits below is a regularization cutoff which roughly corresponds to the energy scale where the structure of the W boson becomes manifest.

VALUE ($e/2m_W$)	*DOCUMENT ID*		*TECN*
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	[23] ABE	95G	
	[24] ALITTI	92C	UA2
	[25] SAMUEL	92	THEO
	[26] SAMUEL	91	THEO
	[27] GRIFOLS	88	THEO
	[28] GROTCH	87	THEO
	[29] VANDERBIJ	87	THEO
	[30] GRAU	85	THEO
	[31] SUZUKI	85	THEO
	[32] HERZOG	84	THEO

[23] ABE 95G report $-1.3 < \kappa < 3.2$ for λ=0 and $-0.7 < \lambda < 0.7$ for κ=1 in $p\overline{p} \to e\nu_e\gamma X$ and $\mu\nu_\mu\gamma X$ at $\sqrt{s} = 1800$ GeV.

[24] ALITTI 92C measure $\kappa = 1^{+2.6}_{-2.2}$ and $\lambda = 0^{+1.7}_{-1.8}$ in $p\overline{p} \to e\nu\gamma + X$ at $\sqrt{s} = 630$ GeV. At 95%CL they report $-3.5 < \kappa < 5.9$ and $-3.6 < \lambda < 3.5$.

[25] SAMUEL 92 use preliminary CDF and UA2 data and find $-2.4 < \kappa < 3.7$ at 96%CL and $-3.1 < \kappa < 4.2$ at 95%CL respectively. They use data for $W\gamma$ production and radiative W decay.

[26] SAMUEL 91 use preliminary CDF data for $p\overline{p} \to W\gamma X$ to obtain $-11.3 \leq \Delta\kappa \leq 10.9$. Note that their $\kappa = 1-\Delta\kappa$.

[27] GRIFOLS 88 uses deviation from ρ parameter to set limit $\Delta\kappa \lesssim 65\ (M_W^2/\Lambda^2)$.

[28] GROTCH 87 finds the limit $-37 < \Delta\kappa < 73.5$ (90% CL) from the experimental limits on $e^+e^- \to \nu\overline{\nu}\gamma$ assuming three neutrino generations and $-19.5 < \Delta\kappa < 56$ for four generations. Note their $\Delta\kappa$ has the opposite sign as our definition.

[29] VANDERBIJ 87 uses existing limits to the photon structure to obtain $|\Delta\kappa| < 33\ (m_W/\Lambda)$. In addition VANDERBIJ 87 discusses problems with using the ρ parameter of the Standard Model to determine $\Delta\kappa$.

[30] GRAU 85 uses the muon anomaly to derive a coupled limit on the anomalous magnetic dipole and electric quadrupole (λ) moments $1.05 > \Delta\kappa \ln(\Lambda/m_W) + \lambda/2 > -2.77$. In the Standard Model $\lambda = 0$.

[31] SUZUKI 85 uses partial-wave unitarity at high energies to obtain $|\Delta\kappa| \lesssim 190\ (m_W/\Lambda)^2$. From the anomalous magnetic moment of the muon, SUZUKI 85 obtains $|\Delta\kappa| \lesssim 2.2/\ln(\Lambda/m_W)$. Finally SUZUKI 85 uses deviations from the ρ parameter and obtains a very qualitative, order-of-magnitude limit $|\Delta\kappa| \lesssim 150\ (m_W/\Lambda)^4$ if $|\Delta\kappa| \ll 1$.

[32] HERZOG 84 consider the contribution of W-boson to muon magnetic moment including anomalous coupling of $WW\gamma$. Obtain a limit $-1 < \Delta\kappa < 3$ for $\Lambda \gtrsim 1$ TeV.

W^+ DECAY MODES

W^- modes are charge conjugates of the modes below.

	Mode		Fraction (Γ_i/Γ)	Confidence level
Γ_1	$\ell^+\nu$	[a]	(10.8±0.4) %	
Γ_2	$e^+\nu$		(10.8±0.4) %	
Γ_3	$\mu^+\nu$		(10.4±0.6) %	
Γ_4	$\tau^+\nu$		(10.9±1.0) %	
Γ_5	hadrons		(67.9±1.5) %	
Γ_6	$\pi^+\gamma$		$< 5 \times 10^{-4}$	95%

[a] ℓ indicates each type of lepton (e, μ, and τ), not sum over them.

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 8 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.7$ for 5 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\rm total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	44		
x_4	43	19	
x_5	−73	−65	−84
	x_2	x_3	x_4

W BRANCHING RATIOS

$\Gamma(\ell^+\nu)/\Gamma_{\rm total}$ Γ_1/Γ

ℓ indicates average over e, μ, and τ modes, not sum over modes.

Currently only e and μ data enter this average because there are no absolute τ data, only the τ/e ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.108±0.004 OUR AVERAGE		Includes data from the 2 datablocks that follow this one.		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.104±0.008	3642	[33] ABE	92I CDF	$E_{\rm cm}^{p\overline{p}}$= 1.8 TeV

[33] $1216 \pm 38^{+27}_{-31}$ $W \to \mu\nu$ events from ABE 92I and 2426 $W \to e\nu$ events of ABE 91C. ABE 92I give the inverse quantity as 9.6 ± 0.7 and we have inverted.

$\Gamma(e^+\nu)/\Gamma_{\rm total}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				
0.108 ±0.004 OUR FIT				
0.109 ±0.004 OUR AVERAGE				
0.1094±0.0033±0.0031		[34] ABE	95W CDF	$E_{\rm cm}^{p\overline{p}}$= 1800 GeV
0.10 ±0.014 $^{+0.02}_{-0.03}$	248	[35] ANSARI	87C UA2	$E_{\rm cm}^{p\overline{p}}$= 546,630 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.106 ±0.0096	2426	[36] ABE	91C	CDF	Repl. by ABE 94B
seen	299	[37] ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}}$ = 546,630 GeV
seen	119	APPEL	86	UA2	$E_{cm}^{p\overline{p}}$ = 546,630 GeV
seen	172	ARNISON	86	UA1	Repl. by ALBAJAR 89

[34] ABE 95W result is from a measurement of $\sigma B(W \to e\nu)/\sigma B(Z \to e^+e^-) = 10.90 \pm 0.32 \pm 0.29$, the theoretical prediction for the cross section ratio, the experimental knowledge of $\Gamma(Z \to e^+e^-) = 83.98 \pm 0.18$ MeV, and $\Gamma(Z) = 2.4969 \pm 0.0038$.

[35] The first error was obtained by adding the statistical and systematic experimental uncertainties in quadrature. The second error reflects the dependence on theoretical prediction of total W cross section: $\sigma(546\text{ GeV}) = 4.7^{+1.4}_{-0.7}$ nb and $\sigma(630\text{ GeV}) = 5.8^{+1.8}_{-1.0}$ nb. See ALTARELLI 85B.

[36] ABE 91C result is from a measurement of $\sigma B(W \to e\nu)/\sigma B(Z \to e^+e^-)$, the theoretical prediction for the cross section ratio, and the experimental knowledge of $\Gamma(Z \to e^+e^-)/\Gamma(Z \to \text{all})$.

[37] ALBAJAR 89 experiment determines values of branching ratio times production cross section.

$\Gamma(\mu^+\nu)/\Gamma_{total}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

The data in this block is included in the average printed for a previous datablock.

0.104±0.006 OR FIT

0.10 ±0.01	1216	[38] ABE	92I	CDF	$E_{cm}^{p\overline{p}}$ = 1.8 TeV

[38] ABE 92I quote the inverse quantity as 9.9 ± 1.2 which we have inverted.

$\Gamma(\tau^+\nu)/\Gamma_{total}$ Γ_4/Γ

VALUE	DOCUMENT ID
0.109±0.010 OR FIT	

$\Gamma(\text{hadrons})/\Gamma_{total}$ Γ_5/Γ

VALUE	DOCUMENT ID
0.679±0.015 OR FIT	

$\Gamma(\mu^+\nu)/\Gamma(e^+\nu)$ Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.95±0.05 OR FIT				
0.97±0.06 OR AVERAGE				
0.89±0.10	13k	[39] ABACHI	95D	D0	$E_{cm}^{p\overline{p}}$ = 1.8 TeV
1.02±0.08	1216	[40] ABE	92I	CDF	$E_{cm}^{p\overline{p}}$ = 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.00±0.14±0.08	67	ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}}$ = 546,630 GeV
$1.24^{+0.6}_{-0.4}$	14	ARNISON	84D	UA1	Repl. by ALBAJAR 89

[39] ABACHI 95D obtain this result from the measured $\sigma_W B(W \to \mu\nu) = 2.09 \pm 0.23 \pm 0.11$ nb and $\sigma_W B(W \to e\nu) = 2.36 \pm 0.07 \pm 0.13$ nb in which the first error is the combined statistical and systematic uncertainty, the second reflects the uncertainty in the luminosity.

[40] ABE 92I obtain $\sigma_W B(W \to \mu\nu) = 2.21 \pm 0.07 \pm 0.21$ and combine with ABE 91C $\sigma_W B((W \to e\nu))$ to give a ratio of the couplings from which we derive this measurement.

$\Gamma(\tau^+\nu)/\Gamma(e^+\nu)$ Γ_4/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ±0.08 OR FIT				
1.00 ±0.08 OR AVERAGE				
0.94 ±0.14	179	[41] ABE	92E	CDF	$E_{cm}^{p\overline{p}}$ = 1.8 TeV
1.04 ±0.08 ±0.08	754	[42] ALITTI	92F	UA2	$E_{cm}^{p\overline{p}}$ = 630 GeV
1.02 ±0.20 ±0.12	32	ALBAJAR	89	UA1	$E_{cm}^{p\overline{p}}$ = 546,630 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.995±0.112±0.083	198	ALITTI	91C	UA2	Repl. by ALITTI 92F
1.02 ±0.20 ±0.10	32	ALBAJAR	87	UA1	Repl. by ALBAJAR 89

[41] ABE 92E use two procedures for selecting $W \to \tau\nu_\tau$ events. The missing E_T trigger leads to 132 ± 14 ± 8 events and the τ trigger to 47 ± 9 ± 4 events. Proper statistical and systematic correlations are taken into account to arrive at $\sigma B(W \to \tau\nu) = 2.05 \pm 0.27$ nb. Combined with ABE 91C result on $\sigma B(W \to e\nu)$, ABE 92E quote a ratio of the couplings from which we derive this measurement.

[42] This measurement is derived by us from the ratio of the couplings of ALITTI 92F.

$\Gamma(\pi^+\gamma)/\Gamma(e^+\nu)$ Γ_6/Γ_2

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$< 7.5 \times 10^{-3}$	95	ABE		92K CDF	$E_{\rm cm}^{p\overline{p}} = 1.8$ TeV
$< 4.9 \times 10^{-3}$	95	[43] ALITTI		92D UA2	$E_{\rm cm}^{p\overline{p}} = 630$ GeV
$<58 \times 10^{-3}$	95	[44] ALBAJAR		90 UA1	$E_{\rm cm}^{p\overline{p}} = 546, 630$ GeV

[43] ALITTI 92D limit is 3.8×10^{-3} at 90%CL.
[44] ALBAJAR 90 obtain $<$ 0.048 at 90%CL.

W REFERENCES

ABACHI	95D	PRL 75 1456	+Abbott, Abolins, Acharya+	(D0 Collab.)
ABE	95C	PRL 74 341	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95G	PRL 74 1936	+Albrow, Amidei, Antos+	(CDF Collab.)
ABE	95P	PRL 75 11	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
Also	95Q	PR D52 4784	Abe, Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95W	PR D52 2624	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
Also	94B	PRL 73 220	Abe, Albrow, Amidei, Anway-Wiese+	(CDF Collab.)
ABE	94B	PRL 73 220	+Albrow, Amidei, Anway-Wiese+	(CDF Collab.)
ROSNER	94	PR D49 1363	+Worah, Takeuchi	(EFI, FNAL)
ABE	92E	PRL 68 3398	+Amidei, Apollinari, Atac+	(CDF Collab.)
ABE	92I	PRL 69 28	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	92K	PRL 69 2160	+Amidei, Anway-Weiss+	(CDF Collab.)
ALITTI	92	PL B276 365	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92B	PL B276 354	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92C	PL B277 194	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92D	PL B277 203	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92F	PL B280 137	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
LEP	92	PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collabs.)
SAMUEL	92	PL B280 124	+Li, Sinha, Sinha, Sundaresan	(OKSU, CARL)
ABE	91C	PR D44 29	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR	91	PL B253 503	+Albrow, Allkofer, Ankoviak, Apsimon+	(UA1 Collab.)
ALITTI	91C	ZPHY C52 209	+Ambrosini, Ansari, Autiero+	(UA2 Collab.)
SAMUEL	91	PRL 67 9	+Li, Sinha, Sinha, Sundaresan	(OKSU, CARL)
Also	91C	PRL 67 2920 erratum		
ABE	90	PRL 64 152	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
Also	91C	PR D44 29	Abe, Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	90G	PRL 65 2243	+Amidei, Apollinari, Atac+	(CDF Collab.)
Also	91B	PR D43 2070	Abe, Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR	90	PL B241 283	+Albrow, Allkofer+	(UA1 Collab.)
ALITTI	90B	PL B241 150	+Ansari, Ansorge, Autiero+	(UA2 Collab.)
ALITTI	90C	ZPHY C47 11	+Ansari, Ansorge, Bagnaia+	(UA2 Collab.)
ABE	89I	PRL 62 1005	+Amidei, Apollinari, Ascoli, Atac+	(CDF Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
BAUR	88	NP B308 127	+Zeppenfeld	(FSU, WISC)
GRIFOLS	88	IJMP A3 225	+Peris, Sola	(BARC, DESY)
Also	87	PL B197 437	Grifols, Peris, Sola	(BARC, DESY)
ALBAJAR	87	PL B185 233	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
ANSARI	87	PL B186 440	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
ANSARI	87C	PL B194 158	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
GROTCH	87	PR D36 2153	+Robinett	(PSU)
HAGIWARA	87	NP B282 253	+Peccei, Zeppenfeld, Hikasa	(KEK, UCLA, FSU)
VANDERBIJ	87	PR D35 1088	van der Bij	(FNAL)
APPEL	86	ZPHY C30 1	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
ARNISON	86	PL 166B 484	+Albrow, Allkofer, Astbury+	(UA1 Collab.) J
ALTARELLI	85B	ZPHY C27 617	+Ellis, Martinelli	(CERN, FNAL, FRAS)
GRAU	85	PL 154B 283	+Grifols	(BARC)
SUZUKI	85	PL 153B 289		(LBL)
ARNISON	84D	PL 134B 469	+Astbury, Aubert, Bacci+	(UA1 Collab.)
BAGNAIA	84	ZPHY C24 1	+Banner, Battiston, Blech+	(UA2 Collab.)
HERZOG	84	PL 148B 355		(WISC)
Also	84B	PL 155B 468 erratum	Herzog	(WISC)
ARNISON	83	PL 122B 103	+Astbury, Aubert, Bacci+	(UA1 Collab.)
ARNISON	83D	PL 129B 273	+Astbury, Aubert, Bacci+	(UA1 Collab.)
BAGNAIA	83	PL 129B 130	+Banner, Battiston, Bloch+	(UA2 Collab.)
BANNER	83B	PL 122B 476	+Battiston, Bloch, Bonaudi+	(UA2 Collab.)