$f_4(2300)$

$I^G(J^{PC}) = 0^+(4^{++})$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_0(2350)$. Contains results only from formation experiments. For production experiments see the $\overline{N}N(1100\text{–}3600)$ entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $\rho_5(2350)$.

$f_4(2300)$ MASS

$\overline{p}p \rightarrow \pi\pi$ or $\overline{K}K$

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 2314	HASAN	94	RVUE	$\overline{p}p \rightarrow \pi\pi$
~ 2300	[1] MARTIN	80B	RVUE	
~ 2300	[1] MARTIN	80C	RVUE	
~ 2340	[2] CARTER	78B	CNTR	0.7–2.4 $\overline{p}p \rightarrow K^-K^+$
~ 2330	DULUDE	78B	OSPK	1–2 $\overline{p}p \rightarrow \pi^0\pi^0$
~ 2310	[3] CARTER	77	CNTR	0.7–2.4 $\overline{p}p \rightarrow \pi\pi$

[1] $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\overline{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
[2] $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.
[3] $I(J^P) = 0(4^+)$ from amplitude analysis.

S-CHANNEL $\overline{p}p$ or $\overline{N}N$

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 2380	[4] CUTTS	78B	CNTR	0.97–3 $\overline{p}p \rightarrow \overline{N}N$
2345 ± 15	[4,5] COUPLAND	77	CNTR	0.7–2.4 $\overline{p}p \rightarrow \overline{p}p$
2359 ± 2	[4,6] ALSPECTOR	73	CNTR	$\overline{p}p$ S channel
2375 ± 10	ABRAMS	70	CNTR	S channel $\overline{N}N$

[4] Isospins 0 and 1 not separated.
[5] From a fit to the total elastic cross section.
[6] Referred to as U or U region by ALSPECTOR 73.

$f_4(2300)$ WIDTH

$\overline{p}p \rightarrow \pi\pi$ or $\overline{K}K$

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 278	HASAN	94	RVUE	$\overline{p}p \rightarrow \pi\pi$
~ 200	[7] MARTIN	80C	RVUE	
~ 150	[8] CARTER	78B	CNTR	0.7–2.4 $\overline{p}p \rightarrow K^-K^+$
~ 210	[9] CARTER	77	CNTR	0.7–2.4 $\overline{p}p \rightarrow \pi\pi$

[7] $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\overline{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
[8] $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.
[9] $I(J^P) = 0(4^+)$ from amplitude analysis.

S-CHANNEL $\overline{p}p$ or $\overline{N}N$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
• • • We do not use the following data for averages, fits, limits, etc. • • •				
135^{+150}_{-65}	[10,11] COUPLAND	77	CNTR	0.7–2.4 $\overline{p}p \rightarrow \overline{p}p$
165^{+18}_{-8}	[11] ALSPECTOR	73	CNTR	$\overline{p}p$ S channel
~ 190	ABRAMS	70	CNTR	S channel $\overline{N}N$

[10] From a fit to the total elastic cross section.
[11] Isospins 0 and 1 not separated.

$f_4(2300)$ REFERENCES

HASAN	94	PL B334 215	+Bugg	(LOQM)
MARTIN	80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	+Pennington	(DURH) JP
CARTER	78B	NP B141 467		(LOQM)
CUTTS	78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
DULUDE	78B	PL 79B 335	+Lanou, Massimo, Peaslee+	(BROW, MIT, BARI) JP
CARTER	77	PL 67B 117	+Coupland, Eisenhandler, Astbury+	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)

OTHER RELATED PAPERS

FIELDS	71	PRL 27 1749	+Cooper, Rhines, Allison	(ANL, OXF)
YOH	71	PRL 26 922	+Barish, Caroll, Lobkowicz+	(CIT, BNL, ROCH)
BRICMAN	69	PL 29B 451	+Ferro-Luzzi, Bizard+	(CERN, CAEN, SACL)