$f_2(2010)$

$I^G(J^{PC}) = 0^+(2^{++})$

See also the mini-review under non-$q\overline{q}$ candidates. (See the index for the page number.)

$f_2(2010)$ MASS

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
2011^{+62}_{-76}	[1] ETKIN	88	MPS	22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1980 ± 20	[2] BOLONKIN	88	SPEC	40 $\pi^- p \rightarrow K^0_S K^0_S n$
2050^{+90}_{-50}	ETKIN	85	MPS	22 $\pi^- p \rightarrow 2\phi n$
2120^{+20}_{-120}	LINDENBAUM	84	RVUE	
2160 ± 50	ETKIN	82	MPS	22 $\pi^- p \rightarrow 2\phi n$

[1] Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi$ 2^{++} S_2, D_2, and D_0 is 98^{+1}_{-3}, 0^{+1}_{-0}, and 2^{+2}_{-1}, respectively.

[2] Statistically very weak, only 1.4 s.d.

$f_2(2010)$ WIDTH

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
202^{+67}_{-62}	[3] ETKIN	88	MPS	22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
145 ± 50	[4] BOLONKIN	88	SPEC	40 $\pi^- p \rightarrow K^0_S K^0_S n$
200^{+160}_{-50}	ETKIN	85	MPS	22 $\pi^- p \rightarrow 2\phi n$
300^{+150}_{-50}	LINDENBAUM	84	RVUE	
310 ± 70	ETKIN	82	MPS	22 $\pi^- p \rightarrow 2\phi n$

[3] Includes data of ETKIN 85.

[4] Statistically very weak, only 1.4 s.d.

$f_2(2010)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)
Γ_1	$\phi\phi$	seen

$f_2(2010)$ REFERENCES

BOLONKIN	88	NP B309 426	+Bloshenko, Gorin+	(ITEP, SERP)
ETKIN	88	PL B201 568	+Foley, Lindenbaum+	(BNL, CUNY)
ETKIN	85	PL 165B 217	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285		(CUNY)
ETKIN	82	PRL 49 1620	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
Also	83	Brighton Conf. 351	Lindenbaum	(BNL, CUNY)

OTHER RELATED PAPERS

LANDBERG	96	PR D53 2839	+Adams, Chan+ (BNL, CUNY, RPI)
ARMSTRONG	89B	PL B221 221	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)
GREEN	86	PRL 56 1639	+Lai+ (FNAL, ARIZ, FSU, NDAM, TUFTS, VAND+)
BOOTH	84	NP B242 51	+Ballance, Carroll, Donald+ (LIVP, GLAS, CERN)