

**$\Delta(1700) D_{33}$**  $I(J^P) = \frac{3}{2}(\frac{3}{2}^-)$  Status: \*\*\*

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 **$\Delta(1700)$  BREIT-WIGNER MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1670 to 1770 (<math>\approx 1700</math>) OUR ESTIMATE</b>			
1762 $\pm 44$	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
1710 $\pm 30$	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1680 $\pm 70$	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1690 $\pm 15$	ARNDT 96	IPWA	$\gamma N \rightarrow \pi N$
1680	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$
1655	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1650	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
1718.4 $^{+13.1}_{-13.0}$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1622	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1629	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1600	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1680	<sup>3</sup> LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 **$\Delta(1700)$  BREIT-WIGNER WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>200 to 400 (<math>\approx 300</math>) OUR ESTIMATE</b>			
600 $\pm 250$	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
280 $\pm 80$	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
230 $\pm 80$	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
285 $\pm 20$	ARNDT 96	IPWA	$\gamma N \rightarrow \pi N$
272	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$
348	LI 93	IPWA	$\gamma N \rightarrow \pi N$
160	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
193.3 $\pm 26.0$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
209	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
216	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
200	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
240	<sup>3</sup> LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

**$\Delta(1700)$  POLE POSITION****REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1620 to 1700 (<math>\approx</math> 1660) OUR ESTIMATE</b>			
1655	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1651	<sup>4</sup> HOEHLER	93	SPED $\pi N \rightarrow \pi N$
$1675 \pm 25$	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1646	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1681 or 1672	<sup>5</sup> LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1600 or 1594	<sup>2</sup> LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

**-2×IMAGINARY PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>150 to 250 (<math>\approx</math> 200) OUR ESTIMATE</b>			
242	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
159	<sup>4</sup> HOEHLER	93	SPED $\pi N \rightarrow \pi N$
$220 \pm 40$	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
208	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
245 or 241	<sup>5</sup> LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
208 or 201	<sup>2</sup> LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

 **$\Delta(1700)$  ELASTIC POLE RESIDUE****MODULUS  $|r|$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
16	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
10	HOEHLER	93	SPED $\pi N \rightarrow \pi N$
$13 \pm 3$	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
13	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

**PHASE  $\theta$** 

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-12	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
$-20 \pm 25$	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-22	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

**$\Delta(1700)$  DECAY MODES**

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 N\pi$	10–20 %
$\Gamma_2 \Sigma K$	
$\Gamma_3 N\pi\pi$	80–90 %
$\Gamma_4 \Delta\pi$	30–60 %
$\Gamma_5 \Delta(1232)\pi$ , <i>S</i> -wave	25–50 %
$\Gamma_6 \Delta(1232)\pi$ , <i>D</i> -wave	1–7 %
$\Gamma_7 N\rho$	30–55 %
$\Gamma_8 N\rho$ , <i>S</i> =1/2, <i>D</i> -wave	
$\Gamma_9 N\rho$ , <i>S</i> =3/2, <i>S</i> -wave	5–20 %
$\Gamma_{10} N\rho$ , <i>S</i> =3/2, <i>D</i> -wave	
$\Gamma_{11} N\gamma$	0.12–0.26 %
$\Gamma_{12} N\gamma$ , helicity=1/2	0.08–0.16 %
$\Gamma_{13} N\gamma$ , helicity=3/2	0.025–0.12 %

 **$\Delta(1700)$  BRANCHING RATIOS**

$\Gamma(N\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.10 to 0.20 OUR ESTIMATE</b>			
0.14±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.12±0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
0.20±0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.16	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$
0.16	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow \Sigma K$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.002	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$
0.001 to 0.011	<sup>6</sup> DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from  $\pi N \rightarrow N\pi\pi$  analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the  $\Delta(1620)$   $S_{31}$  coupling to  $\Delta(1232)\pi$ .

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow \Delta(1232)\pi$ , <i>S</i> -wave	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>+0.21 to +0.29 OUR ESTIMATE</b>			
+0.32±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.18±0.04	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
+0.30	<sup>2,7</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.24	<sup>3</sup> LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow \Delta(1232)\pi, D\text{-wave}$	$(\Gamma_1 \Gamma_6)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>+0.05 to +0.11 OUR ESTIMATE</b>			
+0.08 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
0.14 ± 0.04	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
+0.05	<sup>2,7</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.10	<sup>3</sup> LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow N\rho, S=1/2, D\text{-wave}$	$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.17 ± 0.05	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow N\rho, S=3/2, S\text{-wave}$	$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>±0.11 to ±0.19 OUR ESTIMATE</b>			
+0.10 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
+0.04	<sup>2,7</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.30	<sup>3</sup> LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow N\rho, S=3/2, D\text{-wave}$	$(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.07	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$

### $\Delta(1700)$ PHOTON DECAY AMPLITUDES

#### $\Delta(1700) \rightarrow N\gamma$ , helicity-1/2 amplitude $A_{1/2}$

VALUE ( $\text{GeV}^{-1/2}$ )	DOCUMENT ID	TECN	COMMENT
<b>+0.104 ± 0.015 OUR ESTIMATE</b>			
0.090 ± 0.025	ARNDT 96	IPWA	$\gamma N \rightarrow \pi N$
0.111 ± 0.017	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.089 ± 0.033	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.112 ± 0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.130 ± 0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.123 ± 0.022	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.121 ± 0.004	LI 93	IPWA	$\gamma N \rightarrow \pi N$
+0.130 ± 0.037	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
+0.072 ± 0.033	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

#### $\Delta(1700) \rightarrow N\gamma$ , helicity-3/2 amplitude $A_{3/2}$

VALUE ( $\text{GeV}^{-1/2}$ )	DOCUMENT ID	TECN	COMMENT
<b>+0.085 ± 0.022 OUR ESTIMATE</b>			
0.097 ± 0.020	ARNDT 96	IPWA	$\gamma N \rightarrow \pi N$
0.107 ± 0.015	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.060 ± 0.015	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.047 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.050 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.102 ± 0.015	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.115 \pm 0.004$	LI	93	IPWA	$\gamma N \rightarrow \pi N$
$+0.098 \pm 0.036$	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
$+0.087 \pm 0.023$	FELLER	76	DPWA	$\gamma N \rightarrow \pi N$

## **Δ(1700) FOOTNOTES**

<sup>1</sup> Problems with CHEW 80 are discussed in section 2.1.11 of HOEHLER 83.

<sup>2</sup> LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to  $\pi N \rightarrow N\pi\pi$  data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

<sup>3</sup> From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

<sup>4</sup> See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of  $N$  and  $\Delta$  resonances as determined from Argand diagrams of  $\pi N$  elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

<sup>5</sup> LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to  $\pi N \rightarrow N\pi\pi$  data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

<sup>6</sup> The range given is from the four best solutions. DEANS 75 disagrees with  $\pi^+ p \rightarrow \Sigma^+ K^+$  data of WINNIK 77 around 1920 MeV.

<sup>7</sup> LONGACRE 77 considers this coupling to be well determined.

## **Δ(1700) REFERENCES**

For early references, see Physics Letters **111B** 70 (1982).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
HOEHLER	93	$\pi N$ Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
HOEHLER	83	Landolt-Bornstein 1/9B2		(KARLT)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
BARNHAM	80	NP B168 243	+Glickman, Mier-Jedrzejowicz+	(LOIC)
CHEW	80	Toronto Conf. 123		(LBL) IJP
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu	(SACL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadet	(SACL) IJP
WINNIK	77	NP B128 66	+Toaff, Revel, Goldberg, Berny	(HAIF) I
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP