

$\Lambda(1520) D_{03}$ $I(J^P) = 0(\frac{3}{2}^-)$ Status: ****

Discovered by FERRO-LUZZI 62; the elaboration in WATSON 63 is the classic paper on the Breit-Wigner analysis of a multichannel resonance.

The measurements of the mass, width, and elasticity published before 1975 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** (1982).

Production and formation experiments agree quite well, so they are listed together here.

 $\Lambda(1520)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1519.5 ±1.0 OUR ESTIMATE				
1519.50±0.18 OUR AVERAGE				
1517.3 ±1.5	300	BARBER	80D SPEC	$\gamma p \rightarrow \Lambda(1520) K^+$
1519 ±1		GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
1517.8 ±1.2	5k	BARLAG	79 HBC	$K^- p$ 4.2 GeV/c
1520.0 ±0.5		ALSTON...	78 DPWA	$\bar{K}N \rightarrow \bar{K}N$
1519.7 ±0.3	4k	CAMERON	77 HBC	$K^- p$ 0.96–1.36 GeV/c
1519 ±1		GOPAL	77 DPWA	$\bar{K}N$ multichannel
1519.4 ±0.3	2000	CORDEN	75 DBC	$K^- d$ 1.4–1.8 GeV/c

 $\Lambda(1520)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
15.6 ±1.0 OUR ESTIMATE				
15.59±0.27 OUR AVERAGE				
16.3 ±3.3	300	BARBER	80D SPEC	$\gamma p \rightarrow \Lambda(1520) K^+$
16 ±1		GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
14 ±3	677	¹ BARLAG	79 HBC	$K^- p$ 4.2 GeV/c
15.4 ±0.5		ALSTON...	78 DPWA	$\bar{K}N \rightarrow \bar{K}N$
16.3 ±0.5	4k	CAMERON	77 HBC	$K^- p$ 0.96–1.36 GeV/c
15.0 ±0.5		GOPAL	77 DPWA	$\bar{K}N$ multichannel
15.5 ±1.6	2000	CORDEN	75 DBC	$K^- d$ 1.4–1.8 GeV/c

 $\Lambda(1520)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\bar{K}$	45 ± 1%
$\Gamma_2 \Sigma\pi$	42 ± 1%
$\Gamma_3 \Lambda\pi\pi$	10 ± 1%
$\Gamma_4 \Sigma(1385)\pi$	

Γ_5	$\Sigma(1385)\pi(\rightarrow \Lambda\pi\pi)$
Γ_6	$\Lambda(\pi\pi)_{S\text{-wave}}$
Γ_7	$\Sigma\pi\pi$
Γ_8	$\Lambda\gamma$
Γ_9	$\Sigma^0\gamma$

CONSTRAINED FIT INFORMATION

An overall fit to 9 branching ratios uses 24 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 16.5$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-63				
x_3	-32 -33				
x_7	-4	-3	-1		
x_8	-9	-8	-4	0	
x_9	-24	-21	-10	-1	-2
	x_1	x_2	x_3	x_7	x_8

$\Lambda(1520)$ BRANCHING RATIOS


See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.01 OUR ESTIMATE			
0.448 ± 0.007 OUR FIT	Error includes scale factor of 1.2.		
0.455 ± 0.011 OUR AVERAGE			
0.47 ± 0.02	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.45 ± 0.03	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
0.448 ± 0.014	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.47 ± 0.01	GOPAL	77	DPWA See GOPAL 80
0.42	MAST	76	HBC $K^- p \rightarrow \bar{K}^0 n$

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ± 0.01 OUR ESTIMATE			
0.421 ± 0.007 OUR FIT	Error includes scale factor of 1.2.		
0.423 ± 0.011 OUR AVERAGE			
0.426 ± 0.014	CORDEN	75	DBC $K^- d$ 1.4–1.8 GeV/c
0.418 ± 0.017	BARBARO-...	69B	HBC $K^- p$ 0.28–0.45 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.46	KIM	71	DPWA K-matrix analysis

$\Gamma(\Sigma\pi)/\Gamma(N\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.940±0.026 OUR FIT	Error includes scale factor of 1.3.			
0.95 ±0.04 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.			
0.98 ±0.03	² GOPAL	77 DPWA	$\bar{K}N$ multichannel	
0.82 ±0.08	BURKHARDT	69 HBC	$K^- p$ 0.8–1.2 GeV/c	
1.06 ±0.14	SCHEUER	68 DBC	$K^- N$ 3 GeV/c	
0.96 ±0.20	DAHL	67 HBC	$\pi^- p$ 1.6–4 GeV/c	
0.73 ±0.11	DAUBER	67 HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.06 ±0.12	BERTHON	74 HBC	Quasi-2-body σ	
1.72 ±0.78	MUSGRAVE	65 HBC		

 $\Gamma(\Sigma\pi)/\Gamma(N\bar{K})$ $\Gamma(\Lambda\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.10 ±0.01 OUR ESTIMATE				
0.095±0.005 OUR FIT	Error includes scale factor of 1.2.			
0.096±0.008 OUR AVERAGE	Error includes scale factor of 1.6.			
0.091±0.006	CORDEN	75 DBC	$K^- d$ 1.4–1.8 GeV/c	
0.11 ±0.01	³ MAST	73B IPWA	$K^- p \rightarrow \Lambda\pi\pi$	

$\Gamma(\Lambda\pi\pi)/\Gamma(N\bar{K})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_1
0.213±0.012 OUR FIT	Error includes scale factor of 1.2.			
0.202±0.021 OUR AVERAGE				
0.22 ± 0.03	BURKHARDT 69	HBC	$K^- p$ 0.8–1.2 GeV/c	
0.19 ± 0.04	SCHEUER 68	DBC	$K^- N$ 3 GeV/c	
0.17 ± 0.05	DAHL 67	HBC	$\pi^- p$ 1.6–4 GeV/c	
0.21 ± 0.18	DAUBER 67	HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.27 ± 0.13	BERTHON 74	HBC	Quasi-2-body σ	
0.2	KIM 71	DPWA	K-matrix analysis	

 $\Gamma(\Sigma\pi)/\Gamma(\Lambda\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_3
4.42±0.25 OUR FIT	Error includes scale factor of 1.2.			
3.9 ± 0.6 OUR AVERAGE				
3.9 ± 1.0	UHLIG 67	HBC	$K^- p$ 0.9–1.0 GeV/c	
3.3 ± 1.1	BIRMINGHAM 66	HBC	$K^- p$ 3.5 GeV/c	
4.5 ± 1.0	ARMENTEROS65C	HBC		

 $\Gamma(\Sigma(1385)\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ
0.041±0.005	CHAN 72	HBC	$K^- p \rightarrow \Lambda\pi\pi$	

 $\Gamma(\Sigma(1385)\pi(\rightarrow \Lambda\pi\pi))/\Gamma(\Lambda\pi\pi)$

The $\Lambda\pi\pi$ mode is largely due to $\Sigma(1385)\pi$. Only the values of $(\Sigma(1385)\pi) / (\Lambda\pi\pi)$ given by MAST 73B and CORDEN 75 are based on real 3-body partial-wave analyses. The discrepancy between the two results is essentially due to the different hypotheses made concerning the shape of the $(\pi\pi)_S$ -wave state.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ_3
0.58±0.22	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.82±0.10	⁴ MAST 73B	IPWA	$K^- p \rightarrow \Lambda\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.39±0.10	⁵ BURKHARDT 71	HBC	$K^- p \rightarrow (\Lambda\pi\pi)\pi$	

 $\Gamma(\Lambda(\pi\pi)_S\text{-wave})/\Gamma(\Lambda\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ_3
0.20±0.08	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	

 $\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ
0.009 ± 0.001 OUR ESTIMATE				
0.0086±0.0005 OUR FIT				
0.0086±0.0005 OUR AVERAGE				
0.007 ± 0.002	⁶ CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.0085±0.0006	⁷ MAST 73	MPWA	$K^- p \rightarrow \Sigma\pi\pi$	
0.010 ± 0.0015	BARBARO-... 69B	HBC	$K^- p$ 0.28–0.45 GeV/c	

$\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$				Γ_8/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.008 ± 0.002 OUR ESTIMATE				
0.0079 ± 0.0014 OUR FIT				
0.0080 ± 0.0014	238	MAST	68B HBC	Using $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.45$
$\Gamma(\Sigma^0\gamma)/\Gamma_{\text{total}}$				Γ_9/Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0195 ± 0.0034 OUR FIT				
0.02 ± 0.0035	8	MAST	68B HBC	Not measured; see note

A(1520) FOOTNOTES

- 1 From the best-resolution sample of $\Lambda\pi\pi$ events only.
 2 The $\bar{K}N \rightarrow \Sigma\pi$ amplitude at resonance is $+0.46 \pm 0.01$.
 3 Assumes $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.46 \pm 0.02$.
 4 Both $\Sigma(1385)\pi$ Ds_{03} and $\Sigma(\pi\pi)$ DP_{03} contribute.
 5 The central bin (1514–1524 MeV) gives 0.74 ± 0.10 ; other bins are lower by 2-to-5 standard deviations.
 6 Much of the $\Sigma\pi\pi$ decay proceeds via $\Sigma(1385)\pi$.
 7 Assumes $\Gamma(N\bar{K})/\Gamma_{\text{total}} = 0.46$.
 8 Calculated from $\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$, assuming SU(3). Needed to constrain the sum of all the branching ratios to be unity.

A(1520) REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
BARBER	80D	ZPHY C7 17	+Dainton, Lee, Marshall+	(DARE, LANC, SHEF) (RHEL) IJP
GOPAL	80	Toronto Conf. 159	+Blokzijl, Jongejans+	(AMST, CERN, NIJM, OXF)
BARLAG	79	NP B149 220	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
ALSTON...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	+Franek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
GOPAL	77	NP B119 362	+Alston-Garnjost, Bangerter+	(LBL)
MAST	76	PR D14 13	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM)
CORDEN	75	NP B84 306	+Tristram+	(CDEF, RHEL, SACL, STRB)
BERTHON	74	NC 21A 146	+Bangerter, Alston-Garnjost+	(LBL) IJP
MAST	73	PR D7 3212	+Bangerter, Alston-Garnjost+	(LBL) IJP
MAST	73B	PR D7 5	+Button-Shafer, Hertzbach, Kofler+	(MASA, YALE)
CHAN	72	PRL 28 256	+Filthuth, Kluge+	(HEID, CERN, SACL)
BURKHARDT	71	NP B27 64		(HARV) IJP
KIM	71	PRL 27 356	Kim	(HARV) IJP
Also	70	Duke Conf. 161	Barbaro-Galtieri, Bangerter, Mast, Tripp	(LRL)
BARBARO...	69B	Lund Conf. 352	Tripp	(LRL)
Also	70	Duke Conf. 95	+Filthuth, Kluge+	(HEID, EFI, CERN, SACL)
BURKHARDT	69	NP B14 106	+Alston-Garnjost, Bangerter, Galtieri+	(LRL)
MAST	68B	PRL 21 1715	+Merrill, Verglas, DeWitt+	(SABRE Collab.)
SCHEUER	68	NP B8 503	+Hardy, Hess, Kirz, Miller	(LRL)
DAHL	67	PR 163 1377	+Malamud, Schlein, Slater, Stork	(UCLA)
DAUBER	67	PL 24B 525	+Charlton, Condon, Glasser, Yodh+	(UMD, NRL)
UHLIG	67	PR 155 1448		(BIRM, GLAS, LOIC, OXF, RHEL)
BIRMINGHAM	66	PR 152 1148	+Ferro-Luzzi+	(CERN, HEID, SACL)
ARMENTEROS	65C	PL 19 338	+Petmezas+	(BIRM, CERN, EPOL, LOIC, SACL)
MUSGRAVE	65	NC 35 735	+Ferro-Luzzi, Tripp	(LRL) IJP
WATSON	63	PR 131 2248	+Tripp, Watson	(LRL) IJP
FERRO-LUZZI	62	PRL 8 28		