

$\rho_3(1690)$ $I^G(J^{PC}) = 1^+(3^{--})$ **$\rho_3(1690)$ MASS**

VALUE (MeV)	DOCUMENT ID
1691 ± 5 OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values.
1688.8 ± 2.1 OUR AVERAGE	Includes data from the 5 datablocks that follow this one.

 2π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

1686 ± 4 OUR AVERAGE

1677 ± 14		EVANGELISTA 81	OMEG	-	$12 \pi^- p \rightarrow 2\pi p$
1679 ± 11	476	BALTAY	78B HBC	0	$15 \pi^+ p \rightarrow \pi^+ \pi^- n$
1678 ± 12	175	¹ ANTIPOV	77 CIBS	0	$25 \pi^- p \rightarrow p 3\pi$
1690 ± 7	600	¹ ENGLER	74 DBC	0	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
1693 ± 8		² GRAYER	74 ASPK	0	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
1678 ± 12		MATTHEWS	71C DBC	0	$7 \pi^+ N$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1734 ± 10		³ CORDEN	79 OMEG		$12-15 \pi^- p \rightarrow n 2\pi$
1692 ± 12		^{2,4} ESTABROOKS	75 RVUE		$17 \pi^- p \rightarrow \pi^+ \pi^- n$
1737 ± 23		ARMENISE	70 DBC	0	$9 \pi^+ N$
1650 ± 35	122	BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N 2\pi$
1687 ± 21		STUNTEBECK	70 HDCC	0	$8 \pi^- p, 5.4 \pi^+ d$
1683 ± 13		ARMENISE	68 DBC	0	$5.1 \pi^+ d$
1670 ± 30		GOLDBERG	65 HBC	0	$6 \pi^+ d, 8 \pi^- p$

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.² Uses same data as HYAMS 75.³ From a phase shift solution containing a $f'_2(1525)$ width two times larger than the $K\bar{K}$ result.⁴ From phase-shift analysis. Error takes account of spread of different phase-shift solutions. **$K\bar{K}$ AND $K\bar{K}\pi$ MODES**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

1696 ± 4 OUR AVERAGE

1699 ± 5		ALPER	80 CNTR	0	$62 \pi^- p \rightarrow K^+ K^- n$
1698 ± 12	6k	^{5,6} MARTIN	78D SPEC		$10 \pi p \rightarrow K_S^0 K^- p$
1692 ± 6		BLUM	75 ASPK	0	$18.4 \pi^- p \rightarrow n K^+ K^-$
1690 ± 16		ADERHOLZ	69 HBC	+	$8 \pi^+ p \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1694 ± 8	⁷ COSTA...	80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
--------------	-----------------------	----	------	------------------------------------

⁵ From a fit to $J^P = 3^-$ partial wave.

⁶ Systematic error on mass scale subtracted.

⁷ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

(4π) $^\pm$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

1686 \pm 5 OUR AVERAGE Error includes scale factor of 1.1.

1694 ± 6		⁸ EVANGELISTA 81	OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1665 ± 15	177	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$
1670 ± 10		THOMPSON	74 HBC	+	$13 \pi^+ p$
1687 ± 20		CASON	73 HBC	-	$8,18.5 \pi^- p$
1685 ± 14		⁹ CASON	73 HBC	-	$8,18.5 \pi^- p$
1680 ± 40	144	BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N4\pi$
1689 ± 20	102	⁹ BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N2\rho$
1705 ± 21		CASO	70 HBC	-	$11.2 \pi^- p \rightarrow n\rho2\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1718 ± 10		¹⁰ EVANGELISTA 81	OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1673 ± 9		¹¹ EVANGELISTA 81	OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1733 ± 9	66	⁹ KLIGER	74 HBC	-	$4.5 \pi^- p \rightarrow p4\pi$
1630 ± 15		HOLMES	72 HBC	+	$10-12 K^+ p$
1720 ± 15		BALTAY	68 HBC	+	$7, 8.5 \pi^+ p$

⁸ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

⁹ From $\rho^\pm \rho^0$ mode.

¹⁰ From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

¹¹ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

$\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1681 \pm 7 OUR AVERAGE

1670 ± 25	¹² ALDE	95 GAM2		$38 \pi^- p \rightarrow \omega\pi^0 n$
1690 ± 15		EVANGELISTA 81	OMEG	-
1666 ± 14		GESSAROLI	77 HBC	
1686 ± 9		THOMPSON	74 HBC	+
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1654 ± 24		BARNHAM	70 HBC	+

¹² Supersedes ALDE 92C.

$\eta\pi^+\pi^-$ MODE(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1680±15FUKUI 88 SPEC 0 $8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$


• • • We do not use the following data for averages, fits, limits, etc. • • •

1700±47 13 ANDERSON 69 MMS — $16 \pi^- p$ backward1632±15 13,14 FOCACCI 66 MMS — $7-12 \pi^- p \rightarrow p$ MM1700±15 13,14 FOCACCI 66 MMS — $7-12 \pi^- p \rightarrow p$ MM1748±15 13,14 FOCACCI 66 MMS — $7-12 \pi^- p \rightarrow p$ MM13 Seen in 2.5–3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+\pi^-$ pairs in ρ band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1976)

14 Not seen by BOWEN 72.

 $\rho_3(1690)$ WIDTH **2π , $K\bar{K}$, AND $K\bar{K}\pi$ MODES**

VALUE (MeV)	DOCUMENT ID
160±10 OUR AVERAGE	Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.

 $\rho_3(1690)$ width, 2π , $K\bar{K}$, and $K\bar{K}\pi$ modes (MeV)

2π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

186±14 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.


220±29	DENNEY	83	LASS	10	$\pi^+ N$
246±37	EVANGELISTA	81	OMEG	—	$12 \pi^- p \rightarrow 2\pi p$
116±30	BALTAY	78B	HBC	0	$15 \pi^+ p \rightarrow \pi^+ \pi^- n$
162±50	15 ANTIPOV	77	CIBS	0	$25 \pi^- p \rightarrow p 3\pi$
167±40	ENGLER	74	DBC	0	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
200±18	16 GRAYER	74	ASPK	0	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
156±36	MATTHEWS	71C	DBC	0	$7 \pi^+ N$
171±65	ARMENISE	70	DBC	0	$9 \pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
322±35	17 CORDEN	79	OMEG	12–15	$\pi^- p \rightarrow n 2\pi$
240±30	16,18 ESTABROOKS	75	RVUE	17	$\pi^- p \rightarrow \pi^+ \pi^- n$
180±30	122 BARTSCH	70B	HBC	+	$8 \pi^+ p \rightarrow N 2\pi$
267 ⁺⁷² ₋₄₆	STUNTEBECK	70	HDBC	0	$8 \pi^- p, 5.4 \pi^+ d$
188±49	ARMENISE	68	DBC	0	$5.1 \pi^+ d$
180±40	GOLDBERG	65	HBC	0	$6 \pi^+ d, 8 \pi^- p$

¹⁵ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹⁶ Uses same data as HYAMS 75 and BECKER 79.

¹⁷ From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$ result.

¹⁸ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.



$\rho_3(1690)$ width, 2π mode (MeV)

$K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

204±18 OUR AVERAGE

199±40	6000	19 MARTIN	78D SPEC	10 $\pi^- p \rightarrow K_S^0 K^- p$
205±20		BLUM	75 ASPK 0	18.4 $\pi^- p \rightarrow n K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
219± 4		ALPER	80 CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
186±11		20 COSTA...	80 OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
112±60		ADERHOLZ	69 HBC +	8 $\pi^+ p \rightarrow K\bar{K}\pi$

¹⁹ From a fit to $J^P = 3^-$ partial wave.

²⁰ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

$(4\pi)^{\pm}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

129±10 OUR AVERAGE

123±13		21 EVANGELISTA 81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
105±30	177	BALTAY	78B HBC	+	15 $\pi^+ p \rightarrow p 4\pi$
169 ⁺⁷⁰ ₋₄₈		CASON	73 HBC	—	8, 18.5 $\pi^- p$
135±30	144	BARTSCH	70B HBC	+	8 $\pi^+ p \rightarrow N 4\pi$
160±30	102	BARTSCH	70B HBC	+	8 $\pi^+ p \rightarrow N 2\rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •

230±28	22	EVANGELISTA 81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
184±33	23	EVANGELISTA 81	OMEG	—	12 $\pi^- p \rightarrow p 4\pi$
150	66	24 KLIGER	74	HBC	— 4.5 $\pi^- p \rightarrow p 4\pi$
106±25		THOMPSON	74	HBC	+ 13 $\pi^+ p$
125 ⁺⁸³ -35	24	CASON	73	HBC	— 8, 18.5 $\pi^- p$
130±30		HOLMES	72	HBC	+ 10–12 $K^+ p$
180±30	90	24 BARTSCH	70B	HBC	+ 8 $\pi^+ p \rightarrow N a_2 \pi$
100±35		BALTAY	68	HBC	+ 7, 8.5 $\pi^+ p$

21 From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

22 From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

23 From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

24 From $\rho^\pm \rho^0$ mode.

$\omega\pi$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	------------	----------------

The data in this block is included in the average printed for a previous datablock.

190±40 OUR AVERAGE

230±65	25	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$
190±65		EVANGELISTA 81	OMEG	—	12 $\pi^- p \rightarrow \omega \pi p$
160±56		GESSAROLI	77	HBC	11 $\pi^- p \rightarrow \omega \pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
89±25		THOMPSON	74	HBC	+ 13 $\pi^+ p$
130 ⁺⁷³ -43		BARNHAM	70	HBC	+ 10 $K^+ p \rightarrow \omega \pi X$

25 Supersedes ALDE 92C.

$\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	------------	----------------

The data in this block is included in the average printed for a previous datablock.

106±27

FUKUI	88	SPEC	0	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
-------	----	------	---	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

195	26	ANDERSON	69	MMS	— 16 $\pi^- p$ backward
< 21	26,27	FOCACCI	66	MMS	— 7–12 $\pi^- p \rightarrow p MM$
< 30	26,27	FOCACCI	66	MMS	— 7–12 $\pi^- p \rightarrow p MM$
< 38	26,27	FOCACCI	66	MMS	— 7–12 $\pi^- p \rightarrow p MM$

26 Seen in 2.5–3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+ \pi^-$ pairs in ρ^0 band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1979)

27 Not seen by BOWEN 72.

$\rho_3(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 4π	(71.1 \pm 1.9) %	
Γ_2 $\pi^\pm \pi^+ \pi^- \pi^0$	(67 \pm 22) %	
Γ_3 $\omega \pi$	(16 \pm 6) %	
Γ_4 $\pi \pi$	(23.6 \pm 1.3) %	
Γ_5 $K \bar{K} \pi$	(3.8 \pm 1.2) %	
Γ_6 $K \bar{K}$	(1.58 \pm 0.26) %	1.2
Γ_7 $\eta \pi^+ \pi^-$	seen	
Γ_8 $\pi \pi \rho$ Excluding 2ρ and $a_2(1320)\pi$.		
Γ_9 $a_2(1320)\pi$		
Γ_{10} $\rho \rho$		
Γ_{11} $\phi \pi$		
Γ_{12} $\eta \pi$		
Γ_{13} $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 14.7$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|ccc} & & -77 & \\ x_4 & & -74 & 17 \\ x_5 & & -15 & 2 & 0 \\ x_6 & & & x_1 & x_4 & x_5 \end{array}$$

 $\rho_3(1690)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.236 ± 0.013 OUR FIT				
0.243 ± 0.013 OUR AVERAGE				
$0.259^{+0.018}_{-0.019}$	BECKER	79	ASPK 0	17 $\pi^- p$ polarized
0.23 ± 0.02	CORDEN	79	OMEG	12–15 $\pi^- p \rightarrow$
0.22 ± 0.04	²⁸ MATTHEWS	71c	HDBC 0	$n^2 \pi$ $7 \pi^+ n \rightarrow \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.245 ± 0.006	²⁹ ESTABROOKS	75	RVUE	17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$

28 One-pion-exchange model used in this estimation.

29 From phase-shift analysis of HYAMS 75 data.

$\Gamma(\pi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_2
0.35±0.11	CASON	73	HBC	—	8,18.5 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.2

HOLMES 72 HBC + 10–12 $K^+ p$

<0.12

BALLAM 71B HBC — 16 $\pi^- p$

$\Gamma(\pi\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
0.332±0.026 OUR FIT	Error includes scale factor of 1.1.				
0.30 ±0.10	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow p4\pi$


$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_4
0.067±0.011 OUR FIT	Error includes scale factor of 1.2.				

0.118^{+0.039}_{-0.032} OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.

$0.191^{+0.040}_{-0.037}$	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ± 0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
$0.08^{+0.08}_{-0.03}$	CRENNELL	68B	HBC		6.0 $\pi^- p$

WEIGHTED AVERAGE
0.118+0.039-0.032 (Error scaled by 1.7)



$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

$\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_5/Γ_4
0.16±0.05 OUR FIT					
0.16±0.05	30 BARTSCH	70B HBC	+	$8 \pi^+ p$	
30 Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.					

 $[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	$(\Gamma_8+\Gamma_9+\Gamma_{10})/\Gamma_2$
0.94±0.09 OUR AVERAGE					
0.96±0.21	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$	
0.88±0.15	BALLAM	71B HBC	-	$16 \pi^- p$	
1 ± 0.15	BARTSCH	70B HBC	+	$8 \pi^+ p$	
consistent with 1	CASO	68 HBC	-	$11 \pi^- p$	

 $\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_{10}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.12±0.11		BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$	
0.56	66	KLIGER	74 HBC	-	$4.5 \pi^- p \rightarrow p4\pi$	
0.13±0.09		31 THOMPSON	74 HBC	+	$13 \pi^+ p$	
0.7 ± 0.15		BARTSCH	70B HBC	+	$8 \pi^+ p$	

31 $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable. $\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	$\Gamma_{10}/(\Gamma_8+\Gamma_9+\Gamma_{10})$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.48±0.16	CASO	68 HBC	-	$11 \pi^- p$	

 $\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_9/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.66±0.08	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$	
0.36±0.14	32 THOMPSON	74 HBC	+	$13 \pi^+ p$	
not seen	CASON	73 HBC	-	$8, 18.5 \pi^- p$	
0.6 ± 0.15	BARTSCH	70B HBC	+	$8 \pi^+ p$	
0.6	BALTAY	68 HBC	+	$7, 8.5 \pi^+ p$	

32 $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable. $\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_3/Γ_2
0.23±0.05 OUR AVERAGE					Error includes scale factor of 1.2.	
0.33±0.07		THOMPSON	74 HBC	+	$13 \pi^+ p$	
0.12±0.07		BALLAM	71B HBC	-	$16 \pi^- p$	
0.25±0.10		BALTAY	68 HBC	+	$7, 8.5 \pi^+ p$	
0.25±0.10		JOHNSTON	68 HBC	-	$7.0 \pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	95	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$
<0.09		KLIGER	74 HBC	-	$4.5 \pi^- p \rightarrow p4\pi$

$\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.11	BALTAY	68	HBC	+ 7.8.5 $\pi^+ p$

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.15	BALTAY	68	HBC	+ 7.8.5 $\pi^+ p$

 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.02	THOMPSON	74	HBC	+ 13 $\pi^+ p$

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.0158±0.0026 OUR FIT Error includes scale factor of 1.2.				
0.0130±0.0024 OUR AVERAGE				
0.013 ± 0.003	COSTA...	80	OMEG 0	+ 10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ± 0.004	33 MARTIN	78B	SPEC	- 10 $\pi p \rightarrow K_S^0 K^- p$

³³ From $(\Gamma_4\Gamma_6)^{1/2} = 0.056 \pm 0.034$ assuming $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

 $\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.22±0.08	CASON	73	HBC	- 8,18.5 $\pi^- p$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	FUKUI	88	SPEC 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

 $\rho_3(1690)$ REFERENCES

ALDE	95	ZPHY C66 379	+Binon, Bricman+	(GAMS Collab.) JP
ALDE	92C	ZPHY C54 553	+Bencheikh, Binon+	(BELG, SERP, KEK, LANL, LAPP)
FUKUI	88	PL B202 441	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIYA)
DENNEY	83	PR D28 2726	+Cranley, Firestone, Chapman+	(IOWA, MICH)
EVANGELISTA	81	NP B178 197	+ (BARI, BONN, CERN, DARE, LIVP+)	
ALPER	80	PL 94B 422	+Becker+ (AMST, CERN, CRAC, MPIM, OXF+)	
COSTA...	80	NP B175 402	Costa De Beauregard+ (BARI, BONN, CERN+)	
GORLICH	80	NP B174 16	+Niczyporuk+ (CRAC, MPIM, CERN, ZEEM)	
BECKER	79	NP B151 46	+Blanar, Blum+ (MPIM, CERN, ZEEM, CRAC)	
CORDEN	79	NP B157 250	+Dowell, Garvey+ (BIRM, RHEL, TELA, LOWC) JP	
BALTAY	78B	PR D17 62	+Cautis, Cohen, Csorna+ (COLU, BING)	
MARTIN	78B	NP B140 158	+Ozmutlu, Baldi, Bohringer, Dorsaz+ (DURH, GEVA)	
MARTIN	78D	PL 74B 417	+Ozmutlu, Baldi, Bohringer, Dorsaz+ (DURH, GEVA)	
ANTIPOV	77	NP B119 45	+Busnello, Damgaard, Kienzle+ (SERP, GEVA)	
GESSAROLI	77	NP B126 382	+ (BGNA, FIRZ, GENO, MILA, OXF, PAVI)	
BLUM	75	PL 57B 403	+Chabaud, Dietl, Garelick, Grayer+ (CERN, MPIM) JP	
ESTABROOKS	75	NP B95 322	+Martin (DURH)	
HYAMS	75	NP B100 205	+Jones, Weilhammer, Blum, Dietl+ (CERN, MPIM)	
ENGLER	74	PR D10 2070	+Kraemer, Toaff, Weisser, Diaz+ (CMU, CASE)	

GRAYER	74	NP B75 189	+Hyams, Blum, Dietl+	(CERN, MPIM)
KLIGER	74	SJNP 19 428	+Beketov, Grechko, Guzhanin, Dubovikov+	(ITEP)
		Translated from YAF 19 839.		
OREN	74	NP B71 189	+Cooper, Fields, Rhines, Allison+	(ANL, OXF)
THOMPSON	74	NP B69 220	+Gaidos, McIlwain, Miller, Mulera+	(PURD)
CASON	73	PR D7 1971	+Biswas, Kenney, Madden+	(NDAM)
BOWEN	72	PRL 29 890	+Earles, Faissler, Bliden+	(NEAS, STON)
HOLMES	72	PR D6 3336	+Ferbel, Slattery, Werner	(ROCH)
BALLAM	71B	PR D3 2606	+Chadwick, Guiragossian, Johnson+	(SLAC)
MATTHEWS	71C	NP B33 1	+Prentice, Yoon, Carroll+	(TNTO, WISC) JP
ARMENISE	70	LNC 4 199	+Ghidini, Foring, Cartacci+	(BARI, BGNA, FIRZ)
BARNHAM	70	PRL 24 1083	+Colley, Jobes, Kenyon, Pathak, Riddiford	(BIRM)
BARTSCH	70B	NP B22 109	+Kraus, Tsanos, Grote+	(AACH, BERL, CERN)
CASO	70	LNC 3 707	+Conte, Tomasini+	(GENO, HAMB, MILA, SACL)
STUNTEBECK	70	PL 32B 391	+Kenney, Deery, Biswas, Cason+	(NDAM)
ADERHOLZ	69	NP B11 259	+Bartsch+	(AACH3, BERL, CERN, JAGL, WARS)
ANDERSON	69	PRL 22 1390	+Collins+	(BNL, CMU)
ARMENISE	68	NC 54A 999	+Ghidini, Forino+	(BARI, BGNA, FIRZ, ORSAY) I
BALTAY	68	PRL 20 887	+Kung, Yeh, Ferbel+	(COLU, ROCH, RUTG, YALE) I
CASO	68	NC 54A 983	+Conte, Cords, Diaz+	(GENO, HAMB, MILA, SACL)
CRENNELL	68B	PL 28B 136	+Karshon, Lai, Scarr, Skillicorn	(BNL)
JOHNSTON	68	PRL 20 1414	+Prentice, Steenberg, Yoon	(TNTO, WISC) IJP
FOCACCI	66	PRL 17 890	+Kienzle, Levrat, Maglich, Martin	(CERN)
GOLDBERG	65	PL 17 354	+ (CERN, EPOL, ORSAY, MILA, CEA, SACL)	

OTHER RELATED PAPERS

BARNETT	83B	PL 120B 455	+Blockus, Burka, Chien, Christian+	(JHU)
EHRLICH	66	PR 152 1194	+Selove, Yuta	(PENN)
LEVRAT	66	PL 22 714	+Tolstrup+	(CERN Missing Mass Spect. Collab.)
SEGUINOT	66	PL 19 712	+Martin+	(CERN Missing Mass Spect. Collab.)
BELLINI	65	NC 40A 948	+DiCorato, Duimio, Fiorini	(MILA)
DEUTSCH...	65	PL 18 351	Deutschmann+	(AACH3, BERL, CERN)
FORINO	65	PL 19 65	+Gessaroli+	(BGNA, ORSAY, SACL)