

$K_2(1770)$ $I(J^P) = \frac{1}{2}(2^-)$ **THE $K_2(1770)$ AND THE $K_2(1820)$**

A partial-wave analysis of the $K^-\omega$ system based on about 100,000 $K^-p \rightarrow K^-\omega p$ events (ASTON 93) gives evidence for two $q\bar{q}$ D -wave states near 1.8 GeV. A previous analysis based on about 200,000 diffractively produced $K^-p \rightarrow K^-\pi^+\pi^-p$ events (DAUM 81) gave evidence for two $J^P = 2^-$ states in this region, with masses ~ 1780 MeV and ~ 1840 MeV and widths ~ 200 MeV, in good agreement with the results of ASTON 93. In contrast, the masses obtained using a single resonance do not agree well: ASTON 93 obtains 1728 ± 7 MeV, while DAUM 81 estimates ~ 1820 MeV. We conclude that there are indeed two K_2 resonances here.

We list under the $K_2(1770)$ other measurements that do not resolve the two-resonance structure of the enhancement.

 $K_2(1770)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1773± 8		1 ASTON	93 LASS		$11K^-p \rightarrow K^-\omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1810±20		FRAME	86 OMEG	+	$13K^+p \rightarrow \phi K^+p$
~1730		ARMSTRONG	83 OMEG	-	$18.5K^-p \rightarrow 3Kp$
~1780		2 DAUM	81C CNTR	-	$63K^-p \rightarrow K^-2\pi p$
1710±15	60	CHUNG	74 HBC	-	$7.3K^-p \rightarrow K^-\omega p$
1767± 6		BLIEDEN	72 MMS	-	$11\text{--}16K^-p$
1730±20	306	3 FIRESTONE	72B DBC	+	$12K^+d$
1765±40		4 COLLEY	71 HBC	+	$10K^+p \rightarrow K2\pi N$
1740		DENEGRIS	71 DBC	-	$12.6K^-d \rightarrow \bar{K}2\pi d$
1745±20		AGUILAR...	70C HBC	-	$4.6K^-p$
1780±15		BARTSCH	70C HBC	-	$10.1K^-p$
1760±15		LUDLAM	70 HBC	-	$12.6K^-p$

¹ From a partial wave analysis of the $K^-\omega$ system.

² From a partial wave analysis of the $K^-\pi^+\pi^-$ system.

³ Produced in conjunction with excited deuteron.

⁴ Systematic errors added correspond to spread of different fits.

$K_2(1770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
186±14		5 ASTON	93 LASS		11 $K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
140±40		FRAME	86 OMEG	+	13 $K^+ p \rightarrow \phi K^+ p$
~220		ARMSTRONG	83 OMEG	-	18.5 $K^- p \rightarrow 3K p$
~210		6 DAUM	81C CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
110±50	60	CHUNG	74 HBC	-	7.3 $K^- p \rightarrow K^- \omega p$
100±26		BLIEDEN	72 MMS	-	11–16 $K^- p$
210±30	306	7 FIRESTONE	72B DBC	+	12 $K^+ d$
90±70		8 COLLEY	71 HBC	+	10 $K^+ p \rightarrow K^- 2\pi N$
130		DENEGRIG	71 DBC	-	12.6 $K^- d \rightarrow \bar{K}^- 2\pi d$
100±50		AGUILAR-...	70C HBC	-	4.6 $K^- p$
138±40		BARTSCH	70C HBC	-	10.1 $K^- p$
50 ⁺⁴⁰ ₋₂₀		LUDLAM	70 HBC	-	12.6 $K^- p$

⁵ From a partial wave analysis of the $K^- \omega$ system.⁶ From a partial wave analysis of the $K^- 2\pi$ system.⁷ Produced in conjunction with excited deuteron.⁸ Systematic errors added correspond to spread of different fits. **$K_2(1770)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K_{\pi\pi}$	
$\Gamma_2 K_2^*(1430)\pi$	dominant
$\Gamma_3 K^*(892)\pi$	seen
$\Gamma_4 K f_2(1270)$	seen
$\Gamma_5 K\phi$	seen
$\Gamma_6 K\omega$	seen

 $K_2(1770)$ BRANCHING RATIOS

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	Γ_2/Γ_1
$(K_2^*(1430) \rightarrow K\pi)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
~0.03	DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$
~1.0	9 FIRESTONE 72B DBC + 12 $K^+ d$
<1.0	COLLEY 71 HBC 10 $K^+ p$
0.2 ±0.2	AGUILAR-... 70C HBC - 4.6 $K^- p$
<1.0	BARTSCH 70C HBC - 10.1 $K^- p$
1.0	BARBARO-... 69 HBC + 12.0 $K^+ p$

⁹ Produced in conjunction with excited deuteron.

$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$ Γ_3/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.23	DAUM	81C CNTR 63	$K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$ Γ_4/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.74	DAUM	81C CNTR 63	$K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K\phi)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
seen	ARMSTRONG 83	OMEG	—	$18.5 K^- p \rightarrow K^- \phi N$

 $\Gamma(K\omega)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
seen	OTTER 81	HBC	±	$8.25, 10, 16 K^\pm p$
seen	CHUNG 74	HBC	—	$7.3 K^- p \rightarrow K^- \omega p$

 $K_2(1770)$ REFERENCES

ASTON	93	PL B308 186	+Benz, Bird+ (SLAC, NAGO, CINC, INUS)
FRAME	86	NP B276 667	+Hughes, Lynch, Minto, McFadzean+ (GLAS)
ARMSTRONG	83	NP B221 1	+ (BARI, BIRM, CERN, MILA, CURIN+)
DAUM	81C	NP B187 1	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
OTTER	81	NP B181 1	(AACH3, BERL, LOIC, VIEN, BIRM, BELG, CERN+)
CHUNG	74	PL 51B 413	+Eisner, Protopopescu, Samios, Strand (BNL)
BLIEDEN	72	PL 39B 668	+Finocchiaro, Bowen, Earles+ (STON, NEAS)
FIRESTONE	72B	PR D5 505	+Goldhaber, Lissauer, Trilling (LBL)
COLLEY	71	NP B26 71	+Jobes, Kenyon, Pathak, Hughes+ (BIRM, GLAS)
DENEGRIS	71	NP B28 13	+Antich, Callahan, Carson, Chien, Cox+ (JHU) JP
AGUILAR-...	70C	PRL 25 54	Aguilar-Benitez, Barnes, Bassano, Chung+ (BNL)
BARTSCH	70C	PL 33B 186	+Deutschmann+ (AACH, BERL, CERN, LOIC, VIEN)
LUDLAM	70	PR D2 1234	+Sandweiss, Slaughter (YALE)
BARBARO-...	69	PRL 22 1207	Barbaro-Galtieri, Davis, Flatté+ (LRL)

OTHER RELATED PAPERS

BERLINGHIERI	67	PRL 18 1087	+Farber, Ferbel, Forman (ROCH) I
CARMONY	67	PRL 18 615	+Hendricks, Lander (UCSD)
JOBES	67	PL 26B 49	+Bassompierre, DeBaere+ (BIRM, CERN, BRUX)
BARTSCH	66	PL 22 357	+Deutschmann+ (AACH, BERL, CERN+)