

Heavy Bosons Other Than Higgs Bosons, Searches for

We list here various limits on charged and neutral heavy vector bosons (other than W 's and Z 's), heavy scalar bosons (other than Higgs bosons), vector or scalar leptoquarks, and axiglons.

W_R (Right-Handed W Boson) MASS LIMITS

Assuming a light right-handed neutrino, except for BEALL 82, LANGACKER 89B, and COLANGELO 91. $g_R = g_L$ assumed. [Limits in the section MASS LIMITS for W' below are also valid for W_R if $m_{\nu_R} \ll m_{W_R}$.] Some limits assume manifest left-right symmetry, *i.e.*, the equality of left- and right Cabibbo-Kobayashi-Maskawa matrices. For a comprehensive review, see LANGACKER 89B. Limits on the W_L - W_R mixing angle ζ are found in the next section. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 549		¹ BARENBOIM 97	RVUE	μ decay
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 220	95	² STAHL 97	RVUE	τ decay
> 220	90	³ ALLET 96	CNTR	β^+ decay
> 281	90	⁴ KUZNETSOV 95	CNTR	Polarized neutron decay
> 282	90	⁵ KUZNETSOV 94B	CNTR	Polarized neutron decay
> 439	90	⁶ BHATTACH... 93	RVUE	Z - Z' mixing
> 250	90	⁷ SEVERIJNS 93	CNTR	β^+ decay
		⁸ IMAZATO 92	CNTR	K^+ decay
> 475	90	⁹ POLAK 92B	RVUE	μ decay
> 240	90	¹⁰ AQUINO 91	RVUE	Neutron decay
> 496	90	¹⁰ AQUINO 91	RVUE	Neutron and muon decay
> 700		¹¹ COLANGELO 91	THEO	$m_{K_L^0} - m_{K_S^0}$
> 477	90	¹² POLAK 91	RVUE	μ decay
[none 540-23000]		¹³ BARBIERI 89B	ASTR	SN 1987A; light ν_R
> 300	90	¹⁴ LANGACKER 89B	RVUE	General
> 160	90	¹⁵ BALKE 88	CNTR	$\mu \rightarrow e\nu\bar{\nu}$
> 406	90	¹⁶ JODIDIO 86	ELEC	Any ζ
> 482	90	¹⁶ JODIDIO 86	ELEC	$\zeta = 0$
> 800		MOHAPATRA 86	RVUE	$SU(2)_L \times SU(2)_R \times U(1)$
> 400	95	¹⁷ STOKER 85	ELEC	Any ζ
> 475	95	¹⁷ STOKER 85	ELEC	$\zeta < 0.041$
		¹⁸ BERGSMA 83	CHRM	$\nu_\mu e \rightarrow \mu\nu_e$
> 380	90	¹⁹ CARR 83	ELEC	μ^+ decay
>1600		²⁰ BEALL 82	THEO	$m_{K_L^0} - m_{K_S^0}$
[> 4000]		STEIGMAN 79	COSM	Nucleosynthesis; light ν_R

¹ The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from K_L - K_S mass difference.

² STAHL 97 limit is from fit to τ -decay parameters.

- ³ ALLET 96 measured polarization-asymmetry correlaton in $^{12}\text{N}\beta^+$ decay. The listed limit assumes zero L - R mixing.
- ⁴ KUZNETSOV 95 limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed. See also KUZNETSOV 94B.
- ⁵ KUZNETSOV 94B limit is from measurements of the asymmetry $\langle \vec{p}_\nu \cdot \sigma_n \rangle$ in the β decay of polarized neutrons. Zero mixing assumed.
- ⁶ BHATTACHARYYA 93 uses Z - Z' mixing limit from LEP '90 data, assuming a specific Higgs sector of $\text{SU}(2)_L \times \text{SU}(2)_R \times \text{U}(1)$ gauge model. The limit is for $m_t=200$ GeV and slightly improves for smaller m_t .
- ⁷ SEVERIJNS 93 measured polarization-asymmetry correlation in $^{107}\text{In}\beta^+$ decay. The listed limit assumes zero L - R mixing. Value quoted here is from SEVERIJNS 94 erratum.
- ⁸ IMAZATO 92 measure positron asymmetry in $K^+ \rightarrow \mu^+ \nu_\mu$ decay and obtain $\xi P_\mu > 0.990$ (90%CL). If W_R couples to $u\bar{s}$ with full weak strength ($V_{us}^R=1$), the result corresponds to $m_{W_R} > 653$ GeV. See their Fig.4 for m_{W_R} limits for general $|V_{us}^R|^2=1-|V_{ud}^R|^2$.
- ⁹ POLAK 92B limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta=0$. Supersedes POLAK 91.
- ¹⁰ AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right symmetry assumed. Stronger of the two limits also includes muon decay results.
- ¹¹ COLANGELO 91 limit uses hadronic matrix elements evaluated by QCD sum rule and is less restrictive than BEALL 82 limit which uses vacuum saturation approximation. Manifest left-right symmetry assumed.
- ¹² POLAK 91 limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta=0$. Superseded by POLAK 92B.
- ¹³ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- ¹⁴ LANGACKER 89B limit is for any ν_R mass (either Dirac or Majorana) and for a general class of right-handed quark mixing matrices.
- ¹⁵ BALKE 88 limit is for $m_{\nu_{eR}} = 0$ and $m_{\nu_{\mu R}} \leq 50$ MeV. Limits come from precise measurements of the muon decay asymmetry as a function of the positron energy.
- ¹⁶ JODIDIO 86 is the same TRIUMF experiment as STOKER 85 (and CARR 83); however, it uses a different technique. The results given here are combined results of the two techniques. The technique here involves precise measurement of the end-point e^+ spectrum in the decay of the highly polarized μ^+ .
- ¹⁷ STOKER 85 is same TRIUMF experiment as CARR 83. Here they measure the decay e^+ spectrum asymmetry above 46 MeV/ c using a muon-spin-rotation technique. Assumed a light right-handed neutrino. Quoted limits are from combining with CARR 83.
- ¹⁸ BERGSMA 83 set limit $m_{W_2}/m_{W_1} > 1.9$ at CL = 90%.
- ¹⁹ CARR 83 is TRIUMF experiment with a highly polarized μ^+ beam. Looked for deviation from $V-A$ at the high momentum end of the decay e^+ energy spectrum. Limit from previous world-average muon polarization parameter is $m_{W_R} > 240$ GeV. Assumes a light right-handed neutrino.
- ²⁰ BEALL 82 limit is obtained assuming that W_R contribution to $K_L^0-K_S^0$ mass difference is smaller than the standard one, neglecting the top quark contributions. Manifest left-right symmetry assumed.

Limit on W_L - W_R Mixing Angle ζ

Lighter mass eigenstate $W_1 = W_L \cos \zeta - W_R \sin \zeta$. Light ν_R assumed unless noted.
 Values in brackets are from cosmological and astrophysical considerations.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.0333		21 BARENBOIM	97 RVUE	μ decay
< 0.04	90	22 MISHRA	92 CCFR	νN scattering
-0.0006 to 0.0028	90	23 AQUINO	91 RVUE	
[none 0.00001-0.02]		24 BARBIERI	89B ASTR	SN 1987A
< 0.040	90	25 JODIDIO	86 ELEC	μ decay
-0.056 to 0.040	90	25 JODIDIO	86 ELEC	μ decay

²¹ The quoted limit is from μ decay parameters. BARENBOIM 97 also evaluate limit from K_L - K_S mass difference.

²² MISHRA 92 limit is from the absence of extra large- x , large- y $\bar{\nu}_\mu N \rightarrow \bar{\nu}_\mu X$ events at Tevatron, assuming left-handed ν and right-handed $\bar{\nu}$ in the neutrino beam. The result gives $\zeta^2(1-2m_{W_1}^2/m_{W_2}^2) < 0.0015$. The limit is independent of ν_R mass.

²³ AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right asymmetry is assumed.

²⁴ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.

²⁵ First JODIDIO 86 result assumes $m_{W_R} = \infty$, second is for unconstrained m_{W_R} .

THE W' SEARCHES

Written October 1997 by K.S. Babu, C. Kolda, and J. March-Russell (IAS/Princeton).

Any electrically charged gauge boson outside of the Standard Model is generically denoted W' . A W' always couples to two different flavors of fermions, similar to the W boson. In particular, if a W' couples quarks to leptons it is a leptoquark gauge boson.

The most attractive candidate for W' is the W_R gauge boson associated with the left-right symmetric models [1]. These models seek to provide a spontaneous origin for parity violation in weak interactions. Here the gauge group is extended to $SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ with the Standard Model hypercharge identified as $Y = T_{3R} + (B-L)/2$, T_{3R} being the third component of $SU(2)_R$. The fermions transform under the gauge group in a left-right symmetric fashion: $q_L(3, 2, 1, 1/3) + q_R(3, 1, 2, 1/3)$ for quarks and $\ell_L(1, 2, 1, -1) + \ell_R(1, 1, 2, -1)$

for leptons. Note that the model requires the introduction of right-handed neutrinos, which can facilitate the see-saw mechanism for explaining the smallness of the ordinary neutrino masses. A Higgs bidoublet $\Phi(1, 2, 2, 0)$ is usually employed to generate quark and lepton masses and to participate in the electroweak symmetry breaking. Under left-right (or parity) symmetry, $q_L \leftrightarrow q_R$, $\ell_L \leftrightarrow \ell_R$, $W_L \leftrightarrow W_R$ and $\Phi \leftrightarrow \Phi^\dagger$.

After spontaneous symmetry breaking, the two W bosons of the model, W_L and W_R , will mix. The physical mass eigenstates are denoted as

$$W_1 = \cos \zeta W_L + \sin \zeta W_R, \quad W_2 = -\sin \zeta W_L + \cos \zeta W_R \quad (1)$$

with W_1 identified as the observed W boson. The most general Lagrangian that describes the interactions of the $W_{1,2}$ with the quarks can be written as [2]

$$\begin{aligned} \mathcal{L} = & -\frac{1}{\sqrt{2}} \bar{u} \gamma_\mu \left[\left(g_L \cos \zeta V^L P_L - g_R e^{i\omega} \sin \zeta V^R P_R \right) W_1^\mu \right. \\ & \left. + \left(g_L \sin \zeta V^L P_L + g_R e^{i\omega} \cos \zeta V^R P_R \right) W_2^\mu \right] d + h.c. \quad (2) \end{aligned}$$

where $g_{L,R}$ are the $SU(2)_{L,R}$ gauge couplings, $P_{L,R} = (1 \mp \gamma_5)/2$ and $V^{L,R}$ are the left- and right-handed CKM matrices in the quark sector. The phase ω reflects a possible complex mixing parameter in the W_L - W_R mass-squared matrix. Note that there is CP violation in the model arising from the right-handed currents even with only two generations. The Lagrangian for leptons is identical to that for quarks, with the replacements $u \rightarrow \nu$, $d \rightarrow e$ and the identification of $V^{L,R}$ with the CKM matrices in the leptonic sector.

If parity invariance is imposed on the Lagrangian, then $g_L = g_R$. Furthermore, the Yukawa coupling matrices that arise from coupling to the Higgs bidoublet Φ will be Hermitian. If in

addition the vacuum expectation values of Φ are assumed to be real, the quark and lepton mass matrices will also be Hermitian, leading to the relation $V^L = V^R$. Such models are called *manifest* left-right symmetric models and are approximately realized with a minimal Higgs sector [3]. If instead parity and CP are both imposed on the Lagrangian, then the Yukawa coupling matrices will be real symmetric and, after spontaneous CP violation, the mass matrices will be complex symmetric. In this case, which is known in the literature as *pseudo-manifest* left-right symmetry, $V^L = (V^R)^*$.

Indirect constraints: In minimal version of manifest or pseudo-manifest left-right symmetric models with $\omega = 0$ or π , there are only two free parameters, ζ and M_{W_2} , and they can be constrained from low energy processes. In the large M_{W_2} limit, stringent bounds on the angle ζ arise from three processes. (i) Nonleptonic K decays: The decays $K \rightarrow 3\pi$ and $K \rightarrow 2\pi$ are sensitive to small admixtures of right-handed currents. Assuming the validity of PCAC relations in the Standard Model it has been argued in Ref. 4 that the success in the $K \rightarrow 3\pi$ prediction will be spoiled unless $|\zeta| \leq 4 \times 10^{-3}$. (ii) $b \rightarrow s\gamma$: The amplitude for this process has an enhancement factor m_t/m_b relative to the Standard Model and thus can be used to constrain ζ yielding the limit $-0.01 \leq \zeta \leq 0.003$ [5]. (iii) Universality in weak decays: If the right-handed neutrinos are heavy, the right-handed admixture in the charged current will contribute to β decay and K decay, but not to the μ decay. This will modify the extracted values of V_{ud}^L and V_{us}^L . Demanding that the difference not upset the three generation unitarity of the CKM matrix, a bound $|\zeta| \leq 10^{-3}$ has been derived [6].

If the ν_R are heavy, leptonic and semileptonic processes do not constrain ζ since the emission of ν_R will not be kinematically allowed. However, if the ν_R is light enough to be emitted in μ decay and β decay, stringent limits on ζ do arise. For example, $|\zeta| \leq 0.039$ can be obtained from polarized μ decay [7] in the large M_{W_2} limit of the manifest left-right model. Alternatively, in the $\zeta = 0$ limit, there is a constraint $M_{W_2} \geq 484$ GeV from direct W_2 exchange. For the constraint on the case in which M_{W_2} is not taken to be heavy, see Ref. 2. There are also cosmological and astrophysical constraints on M_{W_2} and ζ in scenarios with a light ν_R . During nucleosynthesis the process $e^+e^- \rightarrow \nu_R\bar{\nu}_R$, proceeding via W_2 exchange, will keep the ν_R in equilibrium leading to an overproduction of ${}^4\text{He}$ unless M_{W_2} is greater than about 1 TeV [8]. Likewise the ν_{eR} produced via $e^-_R p \rightarrow n\nu_R$ inside a supernova must not drain too much of its energy, leading to limits $M_{W_2} > 16$ TeV and $|\zeta| \leq 3 \times 10^{-5}$ [9]. Note that models with light ν_R do not have a see-saw mechanism for explaining the smallness of the neutrino masses, though other mechanisms may arise in variant models [10].

The mass of W_2 is severely constrained (independent of the value of ζ) from K_L-K_S mass-splitting. The box diagram with exchange of one W_L and one W_R has an anomalous enhancement and yields the bound $M_{W_2} \geq 1.6$ TeV [11] for the case of manifest or pseudo-manifest left-right symmetry. If the ν_R have Majorana masses, another constraint arises from neutrinoless double β decay. Combining the experimental limit from ${}^{76}\text{Ge}$ decay with arguments of vacuum stability, a limit of $M_{W_2} \geq 1.1$ TeV has been obtained [12].

Direct search limits: Limits on M_{W_2} from direct searches depend on the available decay channels of W_2 . If ν_R is heavier

than W_2 , the decay $W_2^+ \rightarrow \ell_R^+ \nu_R$ will be forbidden kinematically. Assuming that ζ is small, the dominant decay of W_2 will be into dijets. UA2 [13] has excluded a W_2 in the mass range of 100 to 251 GeV in this channel. DØ excludes the mass range of 340 to 680 GeV [14], while CDF excludes the mass range of 300 to 420 GeV for such a W_2 [15]. If ν_R is lighter than W_2 , the decay $W_2^+ \rightarrow e_R^+ \nu_R$ is allowed. The ν_R can then decay into $e_R W_R^*$, leading to an $eejj$ signature. DØ has a limit of $M_{W_2} > 720$ GeV if $m_{\nu_R} \ll M_{W_2}$; the bound weakens, for example, to 650 GeV for $m_{\nu_R} = M_{W_2}/2$ [16]. CDF finds $M_{W_2} > 652$ GeV if ν_R is stable and much lighter than W_2 [17]. All of these limits assume manifest or pseudo-manifest left-right symmetry. See [16] for some variations in the limits if the assumption of left-right symmetry is relaxed.

Alternative models: W' gauge bosons can also arise in other models. We shall briefly mention some such popular models, but for details we refer the reader to the original literature. The *alternate* left-right model [18] is based on the same gauge group as the left-right model, but arises in the following way: In E_6 unification, there is an option to identify the right-handed down quarks as $SU(2)_R$ singlets or doublets. If they are $SU(2)_R$ doublets, one recovers the conventional left-right model; if they are singlets it leads to the alternate left-right model. A similar ambiguity exists in the assignment of left-handed leptons; the alternate left-right model assigns them to a $(1, 2, 2, 0)$ multiplet. As a consequence, the ordinary neutrino remains exactly massless in the model. One important difference from the usual left-right model is that the limit from the K_L-K_S mass difference is no longer applicable, since the d_R do not couple to the W_R . There is also no limit from polarized μ decay, since the $SU(2)_R$ partner of e_R can receive a large Majorana

mass. Other W' models include the un-unified Standard Model of Ref. 19 where there are two different SU(2) gauge groups, one each for the quarks and leptons; models with separate SU(2) gauge factors for each generation [20]; and the $SU(3)_C \times SU(3)_L \times U(1)$ model of Ref. 21.

Leptoquark gauge bosons: The $SU(3)_C \times U(1)_{B-L}$ part of the gauge symmetry discussed above can be embedded into a simple $SU(4)_C$ gauge group [22]. The model then will contain leptoquark gauge boson as well, with couplings of the type $\{(\bar{e}_L \gamma_\mu d_L + \bar{\nu}_L \gamma_\mu u_L) W'^\mu + (L \rightarrow R)\}$. The best limit on such leptoquark W' comes from nonobservation of $K_L \rightarrow \mu e$, which requires $M_{W'} \geq 1400$ TeV; for the corresponding limits on less conventional leptoquark flavor structures, see Ref. 23. Thus such a W' is inaccessible to direct searches with present machines which are sensitive to vector leptoquark masses of order 300 GeV only.

References

1. J.C. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974);
R.N. Mohapatra and J.C. Pati, Phys. Rev. **D11**, 566 (1975); *ibid.* Phys. Rev. **D11**, 2558 (1975);
G. Senjanovic and R.N. Mohapatra, Phys. Rev. **D12**, 1502 (1975).
2. P. Langacker and S. Uma Sankar, Phys. Rev. **D40**, 1569 (1989).
3. A. Masiero, R.N. Mohapatra, and R. Peccei, Nucl. Phys. **B192**, 66 (1981);
J. Basecq, *et al.*, Nucl. Phys. **B272**, 145 (1986).
4. J. Donoghue and B. Holstein, Phys. Lett. **113B**, 383 (1982).
5. K.S. Babu, K. Fujikawa, and A. Yamada, Phys. Lett. **B333**, 196 (1994);
P. Cho and M. Misiak, Phys. Rev. **D49**, 5894 (1994);

- T.G. Rizzo, Phys. Rev. **D50**, 3303 (1994).
6. L. Wolfenstein, Phys. Rev. **D29**, 2130 (1984).
 7. P. Herczeg, Phys. Rev. **D34**, 3449 (1986).
 8. G. Steigman, K.A. Olive, and D. Schramm, Nucl. Phys. **B180**, 497 (1981).
 9. R. Barbieri and R.N. Mohapatra, Phys. Rev. **D39**, 1229 (1989);
G. Raffelt and D. Seckel, Phys. Rev. Lett. **60**, 1793 (1988).
 10. D. Chang and R.N. Mohapatra, Phys. Rev. Lett. **58**, 1600 (1987);
K.S. Babu and X.G. He, Mod. Phys. Lett. **A4**, 61 (1989).
 11. G. Beall, M. Bender, and A. Soni, Phys. Rev. Lett. **48**, 848 (1982).
 12. R.N. Mohapatra, Phys. Rev. **D34**, 909 (1986).
 13. J. Alitti, *et al.* (UA2 Collaboration), Nucl. Phys. **B400**, 3 (1993).
 14. B. Abbott, *et al.* (DØ Collaboration), International Europhysics Conference on High Energy Physics, August 19-26, 1997, Jerusalem, Israel.
 15. F. Abe, *et al.* (CDF Collaboration), Phys. Rev. **D55**, R5263 (1997).
 16. S. Abachi, *et al.* (DØ Collaboration), Phys. Rev. Lett. **76**, 3271 (1996).
 17. F. Abe, *et al.* (CDF Collaboration), Phys. Rev. Lett. **74**, 2900 (1995).
 18. E. Ma, Phys. Rev. **D36**, 274 (1987);
K.S. Babu, X-G. He and E. Ma, Phys. Rev. **D36**, 878 (1987).
 19. H. Georgi and E. Jenkins, Phys. Rev. Lett. **62**, 2789 (1989);
Nucl. Phys. **B331**, 541 (1990).
 20. X. Li and E. Ma, Phys. Rev. Lett. **47**, 1788 (1981);
R.S. Chivukula, E.H. Simmons, and J. Terning, Phys. Lett. **B331**, 383 (1994);
D.J. Muller and S. Nandi, Phys. Lett. **B383**, 345 (1996).

21. F. Pisano, V. Pleitez, Phys. Rev. **D46**, 410 (1992);
 P. Frampton, Phys. Rev. Lett. **69**, 2889 (1992).
22. J.C. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974).
23. A. Kuznetsov and N. Mikheev, Phys. Lett. **B329**, 295 (1994);
 G. Valencia and S. Willenbrock, Phys. Rev. **D50**, 6843 (1994).

MASS LIMITS for W' (A Heavy-Charged Vector Boson Other Than W) in Hadron Collider Experiments

Couplings of W' to quarks and leptons are taken to be identical with those of W . The following limits are obtained from $p\bar{p} \rightarrow W'X$ with W' decaying to the mode indicated in the comments. New decay channels (e.g., $W' \rightarrow WZ$) are assumed to be suppressed. UA1 and UA2 experiments assume that the $t\bar{b}$ channel is not open.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>720	95	²⁶ ABACHI	96C D0	$W' \rightarrow e\nu_e$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 300–420	95	²⁷ ABE	97G CDF	$W' \rightarrow q\bar{q}$
>610	95	²⁸ ABACHI	95E D0	$W' \rightarrow e\nu_e$ and $W' \rightarrow \tau\nu_\tau \rightarrow e\nu\nu\bar{\nu}$
>652	95	²⁹ ABE	95M CDF	$W' \rightarrow e\nu_e$
>251	90	³⁰ ALITTI	93 UA2	$W' \rightarrow q\bar{q}$
none 260–600	95	³¹ RIZZO	93 RVUE	$W' \rightarrow q\bar{q}$
>520	95	³² ABE	91F CDF	$W' \rightarrow e\nu, \mu\nu$
none 101–158	90	³³ ALITTI	91 UA2	$W' \rightarrow q\bar{q}$
>220	90	³⁴ ALBAJAR	89 UA1	$W' \rightarrow e\nu$
>209	90	³⁵ ANSARI	87D UA2	$W' \rightarrow e\nu$
>210	90	³⁶ ARNISON	86B UA1	$W' \rightarrow e\nu$
>170	90	³⁷ ARNISON	83D UA1	$W' \rightarrow e\nu$

²⁶ For bounds on W_R with nonzero right-handed mass, see Fig. 5 from ABACHI 96C.

²⁷ ABE 97G search for new particle decaying to dijets.

²⁸ ABACHI 95E assume that the decay $W' \rightarrow WZ$ is suppressed and that the neutrino from W' decay is stable and has a mass significantly less $m_{W'}$.

²⁹ ABE 95M assume that the decay $W' \rightarrow WZ$ is suppressed and the (right-handed) neutrino is light, noninteracting, and stable. If $m_\nu=60$ GeV, for example, the effect on the mass limit is negligible.

³⁰ ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $\Gamma(W')/m_{W'} = \Gamma(W)/m_W$ and $B(W' \rightarrow jj) = 2/3$. This corresponds to W_R with $m_{\nu_R} > m_{W_R}$ (no leptonic decay) and $W_R \rightarrow t\bar{b}$ allowed. See their Fig. 4 for limits in the $m_{W'}-B(q\bar{q})$ plane.

³¹ RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

³² ABE 91F assume leptonic branching ratio of 1/12 for each lepton flavor. The limit from the $e\nu$ ($\mu\nu$) mode alone is 490 (435) GeV. These limits apply to W_R if $m_{\nu_R} \lesssim 15$ GeV and ν_R does not decay in the detector. Cross section limit $\sigma \cdot B < (1-10)$ pb is given for $m_{W'} = 100-550$ GeV; see Fig. 2.

- ³³ ALITTI 91 search is based on two-jet invariant mass spectrum, assuming $B(W' \rightarrow q\bar{q}) = 67.6\%$. Limit on $\sigma \cdot B$ as a function of two-jet mass is given in Fig. 7.
- ³⁴ ALBAJAR 89 cross section limit at 630 GeV is $\sigma(W') B(e\nu) < 4.1$ pb (90% CL).
- ³⁵ See Fig. 5 of ANSARI 87D for the excluded region in the $m_{W'q} - [(g_{W'q})^2 B(W' \rightarrow e\bar{\nu})]$ plane. Note that the quantity $(g_{W'q})^2 B(W' \rightarrow e\bar{\nu})$ is normalized to unity for the standard W couplings.
- ³⁶ ARNISON 86B find no excess at large p_T in 148 $W \rightarrow e\nu$ events. Set limit $\sigma \times B(e\nu) < 10$ pb at CL = 90% at $E_{\text{cm}} = 546$ and 630 GeV.
- ³⁷ ARNISON 83D find among 47 $W \rightarrow e\nu$ candidates no event with excess p_T . Also set $\sigma \times B(e\nu) < 30$ pb with CL = 90% at $E_{\text{cm}} = 540$ GeV.

MASS LIMITS for Z' (Heavy Neutral Vector Boson Other Than Z) THE Z' SEARCHES

Written October 1997 by K.S. Babu, C. Kolda, and J. March-Russell (IAS/Princeton).

If the Standard Model is enhanced by additional gauge symmetries or embedded into a larger gauge group, there will arise new heavy gauge bosons, some of which generically are electrically neutral. Such a gauge boson is called a Z' . Consider the most general renormalizable Lagrangian describing the complete set of interactions of the neutral gauge bosons among themselves and with fermions, which is that of the Standard Model plus the following new pieces [1,2,3]:

$$\begin{aligned} \mathcal{L}_{Z'} = & -\frac{1}{4} \widehat{F}'_{\mu\nu} \widehat{F}'^{\mu\nu} - \frac{\sin \chi}{2} \widehat{F}'_{\mu\nu} \widehat{F}^{\mu\nu} + \frac{1}{2} \widehat{M}_{Z'}^2 \widehat{Z}'_\mu \widehat{Z}'^\mu \\ & + \delta \widehat{M}^2 \widehat{Z}'_\mu \widehat{Z}'^\mu - \frac{\widehat{g}'}{2} \sum_i \bar{\psi}_i \gamma^\mu (f_V^i - f_A^i \gamma^5) \psi_i \widehat{Z}'_\mu \end{aligned} \quad (1)$$

where $\widehat{F}_{\mu\nu}, \widehat{F}'_{\mu\nu}$ are the field strength tensors for the hypercharge \widehat{B}_μ gauge boson and the Z' respectively before any diagonalizations are performed, ψ_i are the matter fields with Z' vector and axial charges f_V^i and f_A^i , and \widehat{Z}'_μ is the electroweak Z boson in this basis. (See the Review on “Electroweak Model and Constraints on New Physics” for the Standard Model pieces of the Lagrangian.) The mass terms are assumed to come from

spontaneous symmetry breaking via scalar expectation values. The above Lagrangian is general to all abelian and non-abelian extensions, except that $\chi = 0$ for the non-abelian case since then $\widehat{F}'_{\mu\nu}$ is not gauge invariant. Most analyses take $\chi = 0$ even for the abelian case.

Going to the physical eigenbasis requires diagonalizing both the gauge kinetic and mass terms, with mass eigenstates denoted Z_1 and Z_2 , where we choose Z_1 to be the observed Z boson. The interaction Lagrangian for Z_1 has the form, to leading order in the mixing angle ξ ($s_W \equiv \sin \theta_W$, etc.):

$$\begin{aligned} \mathcal{L}_{Z_1} = & -\frac{e}{2s_W c_W} \left(1 + \frac{\alpha T}{2}\right) \bar{\psi}_i \gamma^\mu \left\{ \left(g_V^i + \xi \tilde{f}_V^i\right) \right. \\ & \left. - \left(g_A^i + \xi \tilde{f}_A^i\right) \gamma^5 \right\} \psi_i Z_{1\mu} \end{aligned} \quad (2)$$

where

$$\xi \simeq \frac{-\cos \chi (\delta \widehat{M}^2 + \widehat{M}_Z^2 s_W \sin \chi)}{\widehat{M}_{Z'}^2 - \widehat{M}_Z^2 \cos^2 \chi + \widehat{M}_Z^2 s_W^2 \sin^2 \chi + 2 \delta \widehat{M}^2 s_W \sin \chi} . \quad (3)$$

We have made the identifications $g_A^i = T_3^i$, $g_V^i = T_3^i - 2Q^i s_*^2$, $\tilde{f}_{V,A}^i = (\widehat{g}' s_W c_W / e \cos \chi) f_{V,A}^i$, and s_W^2 is identified to be the $s_{M_Z}^2$ defined in the ‘‘Electroweak Model and Constraints on New Physics’’ review. Note that the value of the weak angle that appears in the vector coupling is shifted by the S and T oblique parameters:

$$s_*^2 = s_W^2 + \frac{1}{s_W^2 - c_W^2} \left(\frac{1}{4} \alpha S - c_W^2 s_W^2 \alpha T \right) . \quad (4)$$

Recall that $\rho = 1 + \alpha T$ defines the usual ρ parameter. In the presence of Z - Z' mixing, the oblique parameters receive

contributions [4]:

$$\begin{aligned}\alpha S &= 4\xi c_W^2 s_W \tan \chi \\ \alpha T &= \xi^2 \left(\frac{M_{Z_2}^2}{M_{Z_1}^2} - 1 \right) + 2\xi s_W \tan \chi \\ \alpha U &= 0\end{aligned}\tag{5}$$

to leading order in small ξ . These contributions are in addition to those coming from top quark and Higgs boson loops in the Standard Model. (This is in contrast to the ‘‘Electroweak Model and Constraints on New Physics’’ Review in which oblique parameters are defined to be zero for reference values of m_t and M_H .) Note that nonzero Z - Z' contributions to S arise only in the presence of kinetic mixing.

The corresponding $Z_2 \bar{\psi} \psi$ interaction Lagrangian is:

$$\mathcal{L}_{Z_2} = -\frac{e}{2s_W c_W} \bar{\psi}_i \gamma^\mu \{ (h_V^i - g_V^i \xi) - (h_A^i - g_A^i \xi) \gamma^5 \} \psi_i Z_{2\mu}\tag{6}$$

with the following definitions:

$$\begin{aligned}h_V^i &= \tilde{f}_V^i + \tilde{s}(T_3^i - 2Q^i) \tan \chi \\ h_A^i &= \tilde{f}_A^i + \tilde{s}T_3^i \tan \chi \\ \tilde{s} &= s_W + \frac{s_W^3}{c_W^2 - s_W^2} \left(\frac{1}{4c_W^2} \alpha S - \frac{1}{2} \alpha T \right)\end{aligned}\tag{7}$$

where the last equation defines a weak angle appropriate for the Z_2 interactions.

If the Z' charges are generation-dependent, there exist severe constraints in the first two generations coming from precision measurements such as the K_L - K_S mass splitting and $B(\mu \rightarrow 3e)$ owing to the lack of GIM suppression in the Z' interactions; however, constraints on a Z' which couples

differently only to the third generation are somewhat weaker. (It will be assumed in the Z -pole constraint section that the Z' couples identically to all three generations of matter; all other results are general.) If the new Z' interactions commute with the Standard Model gauge group, then per generation, there are only five independent $Z'\bar{\psi}\psi$ couplings; we can choose them to be \tilde{f}_V^u , \tilde{f}_A^u , \tilde{f}_V^d , \tilde{f}_V^e , and \tilde{f}_A^e . All other couplings can be determined in terms of these, *e.g.*, $\tilde{f}_V^\nu = (\tilde{f}_V^e + \tilde{f}_A^e)/2$.

Canonical models: One of the prime motivations for an additional Z' has come from string theory in which certain compactifications lead naturally to an E_6 gauge group, or one of its subgroups. E_6 contains two U(1) factors beyond the Standard Model, a basis for which is formed by the two groups $U(1)_\chi$ and $U(1)_\psi$, defined via the decompositions $E_6 \rightarrow SO(10) \times U(1)_\psi$ and $SO(10) \rightarrow SU(5) \times U(1)_\chi$; one special case often encountered is $U(1)_\eta$ where $Z_\eta = \sqrt{\frac{3}{8}}Z_\chi + \sqrt{\frac{5}{8}}Z_\psi$. The charges of the SM fermions under these U(1)'s, and a discussion of their experimental signals, can be found in Ref. 5.

It is also common to express experimental bounds in terms of a toy Z' usually denoted Z_{SM} . This Z_{SM} , of arbitrary mass, couples to the SM fermions identically to the usual Z .

Almost all analyses of Z' physics have worked with one of these canonical models and have assumed zero kinetic mixing at the weak scale.

Experimental constraints: There are three primary sets of constraints on the existence of a Z' which will be considered here: precision measurements of neutral-current processes at low energies, Z -pole constraints on Z - Z' mixing, and direct search constraints from production at very high energies. In principle, one usually expects other new states to appear at the same scale as the Z' , including its symmetry-breaking sector

and any additional fermions necessary for anomaly cancellation. However, because these states are highly model-dependent, we will not include searches for them, or Z' decays to them, in the bounds that follow.

Low-energy constraints: After the breaking of the new gauge group and the usual electroweak breaking, the Z of the Standard Model can mix with the Z' , with mixing angle ξ defined above. As already discussed, this Z - Z' mixing implies a shift in the usual oblique parameters [S, T, U defined in Eq. (5)]. Current bounds on S and T translate into stringent constraints on the mixing angle, ξ , requiring $\xi \ll 1$; similar constraints on ξ arise from the LEP Z -pole data. Thus we will only consider the small- ξ limit henceforth.

Whether or not the new gauge interactions are parity violating, stringent constraints can arise from atomic parity violation (APV) and polarized electron-nucleon scattering experiments [6]. At low energies, the effective neutral-current Lagrangian is conventionally written:

$$\mathcal{L}_{\text{NC}} = \frac{G_F}{\sqrt{2}} \sum_{q=u,d} \{ C_{1q} (\bar{e} \gamma_\mu \gamma^5 e) (\bar{q} \gamma^\mu q) + C_{2q} (\bar{e} \gamma_\mu e) (\bar{q} \gamma^\mu \gamma^5 q) \} . \quad (8)$$

APV experiments are sensitive only to C_{1u} and C_{1d} (see the ‘‘Electroweak Model and Constraints on New Physics’’ Review for the nuclear weak charge, Q_W , in terms of the C_{1q}) where in the presence of the Z and Z' :

$$C_{1q} = 2(1 + \alpha T) (g_A^e + \xi \tilde{f}_A^e) (g_V^q + \xi \tilde{f}_V^q) + 2r (h_A^e - \xi g_A^e) (h_V^q - \xi g_V^q) \quad (9)$$

where $r = (M_{Z_1}/M_{Z_2})^2$. The r -dependent terms arise from Z_2 exchange and can interfere constructively or destructively with the Z_1 contribution. In the limit $\xi = r = 0$, this reduces to

the Standard Model expression. Polarized electron scattering is sensitive to both the C_{1q} and C_{2q} couplings, again as discussed in the “Electroweak Model and Constraints on New Physics” Review. The C_{2q} can be derived from the expression for C_{1q} with the complete interchange $V \leftrightarrow A$.

Stringent limits also arise from neutrino-hadron scattering. One usually expresses experimental results in terms of the effective 4-fermion operators $(\bar{\nu}\gamma_\mu\nu)(\bar{q}_{L,R}\gamma^\mu q_{L,R})$ with coefficients $(2\sqrt{2}G_F)\epsilon_{L,R}(q)$. (Again, see the “Electroweak Model and Constraints on New Physics” Review.) In the presence of the Z and Z' , the $\epsilon_{L,R}(q)$ are given by:

$$\begin{aligned} \epsilon_{L,R}(q) = & \frac{1 + \alpha T}{2} \left\{ (g_V^q \pm g_A^q)[1 + \xi(\tilde{f}_V^\nu \pm \tilde{f}_A^\nu)] + \xi(\tilde{f}_V^q \pm \tilde{f}_A^q) \right\} \\ & + \frac{r}{2} \left\{ (h_V^q \pm h_A^q)(h_V^\nu \pm h_A^\nu) - \xi(g_V^q \pm g_A^q)(h_V^\nu \pm h_A^\nu) \right. \\ & \left. - \xi(h_V^q \pm h_A^q) \right\} . \end{aligned} \quad (10)$$

Again, the r -dependent terms arise from Z_2 -exchange.

Z-pole constraints: Electroweak measurements made at LEP and SLC while sitting on the Z resonance are generally sensitive to Z' physics only through the mixing with the Z unless the Z and Z' are very nearly degenerate, a possibility we ignore. Constraints on the allowed mixing angle and Z couplings arise by fitting all data simultaneously to the *ansatz* of Z - Z' mixing. For any observable, \mathcal{O} , the shift in that observable, $\Delta\mathcal{O}$, can be expressed (following the procedure of Ref. 7) as:

$$\frac{\Delta\mathcal{O}}{\mathcal{O}} = \mathcal{A}_\mathcal{O}^S \alpha S + \mathcal{A}_\mathcal{O}^T \alpha T + \xi \sum_i \mathcal{B}_\mathcal{O}^{(i)} \tilde{f}^i \quad (11)$$

where i runs over the 5 independent $Z'\bar{\psi}\psi$ couplings listed earlier (assuming a Z' couplings commute with the generation

and gauge symmetries of the Standard Model; this is the only place where we enforce such a restriction). The coefficients $\mathcal{A}_{\mathcal{O}}^{S,T}$ and $\mathcal{B}_{\mathcal{O}}^{(i)}$, which are functions only of the Standard Model parameters, are given in Table 1. The first 5 observables are directly measured at LEP and SLC, while \bar{A}_e , \bar{A}_b and \bar{A}_c are measured via the asymmetries $\bar{A}_{FB}^{(0,f)} = \frac{3}{4}\bar{A}_e\bar{A}_f$ and $A_{LR}^0 = \bar{A}_e$ as defined in the ‘‘Electroweak Model and Constraints on New Physics’’ Review. As an example, the shift in \bar{A}_e due to Z' physics is given by

$$\frac{\Delta\bar{A}_e}{\bar{A}_e} = -24.9\alpha S + 17.7\alpha T - 26.7\xi\tilde{f}_V^e + 2.0\xi\tilde{f}_A^e . \quad (12)$$

Table 1: Expansion coefficients for shifts in Z -pole observables normalized to the Standard Model value of the observable [7,3].

\mathcal{O}	$\mathcal{A}_{\mathcal{O}}^S$	$\mathcal{A}_{\mathcal{O}}^T$	$\mathcal{B}_{\mathcal{O}}^{Vu}$	$\mathcal{B}_{\mathcal{O}}^{Au}$	$\mathcal{B}_{\mathcal{O}}^{Vd}$	$\mathcal{B}_{\mathcal{O}}^{Ve}$	$\mathcal{B}_{\mathcal{O}}^{Ae}$
Γ_Z	-0.49	1.35	-0.89	-0.40	0.37	0.37	0
R_ℓ	-0.39	0.28	-1.3	-0.56	0.52	0.30	4.0
σ_h	0.046	-0.033	0.50	0.22	-0.21	-1.0	-4.0
R_b	0.085	-0.061	-1.4	-2.1	0.29	0	0
R_c	-0.16	0.12	2.7	4.1	-0.59	0	0
\bar{A}_e	-24.9	17.7	0	0	0	-26.7	2.0
\bar{A}_b	-0.32	0.23	0.71	0.71	-1.73	0	0
\bar{A}_c	-2.42	1.72	3.89	-1.49	0	0	0
M_W^2	-0.93	1.43	0	0	0	0	0

High-energy indirect constraints: At $\sqrt{s} < M_{Z_2}$, but off the Z_1 pole, strong constraints on new Z' physics arise from measurements of deviations of asymmetries and leptonic and hadronic cross sections from their Standard Model predictions. These processes are sensitive not only to Z - Z' mixing but also to direct Z_2 exchange primarily through γ - Z_2 and Z_1 - Z_2 interference; therefore information on the Z_2 couplings and mass can be extracted that is not accessible via Z - Z' mixing alone.

Far below the Z_2 mass scale, experiment is only sensitive to the scaled Z_2 couplings $(\sqrt{s}/M_{Z_2}) \cdot h_{V,A}^i$ so the Z_2 mass and overall magnitude of the couplings cannot both be extracted. However as \sqrt{s} approaches M_{Z_2} the Z_2 exchange can no longer be approximated by a contact interaction and the mass and couplings can be simultaneously extracted.

Z' studies done before LEP relied heavily on this approach; see, *e.g.*, Ref. 8. LEP has also done similar work using data collected above the Z peak; see, *e.g.*, Ref. 9. For indirect Z' searches at future facilities, see, *e.g.* Refs. 10 and 11.

Direct-search constraints: Finally, high-energy experiments have searched for on-shell Z' (here Z_2) production and decay. Searches can be classified by the initial state off of which the Z' is produced, and the final state into which the Z' decays; we will not include here exotic decays of a Z' . Experiments to date have been sensitive to Z' production via their coupling to quarks ($p\bar{p}$ colliders), to electrons (e^+e^-) or to both (ep).

For a heavy Z' ($M_{Z_2} \gg M_{Z_1}$), the best limits come from $p\bar{p}$ machines via Drell-Yan production and subsequent decay to charged leptons. For $M_{Z_2} > 600$ GeV, CDF [12] quotes limits on $\sigma(p\bar{p} \rightarrow Z_2 X) \cdot B(Z_2 \rightarrow \ell^+\ell^-) < 0.04$ pb at 95% C.L. for $\ell = e + \mu$ combined; DØ [13] quotes $\sigma \cdot B < 0.025$ pb for $\ell = e$.

For $M_{Z_2} < 600$ GeV, the mass dependence is complicated and one should refer to the original literature. For studies of the search capabilities of future facilities, see *e.g.* Ref. 10.

If the Z' has suppressed, or no, couplings to leptons (*i.e.*, it is leptophobic) then experimental sensitivities are much weaker. In particular, searches for a Z' via hadronic decays at DØ [14] are able to rule out a Z' with quark couplings identical to those of the Z only in the mass range $365 \text{ GeV} < M_{Z_2} < 615 \text{ GeV}$; CDF [15] cannot exclude even this range. Additionally, UA2 [16] finds $\sigma \cdot B(Z' \rightarrow jj) < 11.7 \text{ pb}$ at 90% C.L. for $M_{Z'} > 200 \text{ GeV}$ and more complicated bounds in the range $130 \text{ GeV} < M_{Z'} < 200 \text{ GeV}$.

For a light Z' ($M_{Z'} < M_Z$) direct searches in e^+e^- colliders have ruled out any Z' unless it has extremely weak couplings to leptons. For a combined analysis of the various pre-LEP experiments see Ref. 8.

References

1. B. Holdom, Phys. Lett. **166B**, 196 (1986).
2. F. del Aguila, Acta Phys. Polon. **B25**, 1317 (1994);
F. del Aguila, M. Cvetič and P. Langacker, Phys. Rev. **D52**, 37 (1995).
3. K.S. Babu, C. Kolda and J. March-Russell, Phys. Rev. **D54**, 4635 (1996);
K.S. Babu, C. Kolda, and J. March-Russell, hep-ph/9710441.
4. B. Holdom, Phys. Lett. **B259**, 329 (1991).
5. J. Hewett and T. Rizzo, Phys. Rept. **183**, 193 (1989).
6. J. Kim, *et al.*, Rev. Mod. Phys. **53**, 211 (1981);
U. Amaldi, *et al.*, Phys. Rev. **D36**, 1385 (1987);
W. Marciano and J. Rosner, Phys. Rev. Lett. **65**, 2963 (1990) (*Erratum*: **68** 898 (1992));
K. Mahanthappa and P. Mohapatra, Phys. Rev. **D43**, 3093 (1991) (*Erratum*: **D44** 1616 (1991));

- P. Langacker and M. Luo, Phys. Rev. **D45**, 278 (1992);
 P. Langacker, M. Luo and A. Mann, Rev. Mod. Phys. **64**, 87 (1992).
7. G. Altarelli, *et al.*, Mod. Phys. Lett. **A5**, 495 (1990);
ibid., Phys. Lett. **B263**, 459 (1991).
 8. L. Durkin and P. Langacker, Phys. Lett. **166B**, 436 (1986).
 9. T. Burgsmüller (DELPHI Collaboration), *HEP'97 Conference* (Jerusalem, 1997), <http://wwwcn.cern.ch/~pubxx/www/delsec/conferences/jerusalem>;
 S. Riemann (L3 Collaboration), *Beyond the Standard Model V* (Balholm, 1997), <http://hpl3sn02.cern.ch/conferences/talks97.html>.
 10. M. Cvetič and S. Godfrey, hep-ph/9504216, in *Electroweak Symmetry Breaking and Beyond the Standard Model*, Eds. T. Barklow, *et al.* (World Scientific 1995).
 11. T. Rizzo, Phys. Rev. **D55**, 5483 (1997).
 12. CDF Collaboration, Phys. Rev. Lett. **79**, 2191 (1997).
 13. DØ Collaboration, *XVIII International Conf. on Lepton Photon Interactions* (June 1997), <http://D0sgi0.fnal.gov/public/new/conferences/lp97.html>.
 14. DØ Collaboration, *XVIII International Conference on Lepton Photon Interactions* (June 1997), see URL above.
 15. F. Abe *et al.*, (CDF Collaboration), Phys. Rev. **D55**, 5263R (1997).
 16. J. Alitti, *et al.*, (UA2 Collaboration), Nucl. Phys. **B400**, 3 (1993).

Limits for Z'_{SM}

Z'_{SM} is assumed to have couplings with quarks and leptons which are identical to those of Z .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>690	95	38 ABE	97s CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-, \mu^+\mu^-$
>779	95	39,40 LANGACKER	92B RVUE	Electroweak

••• We do not use the following data for averages, fits, limits, etc. •••

>490	95	ABACHI	96D D0	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>505	95	41 ABE	95 CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>398	95	42 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>237	90	43 ALITTI	93 UA2	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>119	90	44 ALLEN	93 CALO	$\nu e \rightarrow \nu e$
none 490–560	95	45 RIZZO	93 RVUE	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>412	95	ABE	92B CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$, $\mu^+\mu^-$
>387	95	46 ABE	91D CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>307	90	47 GEIREGAT	91 CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>426	90	48 ABE	90F VNS	e^+e^-
>208	90	49 HAGIWARA	90 RVUE	e^+e^-
>173	90	50 ALBAJAR	89 UA1	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>180	90	51 ANSARI	87D UA2	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>160	90	52 ARNISON	86B UA1	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$

³⁸ ABE 97S limit is obtained assuming that Z' decays to known fermions only.

³⁹ LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

⁴⁰ LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0086 < \theta < 0.0005$.

⁴¹ ABE 95 limit is obtained assuming that Z' decays to known fermions only.

⁴² VILAIN 94B assume $m_t = 150$ GeV.

⁴³ ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $B(Z' \rightarrow q\bar{q})=0.7$. See their Fig. 5 for limits in the $m_{Z'}-B(q\bar{q})$ plane.

⁴⁴ ALLEN 93 limit is from total cross section for $\nu e \rightarrow \nu e$, where $\nu = \nu_e, \nu_\mu, \bar{\nu}_\mu$.

⁴⁵ RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

⁴⁶ ABE 91D give $\sigma(Z') \cdot B(e^+e^-) < 1.31$ pb (95%CL) for $m_{Z'} > 200$ GeV at $E_{cm} = 1.8$ TeV. Limits ranging from 2 to 30 pb are given for $m_{Z'} = 100-200$ GeV.

⁴⁷ GEIREGAT 91 limit is from comparison of g_V^e from $\nu_\mu e$ scattering with $\Gamma(Z \rightarrow ee)$ from LEP. Zero mixing assumed.

⁴⁸ ABE 90F use data for $R, R_{\ell\ell},$ and $A_{\ell\ell}$. They fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

⁴⁹ HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-, \tau^+\tau^-$, and hadron cross sections and asymmetries.

⁵⁰ ALBAJAR 89 cross section limit at 630 GeV is $\sigma(Z') B(ee) < 4.2$ pb (90% CL).

⁵¹ See Fig. 5 of ANSARI 87D for the excluded region in the $m_{Z'}-[(g_{Z'q})^2 B(Z' \rightarrow e^+e^-)]$ plane. Note that the quantity $(g_{Z'q})^2 B(Z' \rightarrow e^+e^-)$ is normalized to unity for the standard Z couplings.

⁵² ARNISON 86B find no excess e^+e^- pairs among 13 pairs from Z . Set limit $\sigma \times B(e^+e^-) < 13$ pb at CL = 90% at $E_{cm} = 546$ and 630 GeV.

Limits for Z_{LR}

Z_{LR} is the extra neutral boson in left-right symmetric models. $g_L = g_R$ is assumed unless noted. Values in parentheses assume stronger constraint on the Higgs sector, usually motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>630	95	⁵³ ABE	97S CDF	$\rho\bar{p}; Z'_{LR} \rightarrow e^+e^-, \mu^+\mu^-$
>389	95	^{54,55} LANGACKER	92B RVUE	Electroweak
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>190	95	⁵⁶ BARATE	97B ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>445	95	⁵⁷ ABE	95 CDF	$\rho\bar{p}; Z'_{LR} \rightarrow e^+e^-$
>253	95	⁵⁸ VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>130	95	⁵⁹ ADRIANI	93D L3	Z parameters
(> 1500)	90	⁶⁰ ALTARELLI	93B RVUE	Z parameters
none 490–560	95	⁶¹ RIZZO	93 RVUE	$\rho\bar{p}; Z'_{LR} \rightarrow q\bar{q}$
>310	95	⁶² ABE	92B CDF	$\rho\bar{p}$
>230	95	⁶³ ABE	92B CDF	$\rho\bar{p}$
(> 900)	90	⁶⁴ DELAGUILA	92 RVUE	
(> 1400)		⁶⁵ LAYSSAC	92B RVUE	Z parameters
(> 564)	90	⁶⁶ POLAK	92 RVUE	μ decay
>474	90	⁶⁷ POLAK	92B RVUE	Electroweak
(> 1340)		⁶⁸ RENTON	92 RVUE	
(> 800)	90	⁶⁹ ALTARELLI	91B RVUE	Z parameters
(> 795)	90	⁷⁰ DELAGUILA	91 RVUE	
>382	90	⁷¹ POLAK	91 RVUE	Electroweak
[> 2000]		WALKER	91 COSM	Nucleosynthesis; light ν_R
[> 500]		⁷² GRIFOLS	90 ASTR	SN 1987A; light ν_R
(> 460)	90	⁷³ HE	90B RVUE	
[> 2400–6800]		⁷⁴ BARBIERI	89B ASTR	SN 1987A; light ν_R
>189		⁷⁵ DELAGUILA	89 RVUE	$\rho\bar{p}$
[> 10000]		RAFFELT	88 ASTR	SN 1987A; light ν_R
>325	90	⁷⁶ AMALDI	87 RVUE	
>278	90	⁷⁷ DURKIN	86 RVUE	
>150	95	⁷⁸ ADEVA	85B MRKJ	$e^+e^- \rightarrow \mu^+\mu^-$

⁵³ ABE 97S limit is obtained assuming that Z' decays to known fermions only.

⁵⁴ LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

⁵⁵ LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0025 < \theta < 0.0083$.

⁵⁶ BARATE 97B gives 95% CL limits on Z - Z' mixing $-0.0017 < \theta < 0.0035$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

⁵⁷ ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

⁵⁸ VILAIN 94B assume $m_t = 150$ GeV and $\theta=0$. See Fig.2 for limit contours in the mass-mixing plane.

⁵⁹ ADRIANI 93D give limits on the Z - Z' mixing $-0.002 < \theta < 0.015$ assuming the ABE 92B mass limit.

- ⁶⁰ ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z - Z' mixing angle is in Table 4.
- ⁶¹ RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.
- ⁶² These limits assume that Z' decays to known fermions only.
- ⁶³ These limits assume that Z' decays to all E_6 fermions and their superpartners.
- ⁶⁴ See Fig. 7b and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.
- ⁶⁵ LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.
- ⁶⁶ POLAK 92 limit is from $m_{W_R} > 477$ GeV, which is derived from muon decay parameters assuming light ν_R . Specific Higgs sector is assumed.
- ⁶⁷ POLAK 92B limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using Z parameters, m_W , and low-energy neutral current data as of 1991. Light ν_R assumed and $m_t = m_H = 100$ GeV used. Supersedes POLAK 91.
- ⁶⁸ RENTON 92 limits use LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. Specific Higgs structure is assumed.
- ⁶⁹ ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.
- ⁷⁰ DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, $m_t > 77$ GeV, $m_{H^0} < 1$ TeV assumed.
- ⁷¹ POLAK 91 limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using m_W , m_Z , and low-energy neutral current data as of 1990. Light ν_R assumed and $m_t = m_H = 100$ GeV used. Superseded by POLAK 92B.
- ⁷² GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.
- ⁷³ HE 90B model assumes a specific Higgs sector. Neutral current data of COSTA 88 as well as m_Z is used. g_R is left free in the fit.
- ⁷⁴ BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- ⁷⁵ DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+ e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.
- ⁷⁶ A wide range of neutral current data as of 1986 are used in the fit.
- ⁷⁷ A wide range of neutral current data as of 1985 are used in the fit.
- ⁷⁸ ADEVA 85B measure asymmetry of μ -pair production, following formalism of RIZZO 81.

Limits for Z_χ

Z_χ is the extra neutral boson in $SO(10) \rightarrow SU(5) \times U(1)_\chi$. $g_\chi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>595	95	⁷⁹ ABE	97S CDF	$p\bar{p}; Z'_\chi \rightarrow e^+ e^-, \mu^+ \mu^-$
>321	95	^{80,81} LANGACKER	92B RVUE	Electroweak

••• We do not use the following data for averages, fits, limits, etc. •••

>190	95	82 ARIMA	97 VNS	Bhabha scattering
>236	95	83 BARATE	97B ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>196	95	84 BUSKULIC	96N ALEP	Hadronic cross section
>425	95	85 ABE	95 CDF	$\rho\bar{p}; Z'\chi \rightarrow e^+e^-$
>147	95	86 ABREU	95M DLPH	Z parameters and $e^+e^- \rightarrow \mu^+\mu^- (n\gamma)$
		87 NARDI	95 RVUE	Z parameters
		88 BUSKULIC	94 ALEP	Z parameters
>262	95	89 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>117	95	90 ADRIANI	93D L3	Z parameters
(>900)	90	91 ALTARELLI	93B RVUE	Z parameters
>340	95	92 ABE	92B CDF	$\rho\bar{p}$
>280	95	93 ABE	92B CDF	$\rho\bar{p}$
(>650)	90	94 DELAGUILA	92 RVUE	
(>760)		95 LAYSSAC	92B RVUE	Z parameters
>148	95	96 LEIKE	92 RVUE	Z parameters
(>700)		97 RENTON	92 RVUE	
(> 500)	90	98 ALTARELLI	91B RVUE	Z parameters
(> 570)		99 BUCHMUEL...	91 RVUE	Z parameters
(> 555)	90	100 DELAGUILA	91 RVUE	
[>1470]		101 FARAGGI	91 COSM	Nucleosynthesis; light ν_R
>320	90	102 GONZALEZ-G..	91 RVUE	
>221		103 MAHANTHAP..	91 RVUE	Cs
>231	90	104,105 ABE	90F VNS	e^+e^-
>206	90	105,106 ABE	90F RVUE	$e^+e^-, \nu_\mu e$
>335		107 BARGER	90B RVUE	$\rho\bar{p}$
(> 650)	90	108 GLASHOW	90 RVUE	
[> 1140]		109 GONZALEZ-G..	90D COSM	Nucleosynthesis; light ν_R
[> 2100]		110 GRIFOLS	90 ASTR	SN 1987A; light ν_R
none <150 or > 363	90	111 HAGIWARA	90 RVUE	e^+e^-
>177		112 DELAGUILA	89 RVUE	$\rho\bar{p}$
>280	95	113 DORENBOS...	89 CHRM	$g_\chi = g_Z$
>352	90	114 COSTA	88 RVUE	
>170	90	115 ELLIS	88 RVUE	$\rho\bar{p}$
>273	90	114 AMALDI	87 RVUE	
>266	90	116 MARCIANO	87 RVUE	
>283	90	117 DURKIN	86 RVUE	

⁷⁹ ABE 97S limit is obtained assuming that Z' decays to known fermions only.

⁸⁰ LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

⁸¹ LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0048 < \theta < 0.0097$.

⁸² Z - Z' mixing is assumed to be zero.

⁸³ BARATE 97B gives 95% CL limits on Z - Z' mixing $-0.0016 < \theta < 0.0036$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-90}$ GeV.

See their Fig. 4 for the limit contour in the mass-mixing plane.

⁸⁴ BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}=130, 136$ GeV (ALEPH) and $\sqrt{s}=58$ GeV (TOPAZ). Zero mixing is assumed.

- 85 ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.
- 86 ABREU 95M limit is for $\alpha_s=0.123$, $m_t=150$ GeV, and $m_H=300$ GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.
- 87 NARDI 95 give 90%CL limits on Z - Z' mixing $-0.0032 < \theta < 0.0031$ for $M_{Z'} > 500$ GeV, $m_t=170$ GeV, $m_H=250$ GeV, $\alpha_s=0.12$. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0032 < \theta < 0.0079$.
- 88 BUSKULIC 94 give 95%CL limits on the Z - Z' mixing $-0.0091 < \theta < 0.0023$.
- 89 VILAIN 94B assume $m_t = 150$ GeV and $\theta=0$. See Fig. 2 for limit contours in the mass-mixing plane.
- 90 ADRIANI 93D give limits on the Z - Z' mixing $-0.004 < \theta < 0.015$ assuming the ABE 92B mass limit.
- 91 ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z - Z' mixing angle is in their Fig. 2.
- 92 These limits assume that Z' decays to known fermions only.
- 93 These limits assume that Z' decays to all E_6 fermions and their superpartners.
- 94 See Fig. 7a and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.
- 95 LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.
- 96 LEIKE 92 is based on '90 LEP data published in LEP 92.
- 97 RENTON 92 limits use LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. Specific Higgs structure is assumed.
- 98 ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.
- 99 BUCHMUELLER 91 limit is from LEP data. Specific assumption is made for the Higgs sector.
- 100 DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, $m_t > 77$ GeV, $m_{H^0} < 1$ TeV assumed.
- 101 FARAGGI 91 limit assumes the nucleosynthesis bound on the effective number of neutrinos $\Delta N_\nu < 0.5$ and is valid for $m_{\nu_R} < 1$ MeV.
- 102 GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, Z mass and widths, m_W from ABE 90G. $100 < m_t < 200$ GeV, $m_{H^0} = 100$ GeV assumed. Dependence on m_t is shown in Fig. 7.
- 103 MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W , m_Z .
- 104 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$.
- 105 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 106 e^+e^- data for R , $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
- 107 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
- 108 GLASHOW 90 model assumes a specific Higgs sector. See GLASHOW 90B.
- 109 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
- 110 GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.

- 111 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries. The upper mass limit disappears at 2.7 s.d.
- 112 DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.
- 113 DORENBOSCH 89 obtain the limit $(g_\chi/g_Z)^2 \cdot (m_Z/m_{Z_\chi})^2 < 0.11$ at 95% CL from the processes $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$.
- 114 A wide range of neutral current data as of 1986 are used in the fit.
- 115 Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.
- 116 MARCIANO 87 limit from unitarity of Cabibbo-Kobayashi-Maskawa matrix.
- 117 A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_ψ

Z_ψ is the extra neutral boson in $E_6 \rightarrow SO(10) \times U(1)_\psi$. $g_\psi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>590	95	118 ABE	97S CDF	$p\bar{p}; Z'_\psi \rightarrow e^+e^-, \mu^+\mu^-$
>160	95	119,120 LANGACKER	92B RVUE	Electroweak
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>160	95	121 BARATE	97B ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>148	95	122 BUSKULIC	96N ALEP	Hadronic cross section
>415	95	123 ABE	95 CDF	$p\bar{p}; Z'_\psi \rightarrow e^+e^-$
>105	95	124 ABREU	95M DLPH	Z parameters and $e^+e^- \rightarrow \mu^+\mu^- (n\gamma)$
		125 NARDI	95 RVUE	Z parameters
>135	95	126 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>118	95	127 ADRIANI	93D L3	Z parameters
>320	95	128 ABE	92B CDF	$p\bar{p}$
>180	95	129 ABE	92B CDF	$p\bar{p}$
>122	95	130 LEIKE	92 RVUE	Z parameters
>105	90	131,132 ABE	90F VNS	e^+e^-
>146	90	132,133 ABE	90F RVUE	$e^+e^-, \nu_\mu e$
>320		134 BARGER	90B RVUE	$p\bar{p}$
[> 160]		135 GONZALEZ-G.	90D COSM	Nucleosynthesis; light ν_R
[> 2000]		136 GRIFOLS	90D ASTR	SN 1987A; light ν_R
>136	90	137 HAGIWARA	90 RVUE	e^+e^-
>154	90	138 AMALDI	87 RVUE	
>146	90	139 DURKIN	86 RVUE	

- 118 ABE 97S limit is obtained assuming that Z' decays to known fermions only.
- 119 LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.
- 120 LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0025 < \theta < 0.013$.

- 121 BARATE 97B gives 95% CL limits on Z - Z' mixing $-0.0020 < \theta < 0.0038$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150_{-90}^{+120}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.
- 122 BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}=130, 136$ GeV (ALEPH) and $\sqrt{s}=58$ GeV (TOPAZ). Zero mixing is assumed.
- 123 ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.
- 124 ABREU 95M limit is for $\alpha_s=0.123$, $m_t=150$ GeV, and $m_H=300$ GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.
- 125 NARDI 95 give 90%CL limits on Z - Z' mixing $-0.0056 < \theta < 0.0055$ for $M_{Z'} > 500$ GeV, $m_t=170$ GeV, $m_H=250$ GeV, $\alpha_s=0.12$. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0066 < \theta < 0.0071$.
- 126 VILAIN 94B assume $m_t = 150$ GeV and $\theta=0$. See Fig.2 for limit contours in the mass-mixing plane.
- 127 ADRIANI 93D give limits on the Z - Z' mixing $-0.003 < \theta < 0.020$ assuming the ABE 92B mass limit.
- 128 These limits assume that Z' decays to known fermions only.
- 129 These limits assume that Z' decays to all E_6 fermions and their superpartners.
- 130 LEIKE 92 is based on '90 LEP data published in LEP 92.
- 131 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$.
- 132 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 133 e^+e^- data for R , $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
- 134 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
- 135 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
- 136 GRIFOLS 90D limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also RIZZO 91.
- 137 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries.
- 138 A wide range of neutral current data as of 1986 are used in the fit.
- 139 A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_η

Z_η is the extra neutral boson in E_6 models, corresponding to $Q_\eta = \sqrt{3/8} Q_\chi - \sqrt{5/8} Q_\psi$. $g_\eta = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho=1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>620	95	140 ABE	97S CDF	$p\bar{p}; Z'_\eta \rightarrow e^+e^-, \mu^+\mu^-$
>182	95	141,142 LANGACKER	92B RVUE	Electroweak

••• We do not use the following data for averages, fits, limits, etc. •••

>173	95	143 BARATE	97B ALEP	$e^+e^- \rightarrow \mu^+\mu^-$ and hadronic cross section
>167	95	144 BUSKULIC	96N ALEP	Hadronic cross section
>440	95	145 ABE	95 CDF	$\rho\bar{p}; Z'_\eta \rightarrow e^+e^-$
>109	95	146 ABREU	95M DLPH	Z parameters and $e^+e^- \rightarrow \mu^+\mu^- (n\gamma)$
		147 NARDI	95 RVUE	Z parameters
>100	95	148 VILAIN	94B CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>100	95	149 ADRIANI	93D L3	Z parameters
(>500)	90	150 ALTARELLI	93B RVUE	Z parameters
>340	95	151 ABE	92B CDF	$\rho\bar{p}$
>230	95	152 ABE	92B CDF	$\rho\bar{p}$
(>450)	90	153 DELAGUILA	92 RVUE	
(>315)		154 LAYSSAC	92B RVUE	Z parameters
>118	95	155 LEIKE	92 RVUE	Z parameters
(>470)		156 RENTON	92 RVUE	
(> 300)	90	157 ALTARELLI	91B RVUE	Z parameters
>120	90	158 GONZALEZ-G.	91 RVUE	
>125	90	159,160 ABE	90F VNS	e^+e^-
>115	90	160,161 ABE	90F RVUE	$e^+e^-, \nu_\mu e$
>340		162 BARGER	90B RVUE	$\rho\bar{p}$
[> 820]		163 GONZALEZ-G.	90D COSM	Nucleosynthesis; light ν_R
[> 3300]		164 GRIFOLS	90 ASTR	SN 1987A; light ν_R
>100	90	165 HAGIWARA	90 RVUE	e^+e^-
[> 1040]		163 LOPEZ	90 COSM	Nucleosynthesis; light ν_R
>173		166 DELAGUILA	89 RVUE	$\rho\bar{p}$
>129	90	167 COSTA	88 RVUE	
>156	90	168 ELLIS	88 RVUE	
>167	90	169 ELLIS	88 RVUE	$\rho\bar{p}$
>111	90	167 AMALDI	87 RVUE	
>143	90	170 BARGER	86B RVUE	$\rho\bar{p}$
>130	90	171 DURKIN	86 RVUE	
[> 760]		163 ELLIS	86 COSM	Nucleosynthesis; light ν_R
[> 500]		163 STEIGMAN	86 COSM	Nucleosynthesis; light ν_R

140 ABE 97S limit is obtained assuming that Z' decays to known fermions only.

141 LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

142 LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.038 < \theta < 0.002$.

143 BARATE 97B gives 95% CL limits on Z - Z' mixing $-0.021 < \theta < 0.012$. The bounds are computed with $\alpha_s = 0.120 \pm 0.003$, $m_t = 175 \pm 6$ GeV, and $M_H = 150^{+120}_{-90}$ GeV. See their Fig. 4 for the limit contour in the mass-mixing plane.

144 BUSKULIC 96N limit is from a combined fit to the hadronic cross sections measured at $\sqrt{s}=130, 136$ GeV (ALEPH) and $\sqrt{s}=58$ GeV (TOPAZ). Zero mixing is assumed.

145 ABE 95 limit is obtained assuming that Z' decays to known fermions only. See their Fig. 3 for the mass bound of Z' decaying to all allowed fermions and supersymmetric fermions.

146 ABREU 95M limit is for $\alpha_s=0.123$, $m_t=150$ GeV, and $m_H=300$ GeV. For the limit contour in the mass-mixing plane, see their Fig. 13.

- 147 NARDI 95 give 90%CL limits on Z - Z' mixing $-0.0087 < \theta < 0.0075$ for $M_{Z'} > 500$ GeV, $m_t = 170$ GeV, $m_H = 250$ GeV, $\alpha_s = 0.12$. The bound is relaxed under the simultaneous presence of the mixing of the known fermions with new heavy states, $-0.0087 < \theta < 0.010$.
- 148 VILAIN 94B assume $m_t = 150$ GeV and $\theta = 0$. See Fig. 2 for limit contours in the mass-mixing plane.
- 149 ADRIANI 93D give limits on the Z - Z' mixing $-0.029 < \theta < 0.010$ assuming the ABE 92B mass limit.
- 150 ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The 90%CL limit on the Z - Z' mixing angle is in Fig. 2.
- 151 These limits assume that Z' decays to known fermions only.
- 152 These limits assume that Z' decays to all E_6 fermions and their superpartners.
- 153 See Fig. 7d in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane from electroweak fit including '90 LEP data.
- 154 LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.
- 155 LEIKE 92 is based on '90 LEP data published in LEP 92.
- 156 RENTON 92 limits use LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. Specific Higgs structure is assumed.
- 157 ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.
- 158 GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, LEP Z mass and widths, m_W from ABE 90G. $100 < m_t < 200$ GeV, $m_{H^0} = 100$ GeV assumed. Dependence on m_t is shown in Fig. 8.
- 159 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$.
- 160 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 161 e^+e^- data for R , $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
- 162 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
- 163 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
- 164 GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.
- 165 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries.
- 166 DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.
- 167 A wide range of neutral current data as of 1986 are used in the fit.
- 168 Z_η mass limits obtained by combining constraints from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider and the global analysis of neutral current data by COSTA 88. Least favorable spectrum of three (E_6 27) generations of particles and their superpartners are assumed.
- 169 Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.
- 170 BARGER 86B limit is based on UA1/UA2 limit on $p\bar{p} \rightarrow Z'$, $Z' \rightarrow e^+e^-$ (Lepton Photon Symp., Kyoto, '85). Extra decay channels for Z' are assumed not be open.
- 171 A wide range of neutral current data as of 1985 are used in the fit.

Limits for other Z'

$$Z_\beta = Z_\chi \cos\beta + Z_\psi \sin\beta$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		172 DELAGUILA	92	RVUE
>360		173 ALTARELLI	91	RVUE Z_β with $\tan\beta = \sqrt{3/5}$;
>190		174 MAHANTHAP.	91	RVUE Z_β with $\tan\beta = \sqrt{3/5}$;
		175 GRIFOLS	90C	RVUE Cs
		176 DELAGUILA	89	RVUE $p\bar{p}$
>180	90	177,178 COSTA	88	RVUE Z_β with $\tan\beta = \sqrt{15}$
>158	90	179 ELLIS	88	RVUE Z_β ($\tan\beta = \sqrt{15}$), $p\bar{p}$

172 Fig. 7c and 7e in DELAGUILA 92 give limits for $\tan\beta = -1/\sqrt{15}$ and $\sqrt{15}$ from electroweak fit including '90 LEP data.

173 ALTARELLI 91 limit is from atomic parity violation in Cs together with LEP, CDF data. Z - Z' mixing is assumed to be zero to set the limit.

174 MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W , m_Z . See Table III of MAHANTHAPPA 91 (corrected in erratum) for limits on various Z' models.

175 GRIFOLS 90C obtains a limit for Z' mass as a function of mixing angle β (his $\theta = \beta - \pi/2$), which is derived from a LAMPF experiment on $\sigma(\nu_e e)$ (ALLEN 90). The result is shown in Fig. 1.

176 See Table I of DELAGUILA 89 for limits on various Z' models.

177 $g_\beta = e/\cos\theta_W$ and $\rho = 1$ assumed.

178 A wide range of neutral current data as of 1986 are used in the fit.

179 Z' mass limits from non-observation of an excess of $\ell^+ \ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.

LEPTOQUARK QUANTUM NUMBERS

Written December 1997 by M. Tanabashi (Tohoku U.).

Leptoquarks are particles carrying both baryon number (B) and lepton number (L). They are expected to exist in various extensions of the Standard Model (SM). The possible quantum numbers of leptoquark states can be restricted by assuming that their direct interactions with the ordinary SM fermions are dimensionless and invariant under the SM gauge group. Table 1 shows the list of all possible quantum numbers with this assumption [1]. The columns of $SU(3)_C$, $SU(2)_W$, and $U(1)_Y$ in Table 1 indicate the QCD representation, the weak isospin representation, and the weak hypercharge, respectively. Naming conventions of leptoquark states are taken from Ref. 1. The spin

of a leptoquark state is taken to be 1 (vector leptoquark) or 0 (scalar leptoquark).

Table 1: Possible leptoquarks and their quantum numbers.

Leptoquarks	Spin	$3B + L$	$SU(3)_c$	$SU(2)_W$	$U(1)_Y$
S_1	0	-2	$\bar{3}$	1	1/3
\tilde{S}_1	0	-2	$\bar{3}$	1	4/3
S_3	0	-2	$\bar{3}$	3	1/3
V_2	1	-2	$\bar{3}$	2	5/6
\tilde{V}_2	1	-2	$\bar{3}$	2	-1/6
R_2	0	0	3	2	7/6
\tilde{R}_2	0	0	3	2	1/6
U_1	1	0	3	1	2/3
\tilde{U}_1	1	0	3	1	5/3
U_3	1	0	3	3	2/3

If we do not require leptoquark states to couple directly with SM fermions, different assignments of quantum numbers become possible.

The Pati-Salam model [2] is an example predicting the existence of a leptoquark state. In this model a vector leptoquark appears at the scale where the Pati-Salam $SU(4)$ “color” gauge group breaks into the familiar QCD $SU(3)_C$ group (or $SU(3)_C \times U(1)_{B-L}$). The Pati-Salam leptoquark is a weak isosinglet and its hypercharge is 2/3 (U_1 leptoquark in Table 1). The coupling strength of the Pati-Salam leptoquark is given by the QCD coupling at the Pati-Salam symmetry breaking scale.

Bounds on leptoquark states are obtained both directly and indirectly. Direct limits are from their production cross sections at colliders, while indirect limits are calculated from the bounds

on the leptoquark induced four-fermion interactions which are obtained from low energy experiments.

The pair production cross sections of leptoquarks are evaluated from their interactions with gauge bosons. The gauge couplings of a scalar leptoquark are determined uniquely according to its quantum numbers in Table 1. The magnetic-dipole-type and the electric-quadrupole-type interactions of a vector leptoquark are, however, not determined even if we fix its gauge quantum numbers as listed in the table [3]. We need extra assumptions about these interactions to evaluate the pair production cross section for a vector leptoquark.

If a leptoquark couples to fermions of more than a single generation in the mass eigenbasis of the SM fermions, it can induce four-fermion interactions causing flavor-changing-neutral-currents and lepton-family-number violations. Non-chiral leptoquarks, which couple simultaneously to both left- and right-handed quarks, cause four-fermion interactions affecting the $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$ ratio [4]. Indirect limits provide stringent constraints on these leptoquarks. Since the Pati-Salam leptoquark has non-chiral coupling with both e and μ , indirect limits from the bounds on $K_L \rightarrow \mu e$ lead to severe bounds on the Pati-Salam leptoquark mass. For detailed bounds obtained in this way, see the Boson Particle Listings for “Indirect Limits for Leptoquarks” and its references.

It is therefore often assumed that a leptoquark state couples only to a single generation in a chiral interaction, where indirect limits become much weaker. This assumption gives strong constraints on concrete models of leptoquarks, however. Leptoquark states which couple only to left- or right-handed quarks are called chiral leptoquarks. Leptoquark states which couple only to the first (second, third) generation are referred as the first (second, third) generation leptoquarks in this section.

Reference

1. W. Buchmüller, R. Rückl, and D. Wyler, Phys. Lett. **B191**, 442 (1987).
2. J.C. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974).
3. J. Blümlein, E. Boos, and A. Kryukov, Z. Phys. **C76**, 137 (1997).
4. O. Shanker, Nucl. Phys. **B204**, 375 (1982).

MASS LIMITS for Leptoquarks from Pair Production

These limits rely only on the color or electroweak charge of the leptoquark.

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
>225	95	180	ABBOTT	98E D0	First generation
> 99	95	181	ABE	97F CDF	Third generation
>131	95	182	ABE	95U CDF	Second generation
> 45.5	95	183,184	ABREU	93J DLPH	First + second generation
> 44.4	95	185	ADRIANI	93M L3	First generation
> 44.6	95	186	ADRIANI	93M L3	Third generation
> 44	95	185	DECAMP	92 ALEP	First or second generation
> 45	95	185	DECAMP	92 ALEP	Third generation
> 44.2	95	185	ALEXANDER	91 OPAL	First or second generation
> 41.4	95	185	ALEXANDER	91 OPAL	Third generation
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>225	95	187	ABBOTT	97B D0	Result included in ABBOTT 98E
>213	95	187	ABE	97X CDF	First generation
>119	95	188	ABACHI	95G D0	Second generation
>116	95	189	ABACHI	94B D0	First generation
> 80	95	190	ABE	93I CDF	First generation
> 44.5	95	185	ADRIANI	93M L3	Second generation
> 42.1	95	191	ABREU	92F DLPH	Second generation
> 74	95	192	ALITTI	92E UA2	First generation
> 43.2	95	185	ADEVA	91B L3	First generation
> 43.4	95	185	ADEVA	91B L3	Second generation
none 8.9–22.6	95	193	KIM	90 AMY	First generation
none 10.2–23.2	95	193	KIM	90 AMY	Second generation
none 5–20.8	95	194	BARTEL	87B JADE	
none 7–20.5	95	2	195 BEHREND	86B CELL	

180 ABBOTT 98E search for scalar leptoquarks using $e\nu jj$, $eejj$, and $\nu\nu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit above assumes $B(eq)=1$. For $B(eq)=0.5$ and 0, the bound becomes 204 and 79 GeV, respectively.

181 ABE 97F search for third generation scalar and vector leptoquarks in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The quoted limit is for scalar leptoquark with $B(\tau b) = 1$.

182 ABE 95U search for scalar leptoquarks of charge $Q=2/3$ and $-1/3$ using $\mu\mu jj$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit is for $B(\mu q) = 1$. For $B(\mu q) = B(\nu q) = 0.5$, the limit is > 96 GeV.

183 Limit is for charge $-1/3$ isospin-0 leptoquark with $B(\ell q) = 2/3$.

- 184 First and second generation leptoquarks are assumed to be degenerate. The limit is slightly lower for each generation.
- 185 Limits are for charge $-1/3$, isospin-0 scalar leptoquarks decaying to $\ell^- q$ or νq with any branching ratio. See paper for limits for other charge-isospin assignments of leptoquarks.
- 186 ADRIANI 93M limit for charge $-1/3$, isospin-0 leptoquark decaying to τb .
- 187 ABBOTT 97B, ABE 97X search for scalar leptoquarks using $e e j j$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit is for $B(eq)=1$.
- 188 ABACHI 95G search for scalar leptoquarks using $\mu\mu$ +jets and $\mu\nu_\mu$ +jets events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. The limit is for $B(\mu q) = 1$. For $B(\mu q) = B(\nu q) = 0.5$, the limit is > 97 GeV.
- 189 ABACHI 94B search for $e e j j$ and $e \nu j j$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. ABACHI 94B obtain the limit >120 GeV for $B(eq)=B(\nu q)=0.5$ and >133 GeV for $B(eq)=1$. A change in the $D\bar{D}$ luminosity monitor constant reduces the first bound to >116 GeV quoted above (see FERMILAB-TM-1911). This limit does not depend on the electroweak quantum numbers of the leptoquark.
- 190 ABE 93I search for $\ell\ell j j$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=1.8$ TeV. The limit is for $B(eq) = B(\nu q) = 0.5$ and improves to >113 GeV for $B(eq) = 1$. This limit does not depend on electroweak quantum numbers of the leptoquark.
- 191 ABREU 92F limit is for charge $-1/3$ isospin-0 leptoquark with $B(\mu q)=2/3$. If first and second generation leptoquarks are degenerate, the limit is 43.0 GeV, and for a charge $2/3$ second generation leptoquark 43.4 GeV. Cross-section limit for pair production of states decaying to ℓq is given in the paper.
- 192 ALITTI 92E search for $\ell\ell j j$ and $\ell\nu j j$ events in $p\bar{p}$ collisions at $E_{\text{cm}}=630$ GeV. The limit is for $B(eq) = 1$ and is reduced to 67 GeV for $B(eq) = B(\nu q) = 0.5$. This limit does not depend on electroweak quantum numbers of the leptoquark.
- 193 KIM 90 assume pair production of charge $2/3$ scalar-leptoquark via photon exchange. The decay of the first (second) generation leptoquark is assumed to be any mixture of $d e^+$ and $u\bar{\nu}$ ($s\mu^+$ and $c\bar{\nu}$). See paper for limits for specific branching ratios.
- 194 BARTEL 87B limit is valid when a pair of charge $2/3$ spinless leptoquarks X is produced with point coupling, and when they decay under the constraint $B(X \rightarrow c\bar{\nu}_\mu) + B(X \rightarrow s\mu^+) = 1$.
- 195 BEHREND 86B assumed that a charge $2/3$ spinless leptoquark, χ , decays either into $s\mu^+$ or $c\bar{\nu}$: $B(\chi \rightarrow s\mu^+) + B(\chi \rightarrow c\bar{\nu}) = 1$.

MASS LIMITS for Leptoquarks from Single Production

These limits depend on the q - ℓ -leptoquark coupling g_{LQ} . It is often assumed that $g_{LQ}^2/4\pi=1/137$. Limits shown are for a scalar, weak isoscalar, charge $-1/3$ leptoquark.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>237	95	196 AID	96B H1	First generation
> 73	95	197 ABREU	93J DLPH	Second generation
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		198 DERRICK	97 ZEUS	Lepton-flavor violation
>230	95	199 AHMED	94B H1	Sup. by AID 96B
> 65	95	197 ABREU	93J DLPH	First generation
>181	95	200 ABT	93 H1	First generation
>168	95	201 DERRICK	93 ZEUS	First generation

- 196 The quoted limit is for a left-handed scalar leptoquark which solely couples to the first generation with electromagnetic strength. AID 96B also search for leptoquarks with lepton-flavor violating couplings. For limits on states with different quantum numbers and the limits in the coupling-mass plane, see their Fig. 2, Fig. 3, and Table 2. AID 96B supersedes AHMED 94B.

- 197 Limit from single production in Z decay. The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(\ell q) = 2/3$. The limit is 77 GeV if first and second leptoquarks are degenerate.
- 198 DERRICK 97 search for various leptoquarks with lepton-flavor violating couplings. See their Figs. 5–8 and Table 1 for detailed limits.
- 199 AHMED 94B limit is for the left-handed leptoquark decaying to eq and νq with $B(eq) = B(\nu q) = 1/2$. Electromagnetic coupling strength is assumed for the scalar leptoquark interaction. For limits on states with different quantum numbers and the limits in the coupling-mass plane, see their Table 2 and Fig. 6.
- 200 ABT 93 search for single leptoquark production in ep collisions with the decays eq and νq . The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 178 GeV. For limits on states with different quantum numbers, see their Fig. 2. ABT 93 superseded by AHMED 94B.
- 201 DERRICK 93 search for single leptoquark production in ep collisions with the decay eq and νq . The limit is for leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 176 GeV. For limits on states with different quantum numbers, see their Table 3.

Indirect Limits for Leptoquarks

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 0.76	95	202 DEANDREA	97 RVUE	\tilde{R}_2 leptoquark
		203 DERRICK	97 ZEUS	Lepton-flavor violation
		204 GROSSMAN	97 RVUE	$B \rightarrow \tau^+ \tau^- (X)$
		205 JADACH	97 RVUE	$e^+ e^- \rightarrow q \bar{q}$
> 0.31	95	206 AID	95 H1	First generation
>1200		207 KUZNETSOV	95B RVUE	Pati-Salam type
		208 MIZUKOSHI	95 RVUE	Third generation scalar leptoquark
> 0.3	95	209 BHATTACH...	94 RVUE	Spin-0 leptoquark coupled to $\bar{e}_R t_L$
		210 DAVIDSON	94 RVUE	
> 18		211 KUZNETSOV	94 RVUE	Pati-Salam type
> 0.43	95	212 LEURER	94 RVUE	First generation spin-1 leptoquark
> 0.44	95	212 LEURER	94B RVUE	First generation spin-0 leptoquark
		213 MAHANTA	94 RVUE	P and T violation
> 350		214 DESHPANDE	83 RVUE	Sup. by KUZNETSOV 95B
> 1		215 SHANKER	82 RVUE	Nonchiral spin-0 leptoquark
> 125		215 SHANKER	82 RVUE	Nonchiral spin-1 leptoquark

- 202 DEANDREA 97 limit is for \tilde{R}_2 leptoquark obtained from atomic parity violation (APV). The coupling of leptoquark is assumed to be electromagnetic strength. See Table 2 for limits of the four-fermion interactions induced by various scalar leptoquark exchange. DEANDREA 97 combines APV limit and limits from Tevatron and HERA. See Fig. 1–4 for combined limits of leptoquark in mass-coupling plane.
- 203 DERRICK 97 search for lepton-flavor violation in ep collision. See their Tables 2–5 for limits on lepton-flavor violating four-fermion interactions induced by various leptoquarks.
- 204 GROSSMAN 97 estimate the upper bounds on the branching fraction $B \rightarrow \tau^+ \tau^- (X)$ from the absence of the B decay with large missing energy. These bounds can be used to constrain leptoquark induced four-fermion interactions.

- 205 JADACH 97 limit is from $e^+e^- \rightarrow q\bar{q}$ cross section at $\sqrt{s}=172.3$ GeV which can be affected by the t - and u -channel exchanges of leptoquarks. See their Fig. 1 for limits on vector leptoquarks in mass-coupling plane.
- 206 AID 95 limit is for the weak isotriplet spin-1 leptoquark with the electromagnetic coupling strength. For the limits of leptoquarks with different quantum number, see their Table 2. AID 95 limits are from the measurements of the Q^2 spectrum measurement of $e p \rightarrow e X$.
- 207 KUZNETSOV 95B use π , K , B , τ decays and μe conversion and give a list of bounds on the leptoquark mass and the fermion mixing matrix in the Pati-Salam model. The quoted limit is from $K_L \rightarrow \mu e$ decay assuming zero mixing. See also KUZNETSOV 94, DESHPANDE 83, and DIMOPOULOS 81.
- 208 MIZUKOSHI 95 calculate the one-loop radiative correction to the Z -physics parameters in various scalar leptoquark models. See their Fig. 4 for the exclusion plot of third generation leptoquark models in mass-coupling plane.
- 209 BHATTACHARYYA 94 limit is from one-loop radiative correction to the leptonic decay width of the Z . $m_H=250$ GeV, $\alpha_s(m_Z)=0.12$, $m_t=180$ GeV, and the electroweak strength of leptoquark coupling are assumed. For leptoquark coupled to $\bar{e}_L t_R$, $\bar{\mu} t$, and $\bar{\tau} t$, see Fig. 2 in BHATTACHARYYA 94B erratum and Fig. 3.
- 210 DAVIDSON 94 gives an extensive list of the bounds on leptoquark-induced four-fermion interactions from π , K , D , B , μ , τ decays and meson mixings, *etc.* See Table 15 of DAVIDSON 94 for detail.
- 211 KUZNETSOV 94 gives mixing independent bound of the Pati-Salam leptoquark from the cosmological limit on $\pi^0 \rightarrow \bar{\nu}\nu$.
- 212 LEURER 94, LEURER 94B limits are obtained from atomic parity violation and apply to any chiral leptoquark which couples to the first generation with electromagnetic strength. For a nonchiral leptoquark, universality in $\pi_{\ell 2}$ decay provides a much more stringent bound. See also SHANKER 82.
- 213 MAHANTA 94 gives bounds of P - and T -violating scalar-leptoquark couplings from atomic and molecular experiments.
- 214 DESHPANDE 83 used upper limit on $K_L^0 \rightarrow \mu e$ decay with renormalization-group equations to estimate coupling at the heavy boson mass. See also DIMOPOULOS 81.
- 215 From $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$ ratio. SHANKER 82 assumes the leptoquark induced four-fermion coupling $4g^2/M^2 (\bar{\nu}_{eL} u_R) (\bar{d}_L e_R)$ with $g=0.004$ for spin-0 leptoquark and $g^2/M^2 (\bar{\nu}_{eL} \gamma_\mu u_L) (\bar{d}_R \gamma^\mu e_R)$ with $g \simeq 0.6$ for spin-1 leptoquark.

MASS LIMITS for Diquarks

<u>VALUE (GeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 290–420	95	216 ABE	97G CDF	E_6 diquark
none 15–31.7	95	217 ABREU	940 DLPH	SUSY E_6 diquark

216 ABE 97G search for new particle decaying to dijets.

217 ABREU 940 limit is from $e^+e^- \rightarrow \bar{c}\bar{s}cs$. Range extends up to 43 GeV if diquarks are degenerate in mass.

MASS LIMITS for g_A (axigluon)

Axigluons are massive color-octet gauge bosons in chiral color models and have axial-vector coupling to quarks with the same coupling strength as gluons.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 200–980	95	218 ABE	97G CDF	$p\bar{p} \rightarrow g_A X, X \rightarrow 2 \text{ jets}$
none 200–870	95	219 ABE	95N CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow q\bar{q}$
none 240–640	95	220 ABE	93G CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2\text{jets}$
>50	95	221 CUYPERS	91 RVUE	$\sigma(e^+e^- \rightarrow \text{hadrons})$
none 120–210	95	222 ABE	90H CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2\text{jets}$
>29		223 ROBINETT	89 THEO	Partial-wave unitarity
none 150–310	95	224 ALBAJAR	88B UA1	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2\text{jets}$
>20		BERGSTROM	88 RVUE	$p\bar{p} \rightarrow \Upsilon X \text{ via } g_A g$
> 9		225 CUYPERS	88 RVUE	Υ decay
>25		226 DONCHESKI	88B RVUE	Υ decay

218 ABE 97G search for new particle decaying to dijets.

219 ABE 95N assume axigluons decaying to quarks in the Standard Model only.

220 ABE 93G assume $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 10$.

221 CUYPERS 91 compare α_s measured in Υ decay and that from R at PEP/PETRA energies.

222 ABE 90H assumes $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 5$ ($\Gamma(g_A) = 0.09m_{g_A}$). For $N = 10$, the excluded region is reduced to 120–150 GeV.

223 ROBINETT 89 result demands partial-wave unitarity of $J = 0$ $t\bar{t} \rightarrow t\bar{t}$ scattering amplitude and derives a limit $m_{g_A} > 0.5 m_t$. Assumes $m_t > 56$ GeV.

224 ALBAJAR 88B result is from the nonobservation of a peak in two-jet invariant mass distribution. $\Gamma(g_A) < 0.4 m_{g_A}$ assumed. See also BAGGER 88.

225 CUYPERS 88 requires $\Gamma(\Upsilon \rightarrow g g_A) < \Gamma(\Upsilon \rightarrow g g g)$. A similar result is obtained by DONCHESKI 88.

226 DONCHESKI 88B requires $\Gamma(\Upsilon \rightarrow g q\bar{q})/\Gamma(\Upsilon \rightarrow g g g) < 0.25$, where the former decay proceeds via axigluon exchange. A more conservative estimate of < 0.5 leads to $m_{g_A} > 21$ GeV.

X^0 (Heavy Boson) Searches in Z Decays

Searches for radiative transition of Z to a lighter spin-0 state X^0 decaying to hadrons, a lepton pair, a photon pair, or invisible particles as shown in the comments. The limits are for the product of branching ratios.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		227 ACCIARRI	97Q L3	$X^0 \rightarrow \text{invisible particle(s)}$
		228 ACTON	93E OPAL	$X^0 \rightarrow \gamma\gamma$
		229 ABREU	92D DLPH	$X^0 \rightarrow \text{hadrons}$
		230 ADRIANI	92F L3	$X^0 \rightarrow \text{hadrons}$
		231 ACTON	91 OPAL	$X^0 \rightarrow \text{anything}$
$< 1.1 \times 10^{-4}$	95	232 ACTON	91B OPAL	$X^0 \rightarrow e^+e^-$

$<9 \times 10^{-5}$	95	232	ACTON	91B OPAL	$X^0 \rightarrow \mu^+ \mu^-$
$<1.1 \times 10^{-4}$	95	232	ACTON	91B OPAL	$X^0 \rightarrow \tau^+ \tau^-$
$<2.8 \times 10^{-4}$	95	233	ADEVA	91D L3	$X^0 \rightarrow e^+ e^-$
$<2.3 \times 10^{-4}$	95	233	ADEVA	91D L3	$X^0 \rightarrow \mu^+ \mu^-$
$<4.7 \times 10^{-4}$	95	234	ADEVA	91D L3	$X^0 \rightarrow \text{hadrons}$
$<8 \times 10^{-4}$	95	235	AKRAWY	90J OPAL	$X^0 \rightarrow \text{hadrons}$

227 See Fig. 4 of ACCIARRI 97Q for the upper limit on $B(Z \rightarrow \gamma X^0; E_\gamma > E_{\min})$ as a function of E_{\min} .

228 ACTON 93E give $\sigma(e^+ e^- \rightarrow X^0 \gamma) \cdot B(X^0 \rightarrow \gamma \gamma) < 0.4 \text{ pb}$ (95%CL) for $m_{X^0} = 60 \pm 2.5 \text{ GeV}$. If the process occurs via s -channel γ exchange, the limit translates to $\Gamma(X^0) \cdot B(X^0 \rightarrow \gamma \gamma)^2 < 20 \text{ MeV}$ for $m_{X^0} = 60 \pm 1 \text{ GeV}$.

229 ABREU 92D give $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (3-10) \text{ pb}$ for $m_{X^0} = 10-78 \text{ GeV}$. A very similar limit is obtained for spin-1 X^0 .

230 ADRIANI 92F search for isolated γ in hadronic Z decays. The limit $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (2-10) \text{ pb}$ (95%CL) is given for $m_{X^0} = 25-85 \text{ GeV}$.

231 ACTON 91 searches for $Z \rightarrow Z^* X^0$, $Z^* \rightarrow e^+ e^-$, $\mu^+ \mu^-$, or $\nu \bar{\nu}$. Excludes any new scalar X^0 with $m_{X^0} < 9.5 \text{ GeV}/c$ if it has the same coupling to $Z Z^*$ as the MSM Higgs boson.

232 ACTON 91B limits are for $m_{X^0} = 60-85 \text{ GeV}$.

233 ADEVA 91D limits are for $m_{X^0} = 30-89 \text{ GeV}$.

234 ADEVA 91D limits are for $m_{X^0} = 30-86 \text{ GeV}$.

235 AKRAWY 90J give $\Gamma(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < 1.9 \text{ MeV}$ (95%CL) for $m_{X^0} = 32-80 \text{ GeV}$. We divide by $\Gamma(Z) = 2.5 \text{ GeV}$ to get product of branching ratios. For nonresonant transitions, the limit is $B(Z \rightarrow \gamma q \bar{q}) < 8.2 \text{ MeV}$ assuming three-body phase space distribution.

MASS LIMITS for a Heavy Neutral Boson Coupling to $e^+ e^-$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
none 55-61		236 ODAKA	89 VNS	$\Gamma(X^0 \rightarrow e^+ e^-) \cdot B(X^0 \rightarrow \text{hadrons}) \gtrsim 0.2 \text{ MeV}$
>45	95	237 DERRICK	86 HRS	$\Gamma(X^0 \rightarrow e^+ e^-) = 6 \text{ MeV}$
>46.6	95	238 ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+ e^-) = 10 \text{ keV}$
>48	95	238 ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+ e^-) = 4 \text{ MeV}$
		239 BERGER	85B PLUT	
none 39.8-45.5		240 ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+ e^-) = 10 \text{ keV}$
>47.8	95	240 ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+ e^-) = 4 \text{ MeV}$
none 39.8-45.2		240 BEHREND	84C CELL	
>47	95	240 BEHREND	84C CELL	$\Gamma(X^0 \rightarrow e^+ e^-) = 4 \text{ MeV}$

236 ODAKA 89 looked for a narrow or wide scalar resonance in $e^+ e^- \rightarrow \text{hadrons}$ at $E_{\text{cm}} = 55.0-60.8 \text{ GeV}$.

237 DERRICK 86 found no deviation from the Standard Model Bhabha scattering at $E_{\text{cm}} = 29 \text{ GeV}$ and set limits on the possible scalar boson $e^+ e^-$ coupling. See their figure 4 for excluded region in the $\Gamma(X^0 \rightarrow e^+ e^-) - m_{X^0}$ plane. Electronic chiral invariance requires a parity doublet of X^0 , in which case the limit applies for $\Gamma(X^0 \rightarrow e^+ e^-) = 3 \text{ MeV}$.

- 238 ADEVA 85 first limit is from $2\gamma, \mu^+ \mu^-$, hadrons assuming X^0 is a scalar. Second limit is from $e^+ e^-$ channel. $E_{\text{cm}} = 40\text{--}47$ GeV. Supersedes ADEVA 84.
- 239 BERGER 85B looked for effect of spin-0 boson exchange in $e^+ e^- \rightarrow e^+ e^-$ and $\mu^+ \mu^-$ at $E_{\text{cm}} = 34.7$ GeV. See Fig. 5 for excluded region in the $m_{X^0} - \Gamma(X^0)$ plane.
- 240 ADEVA 84 and BEHREND 84C have $E_{\text{cm}} = 39.8\text{--}45.5$ GeV. MARK-J searched X^0 in $e^+ e^- \rightarrow$ hadrons, $2\gamma, \mu^+ \mu^-, e^+ e^-$ and CELLO in the same channels plus τ pair. No narrow or broad X^0 is found in the energy range. They also searched for the effect of X^0 with $m_{X^0} > E_{\text{cm}}$. The second limits are from Bhabha data and for spin-0 singlet. The same limits apply for $\Gamma(X^0 \rightarrow e^+ e^-) = 2$ MeV if X^0 is a spin-0 doublet. The second limit of BEHREND 84C was read off from their figure 2. The original papers also list limits in other channels.

Search for X^0 Resonance in $e^+ e^-$ Collisions

The limit is for $\Gamma(X^0 \rightarrow e^+ e^-) \cdot B(X^0 \rightarrow f)$, where f is the specified final state.

Spin 0 is assumed for X^0 .

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<10^3$	95	241 ABE	93C VNS	$\Gamma(ee)$
$<(0.4\text{--}10)$	95	242 ABE	93C VNS	$f = \gamma\gamma$
$<(0.3\text{--}5)$	95	243,244 ABE	93D TOPZ	$f = \gamma\gamma$
$<(2\text{--}12)$	95	243,244 ABE	93D TOPZ	$f =$ hadrons
$<(4\text{--}200)$	95	244,245 ABE	93D TOPZ	$f = ee$
$<(0.1\text{--}6)$	95	244,245 ABE	93D TOPZ	$f = \mu\mu$
$<(0.5\text{--}8)$	90	246 STERNER	93 AMY	$f = \gamma\gamma$

241 Limit is for $\Gamma(X^0 \rightarrow e^+ e^-) m_{X^0} = 56\text{--}63.5$ GeV for $\Gamma(X^0) = 0.5$ GeV.

242 Limit is for $m_{X^0} = 56\text{--}61.5$ GeV and is valid for $\Gamma(X^0) \ll 100$ MeV. See their Fig. 5 for limits for $\Gamma = 1, 2$ GeV.

243 Limit is for $m_{X^0} = 57.2\text{--}60$ GeV.

244 Limit is valid for $\Gamma(X^0) \ll 100$ MeV. See paper for limits for $\Gamma = 1$ GeV and those for $J = 2$ resonances.

245 Limit is for $m_{X^0} = 56.6\text{--}60$ GeV.

246 STERNER 93 limit is for $m_{X^0} = 57\text{--}59.6$ GeV and is valid for $\Gamma(X^0) < 100$ MeV. See their Fig. 2 for limits for $\Gamma = 1, 3$ GeV.

Search for X^0 Resonance in Two-Photon Process

The limit is for $\Gamma(X^0) \cdot B(X^0 \rightarrow \gamma\gamma)^2$. Spin 0 is assumed for X^0 .

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<2.6	95	247 ACTON	93E OPAL	$m_{X^0} = 60 \pm 1$ GeV
<2.9	95	BUSKULIC	93F ALEP	$m_{X^0} \sim 60$ GeV

247 ACTON 93E limit for a $J = 2$ resonance is 0.8 MeV.

Search for X^0 Resonance in $e^+ e^- \rightarrow X^0 \gamma$

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

248 ADAM 96C DLPH X^0 decaying invisibly

248 ADAM 96C is from the single photon production cross at $\sqrt{s}=130, 136$ GeV. The upper bound is less than 3 pb for X^0 masses between 60 and 130 GeV. See their Fig. 5 for the exact bound on the cross section $\sigma(e^+ e^- \rightarrow \gamma X^0)$.

Search for X^0 Resonance in $Z \rightarrow f\bar{f}X^0$

The limit is for $B(Z \rightarrow f\bar{f}X^0) \cdot B(X^0 \rightarrow F)$ where f is a fermion and F is the specified final state. Spin 0 is assumed for X^0 .

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		249 ABREU	96T DLPH	$f=e,\mu,\tau; F=\gamma\gamma$
$<3.7 \times 10^{-6}$	95	250 ABREU	96T DLPH	$f=\nu; F=\gamma\gamma$
		251 ABREU	96T DLPH	$f=q; F=\gamma\gamma$
$<6.8 \times 10^{-6}$	95	250 ACTON	93E OPAL	$f=e,\mu,\tau; F=\gamma\gamma$
$<5.5 \times 10^{-6}$	95	250 ACTON	93E OPAL	$f=q; F=\gamma\gamma$
$<3.1 \times 10^{-6}$	95	250 ACTON	93E OPAL	$f=\nu; F=\gamma\gamma$
$<6.5 \times 10^{-6}$	95	250 ACTON	93E OPAL	$f=e,\mu; F=\ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
$<7.1 \times 10^{-6}$	95	250 BUSKULIC	93F ALEP	$f=e,\mu; F=\ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
		252 ADRIANI	92F L3	$f=q; F=\gamma\gamma$

249 ABREU 96T obtain limit as a function of m_{X^0} . See their Fig. 6.

250 Limit is for m_{X^0} around 60 GeV.

251 ABREU 96T obtain limit as a function of m_{X^0} . See their Fig. 15.

252 ADRIANI 92F give $\sigma_Z \cdot B(Z \rightarrow q\bar{q}X^0) \cdot B(X^0 \rightarrow \gamma\gamma) < (0.75-1.5)$ pb (95%CL) for $m_{X^0} = 10-70$ GeV. The limit is 1 pb at 60 GeV.

Search for X^0 Resonance in $p\bar{p} \rightarrow WX^0$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	253 ABE	97W CDF	$X^0 \rightarrow b\bar{b}$

253 ABE 97W search for X^0 production associated with W in $p\bar{p}$ collisions at $E_{cm}=1.8$ TeV. The 95%CL upper limit on the production cross section times the branching ratio for $X^0 \rightarrow b\bar{b}$ ranges from 14 to 19 pb for X^0 mass between 70 and 120 GeV. See their Fig. 3 for upper limits of the production cross section as a function of m_{X^0} .

Search for Resonance X, Y in $e^+ e^- \rightarrow XY$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
	254 ALEXANDER 97B	OPAL	$X \rightarrow 2$ jets, $Y \rightarrow 2$ jets
	255 BUSKULIC,D 96	ALEP	$X \rightarrow 2$ jets, $Y \rightarrow 2$ jets
254	ALEXANDER 97B search for the associated production of two massive particles decaying into quarks in $e^+ e^-$ collisions at $\sqrt{s}=130-136$ GeV. The 95%CL upper limits on $\sigma(e^+ e^- \rightarrow XY)$ range from 2.7 to 4.5 pb for $95 < m_X + m_Y < 120$ GeV.		
255	BUSKULIC,D 96 observed an excess of four-jet production cross section in $e^+ e^-$ collisions at $\sqrt{s}=130-136$ GeV and find an enhancement in the sum of two dijet masses around 105 GeV.		

Heavy Particle Production in Quarkonium Decays

Limits are for branching ratios to modes shown.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
$< 1.5 \times 10^{-5}$	90	256 BALEST 95	CLE2	$\Upsilon(1S) \rightarrow X^0 \gamma$, $m_{X^0} < 5$ GeV
$< 3 \times 10^{-5} - 6 \times 10^{-3}$	90	257 BALEST 95	CLE2	$\Upsilon(1S) \rightarrow X^0 \bar{X}^0 \gamma$, $m_{X^0} < 3.9$ GeV
$< 5.6 \times 10^{-5}$	90	258 ANTREASYAN 90C	CBAL	$\Upsilon(1S) \rightarrow X^0 \gamma$, $m_{X^0} < 7.2$ GeV
		259 ALBRECHT 89	ARG	
256	BALEST 95 two-body limit is for pseudoscalar X^0 . The limit becomes $< 10^{-4}$ for $m_{X^0} < 7.7$ GeV.			
257	BALEST 95 three-body limit is for phase-space photon energy distribution and angular distribution same as for $\Upsilon \rightarrow gg\gamma$.			
258	ANTREASYAN 90C assume that X^0 does not decay in the detector.			
259	ALBRECHT 89 give limits for $B(\Upsilon(1S), \Upsilon(2S) \rightarrow X^0 \gamma) \cdot B(X^0 \rightarrow \pi^+ \pi^-, K^+ K^-, p \bar{p})$ for $m_{X^0} < 3.5$ GeV.			

REFERENCES FOR Searches for Heavy Bosons Other Than Higgs Bosons

ABBOTT	98E	PRL 80 2051	B. Abbott+	(D0 Collab.)
ABBOTT	97B	PRL 79 4321	+Abolins, Acharya+	(D0 Collab.)
ABE	97F	PRL 78 2906	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97G	PR D55 R5263	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97S	PRL 79 2192	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97W	PRL 79 3819	F. Abe+	(CDF Collab.)
ABE	97X	PRL 79 4327	+Akimoto, Akopian, Albrow, Amadon+	(CDF Collab.)
ACCIARRI	97Q	PL B412 201	M. Acciarri+	(L3 Collab.)
ALEXANDER	97B	ZPHY C73 201	G. Alexander+	(OPAL Collab.)
ARIMA	97	PR D55 19	+Odaka, Ogawa, Shirai, Tsuboyama+	(VENUS Collab.)
BARATE	97B	PL B399 329	+Buskulic, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BARENBOIM	97	PR D55 4213	+Bernabeu, Prades, Raidal	(VALE, IFIC)
DEANDREA	97	PL B409 277		(MARS)
DERRICK	97	ZPHY C73 613	M. Derrick+	(ZEUS Collab.)
GROSSMAN	97	PR D55 2768	+Ligeti, Nardi	(REHO, CIT)
JADACH	97	PL B408 281	+Ward, Was	(CERN, INPK, TENN, SLAC)
STAHL	97	ZPHY C74 73	A. Stahl, H. Voss	(BONN)
ABACHI	96C	PRL 76 3271	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABACHI	96D	PL B385 471	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABREU	96T	ZPHY C72 179	+Adam, Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
ADAM	96C	PL B380 471	+Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
AID	96B	PL B369 173	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ALLET	96	PL B383 139	+Bodek, Camps, Deutsch+	(VILL, LEUV, LOUV, WISC)

BUSKULIC	96N	PL B378 373	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC,D	96	ZPHY C71 179	D. Buskulic+	(ALEPH Collab.)
ABACHI	95E	PL B358 405	+Abbott, Abolins, Acharya, Adam, Adams+	(D0 Collab.)
ABACHI	95G	PRL 75 3618	+Abbott, Abolins, Acharya, Adam, Adams+	(D0 Collab.)
ABE	95	PR D51 R949	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95M	PRL 74 2900	+Albrow, Amidei, Antos, Anway-Wiese+	(CDF Collab.)
ABE	95N	PRL 74 3538	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
ABE	95U	PRL 75 1012	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
ABREU	95M	ZPHY C65 603	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
AID	95	PL B353 578	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
BALEST	95	PR D51 2053	+Cho, Ford, Johnson+	(CLEO Collab.)
KUZNETSOV	95	PRL 75 794	+Serebrov, Stepanenko+	(PNPI, KIAE, HARV, NIST)
KUZNETSOV	95B	PAN 58 2113	+Mikheev	(YARO)
		Translated from YAF 58 2228.		
MIZUKOSHI	95	NP B443 20	+Eboli, Gonzalez-Garcia	(SPAUL, CERN)
NARDI	95	PL B344 225	+Roulet, Tommasini	(MICH, CERN)
ABACHI	94B	PRL 72 965	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
ABREU	94O	ZPHY C64 183	+Adam, Adye, Agasi, Ajinenko, Aleksan+	(DELPHI Collab.)
AHMED	94B	ZPHY C64 545	+Aid, Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
BHATTACH...	94	PL B336 100	Bhattacharyya, Ellis, Sridhar	(CERN)
Also	94B	PL B338 522 (erratum)	Bhattacharyya, Ellis, Sridhar	(CERN)
BHATTACH...	94B	PL B338 522 (erratum)	Bhattacharyya, Ellis, Sridhar	(CERN)
BUSKULIC	94	ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
DAVIDSON	94	ZPHY C61 613	+Bailey, Campbell	(CFPA, TNT0, ALBE)
KUZNETSOV	94	PL B329 295	+Mikheev	(YARO)
KUZNETSOV	94B	JETPL 60 315	+Serebrov, Stepanenko+	(PNPI, KIAE, HARV, NIST)
		Translated from ZETFP 60 311.		
LEURER	94	PR D50 536		(REHO)
LEURER	94B	PR D49 333		(REHO)
Also	93	PRL 71 1324	Leurer	(REHO)
MAHANTA	94	PL B337 128		(MEHTA)
SEVERIJNS	94	PRL 73 611 (erratum)	+	(LOUV, WISC, LEUV, ETH, MASA)
VILAIN	94B	PL B332 465	+Wilquet, Beyer, Flegel, Grote+	(CHARM II Collab.)
ABE	93C	PL B302 119	+Amako, Arai, Arima, Asano, Chiba+	(VENUS Collab.)
ABE	93D	PL B304 373	+Adachi, Awa, Aoki, Belusevic, Emi+	(TOPAZ Collab.)
ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+	(CDF Collab.)
ABE	93I	PR D48 R3939	+Albrow, Amidei, Anway-Wiese, Apollinari+	(CDF Collab.)
ABREU	93J	PL B316 620	+Adam, Adye, Agasi, Aleksan, Alekseev+	(DELPHI Collab.)
ABT	93	NP B396 3	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ACTON	93E	PL B311 391	+Akers, Alexander+	(OPAL Collab.)
ADRIANI	93D	PL B306 187	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ADRIANI	93M	PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ALITTI	93	NP B400 3	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALLEN	93	PR D47 11	+Chen, Doe, Hausammann+	(UCI, LANL, ANL, UMD)
ALTARELLI	93B	PL B318 139	+Casalbuoni+	(CERN, FIRZ, GEVA, PADO)
BHATTACH...	93	PR D47 R3693	Bhattacharyya+	(CALC, JADA, ICTP, AHMED, BOSE)
BUSKULIC	93F	PL B308 425	+De Bonis, Decamp, Chez, Goy, Lees+	(ALEPH Collab.)
DERRICK	93	PL B306 173	+Krackauer, Magill, Musgrave, Repond+	(ZEUS Collab.)
RIZZO	93	PR D48 4470		(ANL)
SEVERIJNS	93	PRL 70 4047	+Gimeno-Nogues+	(LOUV, WISC, LEUV, ETH, MASA)
Also	94	PRL 73 611 (erratum)	Severijns+	(LOUV, WISC, LEUV, ETH, MASA)
STERNER	93	PL B303 385	+Abashian, Gotow, Haim, Mattson, Morgan+	(AMY Collab.)
ABE	92B	PRL 68 1463	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABREU	92D	ZPHY C53 555	+Adam, Adami, Adye, Akesson, Alekseev+	(DELPHI Collab.)
ABREU	92F	PL B275 222	+Adam, Adami, Adye, Akesson, Alekseev+	(DELPHI Collab.)
ADRIANI	92F	PL B292 472	+Aguilar-Benitez, Ahlen, Akbari, Alcazar+	(L3 Collab.)
ALITTI	92E	PL B274 507	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
DECAMP	92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
DELAGUILA	92	NP B372 3	del Aguila+	(CERN, GRAN, BRUXT, MADE)
Also	91C	NP B361 45	del Aguila, Moreno, Quiros	(BARC, MADE)
IMAZATO	92	PRL 69 877	+Kawashima, Tanaka+	(KEK, INUS, TOKY, TOKMS)
LANGACKER	92B	PR D45 278	+Luo	(PENN)
LAYSSAC	92	ZPHY C53 97	+Renard, Verzeznassi	(MONP, LAPP)
LAYSSAC	92B	PL B287 267	+Renard, Verzeznassi	(MONP, TRSTT)
LEIKE	92	PL B291 187	+Riemann, Riemann	(BERL, CERN)
LEP	92	PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collabs.)
MISHRA	92	PRL 68 3499	+Leung, Arroyo+	(COLU, CHIC, FNAL, ROCH, WISC)
POLAK	92	PL B276 492	+Zralek	(SILES)
POLAK	92B	PR D46 3871	+Zralek	(SILES)
RENTON	92	ZPHY C56 355		(OXF)

ABE	91D	PRL 67 2418	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	91F	PRL 67 2609	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ACTON	91	PL B268 122	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON	91B	PL B273 338	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADEVA	91B	PL B261 169	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
ADEVA	91D	PL B262 155	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
ALEXANDER	91	PL B263 123	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
ALITTI	91	ZPHY C49 17	+Ansari, Ansorge, Autiero, Bareyre+	(UA2 Collab.)
ALTARELLI	91	PL B261 146	+Casalbuoni, De Curtis+	(CERN, FIRZ, GEVA)
ALTARELLI	91B	PL B263 459	+Casalbuoni, De Curtis+	(CERN, FIRZ, GEVA)
Also	90	PL B245 669	Altarelli, Casalbuoni, Feruglio, Gatto	(CERN, LECE, GEVA)
AQUINO	91	PL B261 280	+Fernandez, Garcia	(CINV, PUEB)
BUCHMUEL...	91	PL B267 395	Buchmueller, Greub, Minkowski	(DESY, BERN)
COLANGELO	91	PL B253 154	+Nardulli	(BARI)
CUYPERS	91	PL B259 173	+Falk, Frampton	(DURH, HARV, UNCCH)
DELAGUILA	91	PL B254 497	del Aguila, Moreno, Quiros	(BARC, MADE, CERN)
FARAGGI	91	MPL A6 61	+Nanopoulos	(TAMU)
GEIREGAT	91	PL B259 499	+Vilain, Wilquet, Binder, Burkard+	(CHARM II Collab.)
GONZALEZ-G...	91	PL B259 365	Gonzalez-Garcia, Valle	(VALE)
Also	90C	NP B345 312	Gonzalez-Garcia, Valle	(VALE)
MAHANTHAP...	91	PR D43 3093	Mahanthappa, Mohapatra	(COLO)
Also	91B	PR D44 1616	Mahanthappa, Mohapatra	(COLO)
POLAK	91	NP B363 385	+Zralek	(SILES)
RIZZO	91	PR D44 202		(WISC, ISU)
WALKER	91	APJ 376 51	+Steigman, Schramm, Olive+	(HSCA, OSU, CHIC, MINN)
ABE	90F	PL B246 297	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABE	90G	PRL 65 2243	+Amidei, Apollinari, Atac+	(CDF Collab.)
ABE	90H	PR D41 1722	+Amidei, Apollinari, Ascoli, Atac+	(CDF Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ALLEN	90	PRL 64 1330	+Chen, Doe+	(UCI, LASL, UMD)
ANTREASYAN	90C	PL B251 204	+Bartels, Besset, Bieler, Bienlein+	(Crystal Ball Collab.)
BARGER	90B	PR D42 152	+Hewett, Rizzo	(WISC, ISU)
GLASHOW	90	PR D42 3224	+Sarid	(HARV)
GLASHOW	90B	PRL 64 725	+Sarid	(HARV)
GONZALEZ-G...	90D	PL B240 163	Gonzalez-Garcia, Valle	(VALE)
GRIFOLS	90	NP B331 244	+Masso	(BARC)
GRIFOLS	90C	MPL A5 2657		(BARC)
GRIFOLS	90D	PR D42 3293	+Masso, Rizzo	(BARC, CERN, WISC, ISU)
HAGIWARA	90	PR D41 815	+Najima, Sakuda, Terunuma	(KEK, DURH, YCC, HIRO)
HE	90B	PL B240 441	+Joshi, Volkas	(MELB)
Also	90C	PL B244 580	He, Joshi, Volkas	(MELB)
KIM	90	PL B240 243	+Breedon, Ko, Lander, Maeshima, Malchow+	(AMY Collab.)
LOPEZ	90	PL B241 392	+Nanopoulos	(TAMU)
ALBAJAR	89	ZPHY C44 15	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
ALBRECHT	89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
BARBIERI	89B	PR D39 1229	+Mohapatra	(PISA, UMD)
DELAGUILA	89	PR D40 2481	del Aguila, Moreno, Quiros	(BARC, MADE)
Also	90B	PR D41 134	del Aguila, Moreno, Quiros	(BARC, MADE)
Also	90C	PR D42 262	del Aguila, Moreno, Quiros	(BARC, MADE)
DORENBOS...	89	ZPHY C41 567	Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
GEIREGAT	89	PL B232 539	+Vilain, Wilquet, Bergsma, Binder+	(CHARM II Collab.)
LANGACKER	89B	PR D40 1569	+Uma Sankar	(PENN)
ODAKA	89	JPSJ 58 3037	+Kondo, Abe, Amako+	(VENUS Collab.)
ROBINETT	89	PR D39 834		(PSU)
ALBAJAR	88B	PL B209 127	+Albrow, Allkofer, Astbury, Aubert+	(UA1 Collab.)
BAGGER	88	PR D37 1188	+Schmidt, King	(HARV, BOST)
BALKE	88	PR D37 587	+Gidal, Jodidio+	(LBL, UCB, COLO, NWES, TRIU)
BERGSTROM	88	PL B212 386		(STOH)
COSTA	88	NP B297 244	+Ellis, Fogli+	(PADO, CERN, BARI, WISC, LBL)
CUYPERS	88	PRL 60 1237	+Frampton	(UNCCH)
DONCHESKI	88	PL B206 137	+Grotch, Robinett	(PSU)
DONCHESKI	88B	PR D38 412	+Grotch, Robinett	(PSU)
ELLIS	88	PL B202 417	Ellis, Franzini, Zwirner	(CERN, UCB, LBL)
RAFFELT	88	PRL 60 1793	+Seckel	(UCB, LLL, UCSC)
AMALDI	87	PR D36 1385	+Bohm, Durkin, Langacker+	(CERN, AACH3, OSU+)
ANSARI	87D	PL B195 613	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	87B	ZPHY C36 15	+Becker, Felst+	(JADE Collab.)
MARCIANO	87	PR D35 1672	+Sirlin	(BNL, NYU)
ARNISON	86B	EPL 1 327	+Albrow, Allkofer+	(UA1 Collab.)

BARGER	86B	PRL 56 30	+Deshpande, Whisnant	(WISC, OREG, FSU)
BEHREND	86B	PL B178 452	+Buerger, Criegee, Fenner, Field+	(CELLO Collab.)
DERRICK	86	PL 166B 463	+Gan, Kooijman, Loos+	(HRS Collab.)
Also	86B	PR D34 3286	Derrick, Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
DURKIN	86	PL 166B 436	+Langacker	(PENN)
ELLIS	86	PL 167B 457	+Enqvist, Nanopoulos, Sarkar	(CERN, OXFTP)
JODIDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	Jodidio, Balke, Carr+	(LBL, NWES, TRIU)
MOHAPATRA	86	PR D34 909		(UMD)
STEIGMAN	86	PL B176 33	+Olive, Schramm, Turner	(BART, MINN+)
ADEVA	85	PL 152B 439	+Becker, Becker-Szendy+	(Mark-J Collab.)
ADEVA	85B	PRL 55 665	+Becker, Becker-Szendy+	(Mark-J Collab.)
BERGER	85B	ZPHY C27 341	+Deuter, Genzel, Lackas, Pielorz+	(PLUTO Collab.)
STOKER	85	PRL 54 1887	+Balke, Carr, Gidal+	(LBL, NWES, TRIU)
ADEVA	84	PRL 53 134	+Barber, Becker, Berdugo+	(Mark-J Collab.)
BEHREND	84C	PL 140B 130	+Burger, Criegee, Fenner+	(CELLO Collab.)
ARNISON	83D	PL 129B 273	+Astbury, Aubert, Bacci+	(UA1 Collab.)
BERGSMAN	83	PL 122B 465	+Dorenbosch, Jonker+	(CHARM Collab.)
CARR	83	PRL 51 627	+Gidal, Gobbi, Jodidio, Oram+	(LBL, NWES, TRIU)
DESHPANDE	83	PR D27 1193	+Johnson	(OREG)
BEALL	82	PRL 48 848	+Bander, Soni	(UCI, UCLA)
SHANKER	82	NP B204 375		(TRIU)
DIMOPOUL...	81	NP B182 77	Dimopoulos, Raby, Kane	(STAN, MICH)
RIZZO	81	PR D24 704	+Senjanovic	(BNL)
STEIGMAN	79	PRL 43 239	+Olive, Schramm	(BART, EFI)