


 $I(J^P) = 0(?)$ 

$J^P$  is natural, width and decay modes consistent with  $1^-$ .

### $D_s^{*\pm}$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ , and  $D_s^{*\pm}$  mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2112.4 ± 0.7 OUR FIT</b>	Error includes scale factor of 1.1.		
<b>2106.6 ± 2.1 ± 2.7</b>	<sup>1</sup> BLAYLOCK 87 MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$		

<sup>1</sup> Assuming  $D_s^\pm$  mass =  $1968.7 \pm 0.9$  MeV.

### $m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ , and  $D_s^{*\pm}$  mass and mass difference measurements.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>143.8 ± 0.4 OUR FIT</b>				
<b>143.9 ± 0.4 OUR AVERAGE</b>				
143.76 ± 0.39 ± 0.40		GRONBERG 95	CLE2	$e^+ e^-$
144.22 ± 0.47 ± 0.37		BROWN 94	CLE2	$e^+ e^-$
142.5 ± 0.8 ± 1.5		<sup>2</sup> ALBRECHT 88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
139.5 ± 8.3 ± 9.7	60	AIHARA 84D	TPC	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
143.0 ± 18.0	8	ASRATYAN 85	HLBC	FNAL 15-ft, $\nu^2 H$
110 ± 46		BRANDELIK 79	DASP	$e^+ e^- \rightarrow D_s^\pm \gamma X$

<sup>2</sup> Result includes data of ALBRECHT 84B.

### $D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	GRONBERG 95	CLE2	$e^+ e^-$
< 4.5	90	ALBRECHT 88	ARG	$E_{cm}^{ee} = 10.2$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.9	90	BROWN 94	CLE2	$e^+ e^-$
< 22	90	BLAYLOCK 87	MRK3	$e^+ e^- \rightarrow D_s^\pm \gamma X$

**$D_s^{*+}$  DECAY MODES**

$D_s^{*-}$  modes are charge conjugates of the modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad D_s^+ \gamma$	(94.2±2.5) %
$\Gamma_2 \quad D_s^+ \pi^0$	( 5.8±2.5) %

**CONSTRAINED FIT INFORMATION**

An overall fit to a branching ratio uses 1 measurements and one constraint to determine 2 parameters. The overall fit has a  $\chi^2 = 0.0$  for 0 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{matrix} x_2 & | & -100 \\ & | \\ x_1 & & \end{matrix}$$

 **$D_s^{*+}$  BRANCHING RATIOS**

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$	$\Gamma_1 / \Gamma$
VALUE <b>0.942±0.026 OUR FIT</b>	

• • • We do not use the following data for averages, fits, limits, etc. • • •
seen ASRATYAN 91 HLBC $\bar{\nu}_\mu$ Ne
seen ALBRECHT 88 ARG $e^+ e^- \rightarrow D_s^\pm \gamma X$
seen AIHARA 84D
seen ALBRECHT 84B
seen BRANDELIK 79

$\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$	$\Gamma_2 / \Gamma_1$
VALUE <b>0.062±0.029 OUR FIT</b>	
0.062 <sup>+0.020</sup> <sub>-0.018</sub> ±0.022	GRONBERG 95 CLE2 $e^+ e^-$

 **$D_s^{*\pm}$  REFERENCES**

GRONBERG 95 PRL 75 3232	+Korte, Kutschke+	(CLEO Collab.)
BROWN 94 PR D50 1884	+Fast, McIlwain, Miao+	(CLEO Collab.)
ASRATYAN 91 PL B257 525	+Marage+(ITEP, BELG, SACL, SERP, CRAC, BARI, CERN)	
ALBRECHT 88 PL B207 349	+Binder, Boeckmann+	(ARGUS Collab.)
BLAYLOCK 87 PRL 58 2171	+Bolton, Brown, Bunnell+	(Mark III Collab.)
ASRATYAN 85 PL 156B 441	+Fedotov, Ammosov, Burtovoy+	(ITEP, SERP)
AIHARA 84D PRL 53 2465	+Alston-Garnjost, Badtke, Bakken+	(TPC Collab.)
ALBRECHT 84B PL 146B 111	+Drescher, Heller+	(ARGUS Collab.)
BRANDELIK 79 PL 80B 412	+Braunschweig, Martyn, Sander+	(DASP Collab.)

———— OTHER RELATED PAPERS ——

KAMAL	92	PL B284 421	+Xu	(ALBE)
BRANDELIK	78C	PL 76B 361	+Cords+	(DASP Collab.)
BRANDELIK	77B	PL 70B 132	+Braunschweig, Martyn, Sander+	(DASP Collab.)

---