BOTTOM BARYONS ($B = -1$)

$\Lambda_b^0 = udb$, $\Xi_b^0 = usb$, $\Xi_b^- = dsb$

Λ_b^0

$I(J^P) = 0(\frac{1}{2}^+)$

$I(J^P)$ not yet measured; $0(\frac{1}{2}^+)$ is the quark model prediction.

Mass $m = 5624 \pm 9$ MeV (S = 1.8)

Mean life $\tau = (1.24 \pm 0.08) \times 10^{-12}$ s

$c\tau = 372$ μm

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p}$), branching ratios, and detection efficiencies. They scale with the LEP Λ_b production fraction B($b \rightarrow \Lambda_b$) and are evaluated for our value B($b \rightarrow \Lambda_b$) = $(10.1^{+3.9}_{-3.1})$%.

The branching fractions B(b-baryon $\rightarrow \Lambda \ell^- \overline{\nu}_\ell$ anything) and B($\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \overline{\nu}_\ell$ anything) are not pure measurements because the underlying measured products of these with B($b \rightarrow \Lambda_b$) were used to determine B($b \rightarrow \Lambda_b$), as described in the note "Production and Decay of b-Flavored Hadrons."

Λ_b^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$J/\psi(1S)\Lambda$	$(4.7 \pm 2.8) \times 10^{-4}$		1744
$\Lambda_c^+ \pi^-$	seen		2345
$\Lambda_c^+ a_1(1260)^-$	seen		2156
$\Lambda_c^+ \ell^- \overline{\nu}_\ell$ anything	[p] $(9.0^{+3.1}_{-3.8})$ %		–
$p\pi^-$	$< 5.0 \times 10^{-5}$	90%	2732
pK^-	$< 5.0 \times 10^{-5}$	90%	2711

b-baryon ADMIXTURE (Λ_b, Ξ_b, Σ_b, Ω_b)

Mean life $\tau = (1.20 \pm 0.07) \times 10^{-12}$ s

These branching fractions are actually an average over weakly decaying *b*-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p}$), branching ratios, and detection efficiencies. They scale with the LEP Λ_b production fraction B($b \rightarrow \Lambda_b$) and are evaluated for our value B($b \rightarrow \Lambda_b$) = $(10.1^{+3.9}_{-3.1})$%.

The branching fractions B(*b*-baryon $\rightarrow \Lambda \ell^- \overline{\nu}_\ell$ anything) and B($\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \overline{\nu}_\ell$ anything) are not pure measurements because the underlying measured products of these with B($b \rightarrow \Lambda_b$) were used to determine B($b \rightarrow \Lambda_b$), as described in the note "Production and Decay of *b*-Flavored Hadrons."

***b*-baryon ADMIXTURE (Λ_b,Ξ_b,Σ_b,Ω_b)**	Fraction (Γ_i/Γ)	*p* (MeV/*c*)
$p\mu^-\overline{\nu}$ anything	(4.9± 2.4) %	–
$\Lambda\ell^-\overline{\nu}_\ell$ anything	($3.1^{+1.0}_{-1.2}$) %	–
$\Lambda/\overline{\Lambda}$ anything	(35 $^{+12}_{-14}$) %	–
$\Xi^-\ell^-\overline{\nu}_\ell$ anything	($5.5^{+2.0}_{-2.4}$) $\times 10^{-3}$	–