

## $\Delta(1750)$ $P_{31}$

$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$  Status: \*

OMITTED FROM SUMMARY TABLE

### $\Delta(1750)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>≈ 1750 OUR ESTIMATE</b>			
1744 ± 36	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1715.2 ± 21.0	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1778.4 ± 9.0	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

### $\Delta(1750)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 ± 120	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
93.3 ± 55.0	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
23.0 ± 29.0	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

### $\Delta(1750)$ DECAY MODES

Mode
$\Gamma_1 N\pi$
$\Gamma_2 N\pi\pi$
$\Gamma_3 N(1440)\pi$

### $\Gamma(N\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.08 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.18	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
0.20	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

### $\Gamma_1/\Gamma$

$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow N(1440)\pi$	DOCUMENT ID	TECN	COMMENT
+0.15 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$

### $(\Gamma_1 \Gamma_3)^{1/2}/\Gamma$

### $\Delta(1750)$ FOOTNOTES

<sup>1</sup> CHEW 80 reports four resonances in the  $P_{31}$  wave — see also the  $\Delta(1910)$ . Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.

## **$\Delta(1750)$ REFERENCES**

MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT)
Also	84	PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
HOEHLER	83	Landolt-Bornstein 1/9B2	G. Hohler	(KARLT)
CHEW	80	Toronto Conf.	123	D.M. Chew (LBL)

---