

N BARYONS

$(S = 0, I = 1/2)$

$p, N^+ = uud; \quad n, N^0 = udd$

p

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 938.27200 \pm 0.00004$ MeV [a]

$$= 1.00727646688 \pm 0.00000000013 \text{ u}$$

$$|m_p - m_{\bar{p}}|/m_p < 5 \times 10^{-7} \text{ [b]}$$

$$|\frac{q_p}{m_p}| / (\frac{q_{\bar{p}}}{m_{\bar{p}}}) = 0.99999999991 \pm 0.00000000009$$

$$|q_p + q_{\bar{p}}|/e < 5 \times 10^{-7} \text{ [b]}$$

$$|q_p + q_e|/e < 1.0 \times 10^{-21} \text{ [c]}$$

Magnetic moment $\mu = 2.792847337 \pm 0.000000029 \mu_N$

$$(\mu_p + \mu_{\bar{p}}) / \mu_p = (-2.6 \pm 2.9) \times 10^{-3}$$

Electric dipole moment $d = (-4 \pm 6) \times 10^{-23} \text{ ecm}$

Electric polarizability $\bar{\alpha} = (12.1 \pm 0.9) \times 10^{-4} \text{ fm}^3$

Magnetic polarizability $\bar{\beta} = (2.1 \pm 0.9) \times 10^{-4} \text{ fm}^3$

Mean life $\tau > 1.6 \times 10^{25}$ years (independent of mode)

> 10^{31} to 10^{33} years [d] (mode dependent)

Below, for N decays, p and n distinguish proton and neutron partial lifetimes. See also the "Note on Nucleon Decay" in our 1994 edition (Phys. Rev. **D50**, 1673) for a short review.

The "partial mean life" limits tabulated here are the limits on τ/B_i , where τ is the total mean life and B_i is the branching fraction for the mode in question.

p DECAY MODES	Partial mean life (10^{30} years)	p
		Confidence level
Antilepton + meson		
$N \rightarrow e^+ \pi$	> 158 (n), > 1600 (p)	90% 459
$N \rightarrow \mu^+ \pi$	> 100 (n), > 473 (p)	90% 453
$N \rightarrow \nu \pi$	> 112 (n), > 25 (p)	90% 459
$p \rightarrow e^+ \eta$	> 313	90% 309
$p \rightarrow \mu^+ \eta$	> 126	90% 296
$n \rightarrow \nu \eta$	> 158	90% 310
$N \rightarrow e^+ \rho$	> 217 (n), > 75 (p)	90% 153
$N \rightarrow \mu^+ \rho$	> 228 (n), > 110 (p)	90% 119
$N \rightarrow \nu \rho$	> 19 (n), > 162 (p)	90% 153
$p \rightarrow e^+ \omega$	> 107	90% 142

$p \rightarrow \mu^+ \omega$	> 117	90%	104
$n \rightarrow \nu \omega$	> 108	90%	144
$N \rightarrow e^+ K$	> 17 (n), > 150 (p)	90%	337
$p \rightarrow e^+ K_S^0$	> 76	90%	337
$p \rightarrow e^+ K_L^0$	> 44	90%	337
$N \rightarrow \mu^+ K$	> 26 (n), > 120 (p)	90%	326
$p \rightarrow \mu^+ K_S^0$	> 64	90%	326
$p \rightarrow \mu^+ K_L^0$	> 44	90%	326
$N \rightarrow \nu K$	> 86 (n), > 670 (p)	90%	339
$p \rightarrow e^+ K^*(892)^0$	> 84	90%	45
$N \rightarrow \nu K^*(892)$	> 78 (n), > 51 (p)	90%	45

Antilepton + mesons

$p \rightarrow e^+ \pi^+ \pi^-$	> 82	90%	448
$p \rightarrow e^+ \pi^0 \pi^0$	> 147	90%	449
$n \rightarrow e^+ \pi^- \pi^0$	> 52	90%	449
$p \rightarrow \mu^+ \pi^+ \pi^-$	> 133	90%	425
$p \rightarrow \mu^+ \pi^0 \pi^0$	> 101	90%	427
$n \rightarrow \mu^+ \pi^- \pi^0$	> 74	90%	427
$n \rightarrow e^+ K^0 \pi^-$	> 18	90%	319

Lepton + meson

$n \rightarrow e^- \pi^+$	> 65	90%	459
$n \rightarrow \mu^- \pi^+$	> 49	90%	453
$n \rightarrow e^- \rho^+$	> 62	90%	154
$n \rightarrow \mu^- \rho^+$	> 7	90%	120
$n \rightarrow e^- K^+$	> 32	90%	340
$n \rightarrow \mu^- K^+$	> 57	90%	330

Lepton + mesons

$p \rightarrow e^- \pi^+ \pi^+$	> 30	90%	448
$n \rightarrow e^- \pi^+ \pi^0$	> 29	90%	449
$p \rightarrow \mu^- \pi^+ \pi^+$	> 17	90%	425
$n \rightarrow \mu^- \pi^+ \pi^0$	> 34	90%	427
$p \rightarrow e^- \pi^+ K^+$	> 75	90%	320
$p \rightarrow \mu^- \pi^+ K^+$	> 245	90%	279

Antilepton + photon(s)

$p \rightarrow e^+ \gamma$	> 670	90%	469
$p \rightarrow \mu^+ \gamma$	> 478	90%	463
$n \rightarrow \nu \gamma$	> 28	90%	470
$p \rightarrow e^+ \gamma \gamma$	> 100	90%	469
$n \rightarrow \nu \gamma \gamma$	> 219	90%	470

Three (or more) leptons

$p \rightarrow e^+ e^+ e^-$	> 793	90%	469
$p \rightarrow e^+ \mu^+ \mu^-$	> 359	90%	457
$p \rightarrow e^+ \nu \nu$	> 17	90%	469
$n \rightarrow e^+ e^- \nu$	> 257	90%	470
$n \rightarrow \mu^+ e^- \nu$	> 83	90%	464
$n \rightarrow \mu^+ \mu^- \nu$	> 79	90%	458
$p \rightarrow \mu^+ e^+ e^-$	> 529	90%	464
$p \rightarrow \mu^+ \mu^+ \mu^-$	> 675	90%	439
$p \rightarrow \mu^+ \nu \nu$	> 21	90%	463
$p \rightarrow e^- \mu^+ \mu^+$	> 6	90%	457
$n \rightarrow 3\nu$	> 0.0005	90%	470

Inclusive modes

$N \rightarrow e^+ \text{anything}$	> 0.6 (n, p)	90%	—
$N \rightarrow \mu^+ \text{anything}$	> 12 (n, p)	90%	—
$N \rightarrow e^+ \pi^0 \text{anything}$	> 0.6 (n, p)	90%	—

$\Delta B = 2$ dinucleon modes

The following are lifetime limits per iron nucleus.

$pp \rightarrow \pi^+ \pi^+$	> 0.7	90%	—
$pn \rightarrow \pi^+ \pi^0$	> 2	90%	—
$nn \rightarrow \pi^+ \pi^-$	> 0.7	90%	—
$nn \rightarrow \pi^0 \pi^0$	> 3.4	90%	—
$pp \rightarrow e^+ e^+$	> 5.8	90%	—
$pp \rightarrow e^+ \mu^+$	> 3.6	90%	—
$pp \rightarrow \mu^+ \mu^+$	> 1.7	90%	—
$pn \rightarrow e^+ \bar{\nu}$	> 2.8	90%	—
$pn \rightarrow \mu^+ \bar{\nu}$	> 1.6	90%	—
$nn \rightarrow \nu_e \bar{\nu}_e$	> 0.000012	90%	—
$nn \rightarrow \nu_\mu \bar{\nu}_\mu$	> 0.000006	90%	—

\bar{p} DECAY MODES

\bar{p} DECAY MODES	Partial mean life (years)	Confidence level	p (MeV/c)
$\bar{p} \rightarrow e^- \gamma$	> 7×10^5	90%	469
$\bar{p} \rightarrow \mu^- \gamma$	> 5×10^4	90%	463
$\bar{p} \rightarrow e^- \pi^0$	> 4×10^5	90%	459
$\bar{p} \rightarrow \mu^- \pi^0$	> 5×10^4	90%	453
$\bar{p} \rightarrow e^- \eta$	> 2×10^4	90%	309
$\bar{p} \rightarrow \mu^- \eta$	> 8×10^3	90%	296
$\bar{p} \rightarrow e^- K_S^0$	> 900	90%	337

$\bar{p} \rightarrow \mu^- K_S^0$	$> 4 \times 10^3$	90%	326
$\bar{p} \rightarrow e^- K_L^0$	$> 9 \times 10^3$	90%	337
$\bar{p} \rightarrow \mu^- K_L^0$	$> 7 \times 10^3$	90%	326
$\bar{p} \rightarrow e^- \gamma\gamma$	$> 2 \times 10^4$	90%	469
$\bar{p} \rightarrow \mu^- \gamma\gamma$	$> 2 \times 10^4$	90%	463
$\bar{p} \rightarrow e^- \rho$	> 200	90%	153
$\bar{p} \rightarrow e^- \omega$	> 200	90%	142
$\bar{p} \rightarrow e^- K^*(892)^0$	$> 1 \times 10^3$	90%	141

n

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 939.56533 \pm 0.00004$ MeV [a]
 $= 1.00866491578 \pm 0.00000000055$ u
 $m_n - m_p = 1.2933318 \pm 0.0000005$ MeV
 $= 0.0013884489 \pm 0.0000000006$ u

Mean life $\tau = 886.7 \pm 1.9$ s (S = 1.2)
 $c\tau = 2.658 \times 10^8$ km

Magnetic moment $\mu = -1.9130427 \pm 0.0000005$ μ_N
Electric dipole moment $d < 0.63 \times 10^{-25}$ e cm, CL = 90%
Electric polarizability $\alpha = (0.98^{+0.19}_{-0.23}) \times 10^{-3}$ fm 3 (S = 1.1)
Charge $q = (-0.4 \pm 1.1) \times 10^{-21}$ e
Mean $n\bar{n}$ -oscillation time $> 8.6 \times 10^7$ s, CL = 90% (free n)
 $> 1.2 \times 10^8$ s, CL = 90% [e] (bound n)

Decay parameters [f]

$p e^- \bar{\nu}_e$	$g_A/g_V = -1.2670 \pm 0.0035$ (S = 1.9)
"	$A = -0.1162 \pm 0.0013$ (S = 1.8)
"	$B = 0.983 \pm 0.004$
"	$a = -0.102 \pm 0.005$
"	$\phi_{AV} = (180.07 \pm 0.18)^\circ$ [g]
"	$D = (-0.5 \pm 1.4) \times 10^{-3}$

n DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	$\frac{p}{(MeV/c)}$
$p e^- \bar{\nu}_e$	100 %		1.19
Charge conservation (Q) violating mode			
$p \nu_e \bar{\nu}_e$	$Q < 8 \times 10^{-27}$	68%	1.29

N(1440) P_{11}

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Breit-Wigner mass = 1430 to 1470 (≈ 1440) MeVBreit-Wigner full width = 250 to 450 (≈ 350) MeV

$$\rho_{\text{beam}} = 0.61 \text{ GeV}/c \quad 4\pi\lambda^2 = 31.0 \text{ mb}$$

Re(pole position) = 1345 to 1385 (≈ 1365) MeV- 2Im(pole position) = 160 to 260 (≈ 210) MeV

N(1440) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	60–70 %	397
$N\pi\pi$	30–40 %	342
$\Delta\pi$	20–30 %	143
$N\rho$	<8 %	†
$N(\pi\pi)^{I=0}_{S\text{-wave}}$	5–10 %	—
$p\gamma$	0.035–0.048 %	414
$p\gamma$, helicity=1/2	0.035–0.048 %	414
$n\gamma$	0.009–0.032 %	413
$n\gamma$, helicity=1/2	0.009–0.032 %	413

N(1520) D_{13}

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Breit-Wigner mass = 1515 to 1530 (≈ 1520) MeVBreit-Wigner full width = 110 to 135 (≈ 120) MeV

$$\rho_{\text{beam}} = 0.74 \text{ GeV}/c \quad 4\pi\lambda^2 = 23.5 \text{ mb}$$

Re(pole position) = 1505 to 1515 (≈ 1510) MeV- 2Im(pole position) = 110 to 120 (≈ 115) MeV

N(1520) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	50–60 %	456
$N\pi\pi$	40–50 %	410
$\Delta\pi$	15–25 %	228
$N\rho$	15–25 %	†
$N(\pi\pi)^{I=0}_{S\text{-wave}}$	<8 %	—
$p\gamma$	0.46–0.56 %	470
$p\gamma$, helicity=1/2	0.001–0.034 %	470
$p\gamma$, helicity=3/2	0.44–0.53 %	470
$n\gamma$	0.30–0.53 %	470
$n\gamma$, helicity=1/2	0.04–0.10 %	470
$n\gamma$, helicity=3/2	0.25–0.45 %	470

N(1535) S₁₁

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Breit-Wigner mass = 1520 to 1555 (\approx 1535) MeVBreit-Wigner full width = 100 to 250 (\approx 150) MeV

$$p_{\text{beam}} = 0.76 \text{ GeV}/c \quad 4\pi\lambda^2 = 22.5 \text{ mb}$$

Re(pole position) = 1495 to 1515 (\approx 1505) MeV- 2Im(pole position) = 90 to 250 (\approx 170) MeV

N(1535) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	35–55 %	467
$N\eta$	30–55 %	182
$N\pi\pi$	1–10 %	422
$\Delta\pi$	<1 %	242
$N\rho$	<4 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	<3 %	—
$N(1440)\pi$	<7 %	†
$p\gamma$	0.15–0.35 %	481
$p\gamma$, helicity=1/2	0.15–0.35 %	481
$n\gamma$	0.004–0.29 %	480
$n\gamma$, helicity=1/2	0.004–0.29 %	480

N(1650) S₁₁

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Breit-Wigner mass = 1640 to 1680 (\approx 1650) MeVBreit-Wigner full width = 145 to 190 (\approx 150) MeV

$$p_{\text{beam}} = 0.96 \text{ GeV}/c \quad 4\pi\lambda^2 = 16.4 \text{ mb}$$

Re(pole position) = 1640 to 1680 (\approx 1660) MeV- 2Im(pole position) = 150 to 170 (\approx 160) MeV

N(1650) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	55–90 %	547
$N\eta$	3–10 %	346
ΛK	3–11 %	161
$N\pi\pi$	10–20 %	511
$\Delta\pi$	1–7 %	344
$N\rho$	4–12 %	†

$N(\pi\pi)^{I=0}_{S\text{-wave}}$	<4 %	—
$N(1440)\pi$	<5 %	147
$p\gamma$	0.04–0.18 %	558
$p\gamma$, helicity=1/2	0.04–0.18 %	558
$n\gamma$	0.003–0.17 %	557
$n\gamma$, helicity=1/2	0.003–0.17 %	557

N(1675) D_{15}

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-)$$

Breit-Wigner mass = 1670 to 1685 (≈ 1675) MeVBreit-Wigner full width = 140 to 180 (≈ 150) MeV $p_{\text{beam}} = 1.01 \text{ GeV}/c \quad 4\pi\lambda^2 = 15.4 \text{ mb}$ Re(pole position) = 1655 to 1665 (≈ 1660) MeV–2Im(pole position) = 125 to 155 (≈ 140) MeV

N(1675) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	40–50 %	563
ΛK	<1 %	209
$N\pi\pi$	50–60 %	529
$\Delta\pi$	50–60 %	364
$N\rho$	< 1–3 %	†
$p\gamma$	0.004–0.023 %	575
$p\gamma$, helicity=1/2	0.0–0.015 %	575
$p\gamma$, helicity=3/2	0.0–0.011 %	575
$n\gamma$	0.02–0.12 %	574
$n\gamma$, helicity=1/2	0.006–0.046 %	574
$n\gamma$, helicity=3/2	0.01–0.08 %	574

N(1680) F_{15}

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^+)$$

Breit-Wigner mass = 1675 to 1690 (≈ 1680) MeVBreit-Wigner full width = 120 to 140 (≈ 130) MeV $p_{\text{beam}} = 1.01 \text{ GeV}/c \quad 4\pi\lambda^2 = 15.2 \text{ mb}$ Re(pole position) = 1665 to 1675 (≈ 1670) MeV–2Im(pole position) = 105 to 135 (≈ 120) MeV

N(1680) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	60–70 %	567
$N\pi\pi$	30–40 %	532
$\Delta\pi$	5–15 %	369
$N\rho$	3–15 %	†
$N(\pi\pi)^{I=0}_{S\text{-wave}}$	5–20 %	—

$p\gamma$	0.21–0.32 %	578
$p\gamma$, helicity=1/2	0.001–0.011 %	578
$p\gamma$, helicity=3/2	0.20–0.32 %	578
$n\gamma$	0.021–0.046 %	577
$n\gamma$, helicity=1/2	0.004–0.029 %	577
$n\gamma$, helicity=3/2	0.01–0.024 %	577

N(1700) D_{13}

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Breit-Wigner mass = 1650 to 1750 (≈ 1700) MeVBreit-Wigner full width = 50 to 150 (≈ 100) MeV

$$p_{\text{beam}} = 1.05 \text{ GeV}/c \quad 4\pi\lambda^2 = 14.5 \text{ mb}$$

Re(pole position) = 1630 to 1730 (≈ 1680) MeV–2Im(pole position) = 50 to 150 (≈ 100) MeV

N(1700) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	5–15 %	580
ΛK	<3 %	250
$N\pi\pi$	85–95 %	547
$N\rho$	<35 %	†
$p\gamma$	0.01–0.05 %	591
$p\gamma$, helicity=1/2	0.0–0.024 %	591
$p\gamma$, helicity=3/2	0.002–0.026 %	591
$n\gamma$	0.01–0.13 %	590
$n\gamma$, helicity=1/2	0.0–0.09 %	590
$n\gamma$, helicity=3/2	0.01–0.05 %	590

N(1710) P_{11}

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Breit-Wigner mass = 1680 to 1740 (≈ 1710) MeVBreit-Wigner full width = 50 to 250 (≈ 100) MeV

$$p_{\text{beam}} = 1.07 \text{ GeV}/c \quad 4\pi\lambda^2 = 14.2 \text{ mb}$$

Re(pole position) = 1670 to 1770 (≈ 1720) MeV–2Im(pole position) = 80 to 380 (≈ 230) MeV

N(1710) DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	10–20 %	587
ΛK	5–25 %	264
$N\pi\pi$	40–90 %	554
$\Delta\pi$	15–40 %	393
$N\rho$	5–25 %	48

$N(\pi\pi)_{S\text{-wave}}^{I=0}$	10–40 %	—
$p\gamma$	0.002–0.05%	598
$p\gamma$, helicity=1/2	0.002–0.05%	598
$n\gamma$	0.0–0.02%	597
$n\gamma$, helicity=1/2	0.0–0.02%	597

 $N(1720)$ P_{13}

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

Breit-Wigner mass = 1650 to 1750 (≈ 1720) MeVBreit-Wigner full width = 100 to 200 (≈ 150) MeV

$$p_{\text{beam}} = 1.09 \text{ GeV}/c \quad 4\pi\lambda^2 = 13.9 \text{ mb}$$

Re(pole position) = 1650 to 1750 (≈ 1700) MeV–2Im(pole position) = 110 to 390 (≈ 250) MeV

$N(1720)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	10–20 %	594
ΛK	1–15 %	278
$N\pi\pi$	>70 %	561
$N\rho$	70–85 %	104
$p\gamma$	0.003–0.10 %	604
$p\gamma$, helicity=1/2	0.003–0.08 %	604
$p\gamma$, helicity=3/2	0.001–0.03 %	604
$n\gamma$	0.002–0.39 %	603
$n\gamma$, helicity=1/2	0.0–0.002 %	603
$n\gamma$, helicity=3/2	0.001–0.39 %	603

 $N(2190)$ G_{17}

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^-)$$

Breit-Wigner mass = 2100 to 2200 (≈ 2190) MeVBreit-Wigner full width = 350 to 550 (≈ 450) MeV

$$p_{\text{beam}} = 2.07 \text{ GeV}/c \quad 4\pi\lambda^2 = 6.21 \text{ mb}$$

Re(pole position) = 1950 to 2150 (≈ 2050) MeV–2Im(pole position) = 350 to 550 (≈ 450) MeV

$N(2190)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	10–20 %	888

N(2220) H₁₉

$$I(J^P) = \frac{1}{2}(\frac{9}{2}+)$$

Breit-Wigner mass = 2180 to 2310 (\approx 2220) MeV

Breit-Wigner full width = 320 to 550 (\approx 400) MeV

$$p_{\text{beam}} = 2.14 \text{ GeV}/c \quad 4\pi\lambda^2 = 5.97 \text{ mb}$$

Re(pole position) = 2100 to 2240 (\approx 2170) MeV

- 2Im(pole position) = 370 to 570 (\approx 470) MeV

N(2220) DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\pi$

10–20 %

905

N(2250) G₁₉

$$I(J^P) = \frac{1}{2}(\frac{9}{2}-)$$

Breit-Wigner mass = 2170 to 2310 (\approx 2250) MeV

Breit-Wigner full width = 290 to 470 (\approx 400) MeV

$$p_{\text{beam}} = 2.21 \text{ GeV}/c \quad 4\pi\lambda^2 = 5.74 \text{ mb}$$

Re(pole position) = 2080 to 2200 (\approx 2140) MeV

- 2Im(pole position) = 280 to 680 (\approx 480) MeV

N(2250) DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\pi$

5–15 %

923

N(2600) I_{1,11}

$$I(J^P) = \frac{1}{2}(\frac{11}{2}-)$$

Breit-Wigner mass = 2550 to 2750 (\approx 2600) MeV

Breit-Wigner full width = 500 to 800 (\approx 650) MeV

$$p_{\text{beam}} = 3.12 \text{ GeV}/c \quad 4\pi\lambda^2 = 3.86 \text{ mb}$$

N(2600) DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\pi$

5–10 %

1126

Δ BARYONS ($S = 0, I = 3/2$)

$$\Delta^{++} = uuu, \quad \Delta^+ = uud, \quad \Delta^0 = udd, \quad \Delta^- = ddd$$

$\Delta(1232)$ P_{33}

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Breit-Wigner mass (mixed charges) = 1230 to 1234 (≈ 1232) MeV

Breit-Wigner full width (mixed charges) = 115 to 125 (≈ 120) MeV

$$p_{\text{beam}} = 0.30 \text{ GeV}/c \quad 4\pi\lambda^2 = 94.8 \text{ mb}$$

$\text{Re}(\text{pole position}) = 1209$ to 1211 (≈ 1210) MeV

$-2\text{Im}(\text{pole position}) = 98$ to 102 (≈ 100) MeV

$\Delta(1232)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\pi$	>99 %	227
$N\gamma$	0.52–0.60 %	259
$N\gamma$, helicity=1/2	0.11–0.13 %	259
$N\gamma$, helicity=3/2	0.41–0.47 %	259

$\Delta(1600)$ P_{33}

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Breit-Wigner mass = 1550 to 1700 (≈ 1600) MeV

Breit-Wigner full width = 250 to 450 (≈ 350) MeV

$$p_{\text{beam}} = 0.87 \text{ GeV}/c \quad 4\pi\lambda^2 = 18.6 \text{ mb}$$

$\text{Re}(\text{pole position}) = 1500$ to 1700 (≈ 1600) MeV

$-2\text{Im}(\text{pole position}) = 200$ to 400 (≈ 300) MeV

$\Delta(1600)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\pi$	10–25 %	512
$N\pi\pi$	75–90 %	473
$\Delta\pi$	40–70 %	301
$N\rho$	<25 %	†
$N(1440)\pi$	10–35 %	74
$N\gamma$	0.001–0.02 %	525
$N\gamma$, helicity=1/2	0.0–0.02 %	525
$N\gamma$, helicity=3/2	0.001–0.005 %	525

$\Delta(1620)$ S_{31}

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$$

Breit-Wigner mass = 1615 to 1675 (≈ 1620) MeV
 Breit-Wigner full width = 120 to 180 (≈ 150) MeV
 $p_{\text{beam}} = 0.91 \text{ GeV}/c$ $4\pi\lambda^2 = 17.7 \text{ mb}$
 $\text{Re}(\text{pole position}) = 1580 \text{ to } 1620 \text{ (≈ 1600) MeV}$
 $-2\text{Im}(\text{pole position}) = 100 \text{ to } 130 \text{ (≈ 115) MeV}$

$\Delta(1620)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	20–30 %	526
$N\pi\pi$	70–80 %	488
$\Delta\pi$	30–60 %	318
$N\rho$	7–25 %	†
$N\gamma$	0.004–0.044 %	538
$N\gamma$, helicity=1/2	0.004–0.044 %	538

 $\Delta(1700)$ D_{33}

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-)$$

Breit-Wigner mass = 1670 to 1770 (≈ 1700) MeV
 Breit-Wigner full width = 200 to 400 (≈ 300) MeV
 $p_{\text{beam}} = 1.05 \text{ GeV}/c$ $4\pi\lambda^2 = 14.5 \text{ mb}$
 $\text{Re}(\text{pole position}) = 1620 \text{ to } 1700 \text{ (≈ 1660) MeV}$
 $-2\text{Im}(\text{pole position}) = 150 \text{ to } 250 \text{ (≈ 200) MeV}$

$\Delta(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	10–20 %	580
$N\pi\pi$	80–90 %	547
$\Delta\pi$	30–60 %	385
$N\rho$	30–55 %	†
$N\gamma$	0.12–0.26 %	591
$N\gamma$, helicity=1/2	0.08–0.16 %	591
$N\gamma$, helicity=3/2	0.025–0.12 %	591

 $\Delta(1905)$ F_{35}

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^+)$$

Breit-Wigner mass = 1870 to 1920 (≈ 1905) MeV
 Breit-Wigner full width = 280 to 440 (≈ 350) MeV
 $p_{\text{beam}} = 1.45 \text{ GeV}/c$ $4\pi\lambda^2 = 9.62 \text{ mb}$
 $\text{Re}(\text{pole position}) = 1800 \text{ to } 1860 \text{ (≈ 1830) MeV}$
 $-2\text{Im}(\text{pole position}) = 230 \text{ to } 330 \text{ (≈ 280) MeV}$

$\Delta(1905)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	5–15 %	713
$N\pi\pi$	85–95 %	687
$\Delta\pi$	<25 %	542
$N\rho$	>60 %	421
$N\gamma$	0.01–0.03 %	721
$N\gamma$, helicity=1/2	0.0–0.1 %	721
$N\gamma$, helicity=3/2	0.004–0.03 %	721

$\Delta(1910) P_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$$

Breit-Wigner mass = 1870 to 1920 (≈ 1910) MeV
 Breit-Wigner full width = 190 to 270 (≈ 250) MeV
 $p_{\text{beam}} = 1.46 \text{ GeV}/c \quad 4\pi\lambda^2 = 9.54 \text{ mb}$
 Re(pole position) = 1830 to 1880 (≈ 1855) MeV
 $-2\text{Im}(\text{pole position}) = 200 \text{ to } 500 (\approx 350) \text{ MeV}$

$\Delta(1910)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	15–30 %	716
$N\gamma$	0.0–0.2 %	725
$N\gamma$, helicity=1/2	0.0–0.2 %	725

$\Delta(1920) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Breit-Wigner mass = 1900 to 1970 (≈ 1920) MeV
 Breit-Wigner full width = 150 to 300 (≈ 200) MeV
 $p_{\text{beam}} = 1.48 \text{ GeV}/c \quad 4\pi\lambda^2 = 9.37 \text{ mb}$
 Re(pole position) = 1850 to 1950 (≈ 1900) MeV
 $-2\text{Im}(\text{pole position}) = 200 \text{ to } 400 (\approx 300) \text{ MeV}$

$\Delta(1920)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	5–20 %	722

$\Delta(1930) D_{35}$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^-)$$

Breit-Wigner mass = 1920 to 1970 (≈ 1930) MeV
 Breit-Wigner full width = 250 to 450 (≈ 350) MeV
 $p_{\text{beam}} = 1.50 \text{ GeV}/c \quad 4\pi\lambda^2 = 9.21 \text{ mb}$
 Re(pole position) = 1840 to 1940 (≈ 1890) MeV
 $-2\text{Im}(\text{pole position}) = 200 \text{ to } 300 (\approx 250) \text{ MeV}$

$\Delta(1930)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	10–20 %	729
$N\gamma$	0.0–0.02 %	737
$N\gamma$, helicity=1/2	0.0–0.01 %	737
$N\gamma$, helicity=3/2	0.0–0.01 %	737

$\Delta(1950)$ F_{37}

$$I(J^P) = \frac{3}{2}(\frac{7}{2}+)$$

Breit-Wigner mass = 1940 to 1960 (≈ 1950) MeV

Breit-Wigner full width = 290 to 350 (≈ 300) MeV

$$p_{\text{beam}} = 1.54 \text{ GeV}/c \quad 4\pi\lambda^2 = 8.91 \text{ mb}$$

$\text{Re}(\text{pole position}) = 1880$ to 1890 (≈ 1885) MeV

$-2\text{Im}(\text{pole position}) = 210$ to 270 (≈ 240) MeV

$\Delta(1950)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	35–40 %	741
$N\pi\pi$		716
$\Delta\pi$	20–30 %	574
$N\rho$	<10 %	469
$N\gamma$	0.08–0.13 %	749
$N\gamma$, helicity=1/2	0.03–0.055 %	749
$N\gamma$, helicity=3/2	0.05–0.075 %	749

$\Delta(2420)$ $H_{3,11}$

$$I(J^P) = \frac{3}{2}(\frac{11}{2}+)$$

Breit-Wigner mass = 2300 to 2500 (≈ 2420) MeV

Breit-Wigner full width = 300 to 500 (≈ 400) MeV

$$p_{\text{beam}} = 2.64 \text{ GeV}/c \quad 4\pi\lambda^2 = 4.68 \text{ mb}$$

$\text{Re}(\text{pole position}) = 2260$ to 2400 (≈ 2330) MeV

$-2\text{Im}(\text{pole position}) = 350$ to 750 (≈ 550) MeV

$\Delta(2420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\pi$	5–15 %	1023

Λ BARYONS ($S = -1, I = 0$)

$$\Lambda^0 = uds$$

Λ

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 1115.683 \pm 0.006$ MeV

$(m_\Lambda - m_{\bar{\Lambda}}) / m_\Lambda = (-0.1 \pm 1.1) \times 10^{-5}$ ($S = 1.6$)

Mean life $\tau = (2.632 \pm 0.020) \times 10^{-10}$ s ($S = 1.6$)

$c\tau = 7.89$ cm

Magnetic moment $\mu = -0.613 \pm 0.004$ μ_N

Electric dipole moment $d < 1.5 \times 10^{-16}$ e cm, CL = 95%

Decay parameters

$p\pi^-$	$\alpha_- = 0.642 \pm 0.013$
"	$\phi_- = (-6.5 \pm 3.5)^\circ$
"	$\gamma_- = 0.76$ [h]
"	$\Delta_- = (8 \pm 4)^\circ$ [h]
$n\pi^0$	$\alpha_0 = +0.65 \pm 0.05$
$pe^-\bar{\nu}_e$	$g_A/g_V = -0.718 \pm 0.015$ [f]

Λ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$p\pi^-$	(63.9 ± 0.5) %	101
$n\pi^0$	(35.8 ± 0.5) %	104
$n\gamma$	(1.75 ± 0.15) $\times 10^{-3}$	162
$p\pi^-\gamma$	[i] (8.4 ± 1.4) $\times 10^{-4}$	101
$pe^-\bar{\nu}_e$	(8.32 ± 0.14) $\times 10^{-4}$	163
$p\mu^-\bar{\nu}_\mu$	(1.57 ± 0.35) $\times 10^{-4}$	131

$\Lambda(1405)$ S_{01}

$$I(J^P) = 0(\frac{1}{2}^-)$$

Mass $m = 1406 \pm 4$ MeV

Full width $\Gamma = 50.0 \pm 2.0$ MeV

Below $\bar{K}N$ threshold

$\Lambda(1405)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma\pi$	100 %	152

$\Lambda(1520)$ D_{03}

$$I(J^P) = 0(\frac{3}{2}^-)$$

Mass $m = 1519.5 \pm 1.0$ MeV [j]

Full width $\Gamma = 15.6 \pm 1.0$ MeV [j]

$$p_{\text{beam}} = 0.39 \text{ GeV}/c \quad 4\pi\lambda^2 = 82.8 \text{ mb}$$

$\Lambda(1520)$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	$45 \pm 1\%$	244
$\Sigma\pi$	$42 \pm 1\%$	267
$\Lambda\pi\pi$	$10 \pm 1\%$	252
$\Sigma\pi\pi$	$0.9 \pm 0.1\%$	152
$\Lambda\gamma$	$0.8 \pm 0.2\%$	351

$\Lambda(1600)$ P_{01}

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 1560$ to 1700 (≈ 1600) MeV

Full width $\Gamma = 50$ to 250 (≈ 150) MeV

$$p_{\text{beam}} = 0.58 \text{ GeV}/c \quad 4\pi\lambda^2 = 41.6 \text{ mb}$$

$\Lambda(1600)$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	15–30 %	343
$\Sigma\pi$	10–60 %	336

$\Lambda(1670)$ S_{01}

$$I(J^P) = 0(\frac{1}{2}^-)$$

Mass $m = 1660$ to 1680 (≈ 1670) MeV

Full width $\Gamma = 25$ to 50 (≈ 35) MeV

$$p_{\text{beam}} = 0.74 \text{ GeV}/c \quad 4\pi\lambda^2 = 28.5 \text{ mb}$$

$\Lambda(1670)$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	15–25 %	414
$\Sigma\pi$	20–60 %	393
$\Lambda\eta$	15–35 %	64

$\Lambda(1690)$ D_{03}

$$I(J^P) = 0(\frac{3}{2}^-)$$

Mass $m = 1685$ to 1695 (≈ 1690) MeV

Full width $\Gamma = 50$ to 70 (≈ 60) MeV

$$p_{\text{beam}} = 0.78 \text{ GeV}/c \quad 4\pi\lambda^2 = 26.1 \text{ mb}$$

$\Lambda(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	20–30 %	433
$\Sigma\pi$	20–40 %	409
$\Lambda\pi\pi$	~ 25 %	415
$\Sigma\pi\pi$	~ 20 %	350

$\Lambda(1800)$ S_{01}

$$I(J^P) = 0(\frac{1}{2}^-)$$

Mass $m = 1720$ to 1850 (≈ 1800) MeV

Full width $\Gamma = 200$ to 400 (≈ 300) MeV

$$p_{\text{beam}} = 1.01 \text{ GeV}/c \quad 4\pi\lambda^2 = 17.5 \text{ mb}$$

$\Lambda(1800)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	25–40 %	528
$\Sigma\pi$	seen	493
$\Sigma(1385)\pi$	seen	345
$N\bar{K}^*(892)$	seen	†

$\Lambda(1810)$ P_{01}

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 1750$ to 1850 (≈ 1810) MeV

Full width $\Gamma = 50$ to 250 (≈ 150) MeV

$$p_{\text{beam}} = 1.04 \text{ GeV}/c \quad 4\pi\lambda^2 = 17.0 \text{ mb}$$

$\Lambda(1810)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	20–50 %	537
$\Sigma\pi$	10–40 %	501
$\Sigma(1385)\pi$	seen	356
$N\bar{K}^*(892)$	30–60 %	†

$\Lambda(1820)$ F_{05}

$$I(J^P) = 0(\frac{5}{2}^+)$$

Mass $m = 1815$ to 1825 (≈ 1820) MeV

Full width $\Gamma = 70$ to 90 (≈ 80) MeV

$$p_{\text{beam}} = 1.06 \text{ GeV}/c \quad 4\pi\lambda^2 = 16.5 \text{ mb}$$

$\Lambda(1820)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	55–65 %	545
$\Sigma\pi$	8–14 %	508
$\Sigma(1385)\pi$	5–10 %	362

$\Lambda(1830)$ D_{05}

$$I(J^P) = 0(\frac{5}{2}^-)$$

Mass $m = 1810$ to 1830 (≈ 1830) MeV

Full width $\Gamma = 60$ to 110 (≈ 95) MeV

$$p_{\text{beam}} = 1.08 \text{ GeV}/c \quad 4\pi\lambda^2 = 16.0 \text{ mb}$$

$\Lambda(1830)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	3–10 %	553
$\Sigma\pi$	35–75 %	515
$\Sigma(1385)\pi$	>15 %	371

$\Lambda(1890)$ P_{03}

$$I(J^P) = 0(\frac{3}{2}^+)$$

Mass $m = 1850$ to 1910 (≈ 1890) MeV

Full width $\Gamma = 60$ to 200 (≈ 100) MeV

$$p_{\text{beam}} = 1.21 \text{ GeV}/c \quad 4\pi\lambda^2 = 13.6 \text{ mb}$$

$\Lambda(1890)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	20–35 %	599
$\Sigma\pi$	3–10 %	559
$\Sigma(1385)\pi$	seen	420
$N\bar{K}^*(892)$	seen	233

$\Lambda(2100)$ G_{07}

$$I(J^P) = 0(\frac{7}{2}^-)$$

Mass $m = 2090$ to 2110 (≈ 2100) MeV

Full width $\Gamma = 100$ to 250 (≈ 200) MeV

$$p_{\text{beam}} = 1.68 \text{ GeV}/c \quad 4\pi\lambda^2 = 8.68 \text{ mb}$$

$\Lambda(2100)$ DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\bar{K}$	25–35 %	751
$\Sigma\pi$	~ 5 %	704
$\Lambda\eta$	< 3 %	617
ΞK	< 3 %	483
$\Lambda\omega$	< 8 %	443
$N\bar{K}^*(892)$	10–20 %	514

$\Lambda(2110)$ F_{05}

$$I(J^P) = 0(\frac{5}{2}^+)$$

Mass $m = 2090$ to 2140 (≈ 2110) MeV

Full width $\Gamma = 150$ to 250 (≈ 200) MeV

$$p_{\text{beam}} = 1.70 \text{ GeV}/c \quad 4\pi\lambda^2 = 8.53 \text{ mb}$$

$\Lambda(2110)$ DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\bar{K}$	5–25 %	757
$\Sigma\pi$	10–40 %	711
$\Lambda\omega$	seen	455
$\Sigma(1385)\pi$	seen	589
$N\bar{K}^*(892)$	10–60 %	524

$\Lambda(2350)$ H_{09}

$$I(J^P) = 0(\frac{9}{2}^+)$$

Mass $m = 2340$ to 2370 (≈ 2350) MeV

Full width $\Gamma = 100$ to 250 (≈ 150) MeV

$$p_{\text{beam}} = 2.29 \text{ GeV}/c \quad 4\pi\lambda^2 = 5.85 \text{ mb}$$

$\Lambda(2350)$ DECAY MODES

$$\text{Fraction } (\Gamma_i/\Gamma)$$

$$p \text{ (MeV}/c)$$

$N\bar{K}$	~ 12 %	915
$\Sigma\pi$	~ 10 %	867

Σ BARYONS ($S = -1, I = 1$)

$$\Sigma^+ = uus, \quad \Sigma^0 = uds, \quad \Sigma^- = dds$$

Σ^+

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1189.37 \pm 0.07$ MeV ($S = 2.2$)

Mean life $\tau = (0.8018 \pm 0.0026) \times 10^{-10}$ s

$$c\tau = 2.404 \text{ cm}$$

$$(\tau_{\Sigma^+} - \tau_{\bar{\Sigma}^-}) / \tau_{\Sigma^+} = (-0.6 \pm 1.2) \times 10^{-3}$$

Magnetic moment $\mu = 2.458 \pm 0.010 \mu_N$ ($S = 2.1$)

$$\Gamma(\Sigma^+ \rightarrow n\ell^+\nu)/\Gamma(\Sigma^- \rightarrow n\ell^-\bar{\nu}) < 0.043$$

Decay parameters

$p\pi^0$	$\alpha_0 = -0.980^{+0.017}_{-0.015}$
"	$\phi_0 = (36 \pm 34)^\circ$
"	$\gamma_0 = 0.16$ [h]
"	$\Delta_0 = (187 \pm 6)^\circ$ [h]
$n\pi^+$	$\alpha_+ = 0.068 \pm 0.013$
"	$\phi_+ = (167 \pm 20)^\circ$ ($S = 1.1$)
"	$\gamma_+ = -0.97$ [h]
"	$\Delta_+ = (-73^{+133}_{-10})^\circ$ [h]
$p\gamma$	$\alpha_\gamma = -0.76 \pm 0.08$

Σ^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$p\pi^0$	$(51.57 \pm 0.30) \%$		189
$n\pi^+$	$(48.31 \pm 0.30) \%$		185
$p\gamma$	$(1.23 \pm 0.05) \times 10^{-3}$		225
$n\pi^+\gamma$	[i] $(4.5 \pm 0.5) \times 10^{-4}$		185
$\Lambda e^+ \nu_e$	$(2.0 \pm 0.5) \times 10^{-5}$		71

$\Delta S = \Delta Q$ (*SQ*) violating modes or
 $\Delta S = 1$ weak neutral current (*S1*) modes

$ne^+\nu_e$	<i>SQ</i>	< 5	$\times 10^{-6}$	90%	224
$n\mu^+\nu_\mu$	<i>SQ</i>	< 3.0	$\times 10^{-5}$	90%	202
pe^+e^-	<i>S1</i>	< 7	$\times 10^{-6}$		225

Σ^0

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1192.642 \pm 0.024$ MeV

$m_{\Sigma^-} - m_{\Sigma^0} = 4.807 \pm 0.035$ MeV ($S = 1.1$)

$m_{\Sigma^0} - m_\Lambda = 76.959 \pm 0.023$ MeV

Mean life $\tau = (7.4 \pm 0.7) \times 10^{-20}$ s

$c\tau = 2.22 \times 10^{-11}$ m

Transition magnetic moment $|\mu_{\Sigma\Lambda}| = 1.61 \pm 0.08$ μ_N

Σ^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda\gamma$	100 %		74
$\Lambda\gamma\gamma$	< 3 %	90%	74
$\Lambda e^+ e^-$	[k] 5×10^{-3}		74

Σ^-

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1197.449 \pm 0.030$ MeV ($S = 1.2$)

$m_{\Sigma^-} - m_{\Sigma^+} = 8.08 \pm 0.08$ MeV ($S = 1.9$)

$m_{\Sigma^-} - m_\Lambda = 81.766 \pm 0.030$ MeV ($S = 1.2$)

Mean life $\tau = (1.479 \pm 0.011) \times 10^{-10}$ s ($S = 1.3$)

$c\tau = 4.434$ cm

Magnetic moment $\mu = -1.160 \pm 0.025$ μ_N ($S = 1.7$)

Decay parameters

$n\pi^-$	$\alpha_- = -0.068 \pm 0.008$
"	$\phi_- = (10 \pm 15)^\circ$
"	$\gamma_- = 0.98$ [h]
"	$\Delta_- = (249^{+12}_{-120})^\circ$ [h]
$n e^- \bar{\nu}_e$	$g_A/g_V = 0.340 \pm 0.017$ [f]
"	$f_2(0)/f_1(0) = 0.97 \pm 0.14$
"	$D = 0.11 \pm 0.10$
$\Lambda e^- \bar{\nu}_e$	$g_V/g_A = 0.01 \pm 0.10$ ($S = 1.5$)
"	$g_{WM}/g_A = 2.4 \pm 1.7$ [f]

Σ^- DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$n\pi^-$	$(99.848 \pm 0.005) \%$	193
$n\pi^- \gamma$	$[i] (4.6 \pm 0.6) \times 10^{-4}$	193
$n e^- \bar{\nu}_e$	$(1.017 \pm 0.034) \times 10^{-3}$	230
$n \mu^- \bar{\nu}_\mu$	$(4.5 \pm 0.4) \times 10^{-4}$	210
$\Lambda e^- \bar{\nu}_e$	$(5.73 \pm 0.27) \times 10^{-5}$	79

$\Sigma(1385)$ P_{13}

$$I(J^P) = 1(\frac{3}{2}^+)$$

$\Sigma(1385)^+$ mass $m = 1382.8 \pm 0.4$ MeV ($S = 2.0$)

$\Sigma(1385)^0$ mass $m = 1383.7 \pm 1.0$ MeV ($S = 1.4$)

$\Sigma(1385)^-$ mass $m = 1387.2 \pm 0.5$ MeV ($S = 2.2$)

$\Sigma(1385)^+$ full width $\Gamma = 35.8 \pm 0.8$ MeV

$\Sigma(1385)^0$ full width $\Gamma = 36 \pm 5$ MeV

$\Sigma(1385)^-$ full width $\Gamma = 39.4 \pm 2.1$ MeV ($S = 1.7$)

Below $\bar{K}N$ threshold

$\Sigma(1385)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$\Lambda\pi$

88 ± 2 %

208

$\Sigma\pi$

12 ± 2 %

127

$\Sigma(1660)$ P_{11}

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1630$ to 1690 (≈ 1660) MeV

Full width $\Gamma = 40$ to 200 (≈ 100) MeV

$p_{\text{beam}} = 0.72$ GeV/c $4\pi\lambda^2 = 29.9$ mb

$\Sigma(1660)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\bar{K}$

10–30 %

405

$\Lambda\pi$

seen

439

$\Sigma\pi$

seen

385

$\Sigma(1670)$ D_{13}

$$I(J^P) = 1(\frac{3}{2}^-)$$

Mass $m = 1665$ to 1685 (≈ 1670) MeV

Full width $\Gamma = 40$ to 80 (≈ 60) MeV

$p_{\text{beam}} = 0.74$ GeV/c $4\pi\lambda^2 = 28.5$ mb

$\Sigma(1670)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\bar{K}$

7–13 %

414

$\Lambda\pi$

5–15 %

447

$\Sigma\pi$

30–60 %

393

$\Sigma(1750)$ S_{11}

$$I(J^P) = 1(\frac{1}{2}^-)$$

Mass $m = 1730$ to 1800 (≈ 1750) MeV

Full width $\Gamma = 60$ to 160 (≈ 90) MeV

$$p_{\text{beam}} = 0.91 \text{ GeV}/c \quad 4\pi\lambda^2 = 20.7 \text{ mb}$$

$\Sigma(1750)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\bar{K}$	10–40 %	486
$\Lambda\pi$	seen	507
$\Sigma\pi$	<8 %	455
$\Sigma\eta$	15–55 %	81

$\Sigma(1775)$ D_{15}

$$I(J^P) = 1(\frac{5}{2}^-)$$

Mass $m = 1770$ to 1780 (≈ 1775) MeV

Full width $\Gamma = 105$ to 135 (≈ 120) MeV

$$p_{\text{beam}} = 0.96 \text{ GeV}/c \quad 4\pi\lambda^2 = 19.0 \text{ mb}$$

$\Sigma(1775)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\bar{K}$	37–43%	508
$\Lambda\pi$	14–20%	525
$\Sigma\pi$	2–5%	474
$\Sigma(1385)\pi$	8–12%	324
$\Lambda(1520)\pi$	17–23%	198

$\Sigma(1915)$ F_{15}

$$I(J^P) = 1(\frac{5}{2}^+)$$

Mass $m = 1900$ to 1935 (≈ 1915) MeV

Full width $\Gamma = 80$ to 160 (≈ 120) MeV

$$p_{\text{beam}} = 1.26 \text{ GeV}/c \quad 4\pi\lambda^2 = 12.8 \text{ mb}$$

$\Sigma(1915)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$N\bar{K}$	5–15 %	618
$\Lambda\pi$	seen	622
$\Sigma\pi$	seen	577
$\Sigma(1385)\pi$	<5 %	440

$\Sigma(1940)$ D_{13}

$$I(J^P) = 1(\frac{3}{2}^-)$$

Mass $m = 1900$ to 1950 (≈ 1940) MeV

Full width $\Gamma = 150$ to 300 (≈ 220) MeV

$$p_{\text{beam}} = 1.32 \text{ GeV}/c \quad 4\pi\lambda^2 = 12.1 \text{ mb}$$

$\Sigma(1940)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	<20 %	637
$\Lambda\pi$	seen	639
$\Sigma\pi$	seen	594
$\Sigma(1385)\pi$	seen	460
$\Lambda(1520)\pi$	seen	354
$\Delta(1232)\bar{K}$	seen	410
$N\bar{K}^*(892)$	seen	320

$\Sigma(2030)$ F_{17}

$$I(J^P) = 1(\frac{7}{2}^+)$$

Mass $m = 2025$ to 2040 (≈ 2030) MeV

Full width $\Gamma = 150$ to 200 (≈ 180) MeV

$$p_{\text{beam}} = 1.52 \text{ GeV}/c \quad 4\pi\lambda^2 = 9.93 \text{ mb}$$

$\Sigma(2030)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	17–23 %	702
$\Lambda\pi$	17–23 %	700
$\Sigma\pi$	5–10 %	657
ΞK	<2 %	412
$\Sigma(1385)\pi$	5–15 %	529
$\Lambda(1520)\pi$	10–20 %	430
$\Delta(1232)\bar{K}$	10–20 %	498
$N\bar{K}^*(892)$	<5 %	438

$\Sigma(2250)$

$$I(J^P) = 1(?^?)$$

Mass $m = 2210$ to 2280 (≈ 2250) MeV

Full width $\Gamma = 60$ to 150 (≈ 100) MeV

$$p_{\text{beam}} = 2.04 \text{ GeV}/c \quad 4\pi\lambda^2 = 6.76 \text{ mb}$$

$\Sigma(2250)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	<10 %	851
$\Lambda\pi$	seen	842
$\Sigma\pi$	seen	803

Ξ BARYONS ($S = -2, I = 1/2$)

$$\Xi^0 = uss, \quad \Xi^- = dss$$

Ξ^0

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

Mass $m = 1314.83 \pm 0.20$ MeV

$m_{\Xi^-} - m_{\Xi^0} = 6.48 \pm 0.24$ MeV

Mean life $\tau = (2.90 \pm 0.09) \times 10^{-10}$ s

$c\tau = 8.71$ cm

Magnetic moment $\mu = -1.250 \pm 0.014$ μ_N

Decay parameters

$$\Lambda\pi^0 \quad \alpha = -0.411 \pm 0.022 \quad (S = 2.1)$$

$$\text{"} \quad \phi = (21 \pm 12)^\circ$$

$$\text{"} \quad \gamma = 0.85 [h]$$

$$\text{"} \quad \Delta = (218^{+12}_{-19})^\circ [h]$$

$$\Lambda\gamma \quad \alpha = 0.4 \pm 0.4$$

$$\Sigma^0\gamma \quad \alpha = 0.20 \pm 0.32$$

Ξ^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\Lambda\pi^0$	$(99.51 \pm 0.05) \%$	$S=1.2$	135
$\Lambda\gamma$	$(1.18 \pm 0.30) \times 10^{-3}$	$S=2.0$	184
$\Sigma^0\gamma$	$(3.5 \pm 0.4) \times 10^{-3}$		117
$\Sigma^+ e^- \bar{\nu}_e$	$(2.7 \pm 0.4) \times 10^{-4}$		120
$\Sigma^+ \mu^- \bar{\nu}_\mu$	$< 1.1 \times 10^{-3}$	CL=90%	64

$\Delta S = \Delta Q$ (SQ) violating modes or $\Delta S = 2$ forbidden (S2) modes

$\Sigma^- e^+ \nu_e$	SQ	< 9	$\times 10^{-4}$	CL=90%	112
$\Sigma^- \mu^+ \nu_\mu$	SQ	< 9	$\times 10^{-4}$	CL=90%	49
$p\pi^-$	S2	< 4	$\times 10^{-5}$	CL=90%	299
$p e^- \bar{\nu}_e$	S2	< 1.3	$\times 10^{-3}$		323
$p \mu^- \bar{\nu}_\mu$	S2	< 1.3	$\times 10^{-3}$		309



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

Mass $m = 1321.31 \pm 0.13$ MeV

Mean life $\tau = (1.639 \pm 0.015) \times 10^{-10}$ s

$$c\tau = 4.91$$
 cm

Magnetic moment $\mu = -0.6507 \pm 0.0025$ μ_N

Decay parameters

$$\Lambda\pi^- \quad \alpha = -0.456 \pm 0.014 \quad (S = 1.8)$$

$$" \quad \phi = (4 \pm 4)^\circ$$

$$" \quad \gamma = 0.89 [h]$$

$$" \quad \Delta = (188 \pm 8)^\circ [h]$$

$$\Lambda e^- \bar{\nu}_e \quad g_A/g_V = -0.25 \pm 0.05 [f]$$

Ξ^- DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda\pi^-$	$(99.887 \pm 0.035) \%$		139
$\Sigma^-\gamma$	$(1.27 \pm 0.23) \times 10^{-4}$		118
$\Lambda e^- \bar{\nu}_e$	$(5.63 \pm 0.31) \times 10^{-4}$		190
$\Lambda\mu^- \bar{\nu}_\mu$	$(3.5 \pm 3.5) \times 10^{-4}$		163
$\Sigma^0 e^- \bar{\nu}_e$	$(8.7 \pm 1.7) \times 10^{-5}$		122
$\Sigma^0 \mu^- \bar{\nu}_\mu$	$< 8 \times 10^{-4}$	90%	70
$\Xi^0 e^- \bar{\nu}_e$	$< 2.3 \times 10^{-3}$	90%	6

$\Delta S = 2$ forbidden (S2) modes

$n\pi^-$	S2	< 1.9	$\times 10^{-5}$	90%	303
$n e^- \bar{\nu}_e$	S2	< 3.2	$\times 10^{-3}$	90%	327
$n\mu^- \bar{\nu}_\mu$	S2	< 1.5	%	90%	314
$p\pi^-\pi^-$	S2	< 4	$\times 10^{-4}$	90%	223
$p\pi^- e^- \bar{\nu}_e$	S2	< 4	$\times 10^{-4}$	90%	304
$p\pi^- \mu^- \bar{\nu}_\mu$	S2	< 4	$\times 10^{-4}$	90%	250
$p\mu^- \mu^-$	L	< 4	$\times 10^{-4}$	90%	272

$\Xi(1530) P_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}+)$$

$\Xi(1530)^0$ mass $m = 1531.80 \pm 0.32$ MeV ($S = 1.3$)

$\Xi(1530)^-$ mass $m = 1535.0 \pm 0.6$ MeV

$\Xi(1530)^0$ full width $\Gamma = 9.1 \pm 0.5$ MeV

$\Xi(1530)^-$ full width $\Gamma = 9.9^{+1.7}_{-1.9}$ MeV

$\Xi(1530)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Xi\pi$	100 %		152
$\Xi\gamma$	<4 %	90%	200

$\Xi(1690)$

$$I(J^P) = \frac{1}{2}(??)$$

Mass $m = 1690 \pm 10$ MeV [j]

Full width $\Gamma < 30$ MeV

$\Xi(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda\bar{K}$	seen	240
$\Sigma\bar{K}$	seen	51
$\Xi\pi$	seen	—
$\Xi^-\pi^+\pi^-$	possibly seen	214

$\Xi(1820) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}-)$$

Mass $m = 1823 \pm 5$ MeV [j]

Full width $\Gamma = 24^{+15}_{-10}$ MeV [j]

$\Xi(1820)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda\bar{K}$	large	400
$\Sigma\bar{K}$	small	320
$\Xi\pi$	small	413
$\Xi(1530)\pi$	small	234

$\Xi(1950)$

$$I(J^P) = \frac{1}{2}(??)$$

Mass $m = 1950 \pm 15$ MeV [j]
 Full width $\Gamma = 60 \pm 20$ MeV [j]

$\Xi(1950)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda\bar{K}$	seen	522
$\Sigma\bar{K}$	possibly seen	460
$\Xi\pi$	seen	518

$\Xi(2030)$

$$I(J^P) = \frac{1}{2}(\geq \frac{5}{2}?)$$

Mass $m = 2025 \pm 5$ MeV [j]
 Full width $\Gamma = 20^{+15}_{-5}$ MeV [j]

$\Xi(2030)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda\bar{K}$	$\sim 20\%$	589
$\Sigma\bar{K}$	$\sim 80\%$	533
$\Xi\pi$	small	573
$\Xi(1530)\pi$	small	421
$\Lambda\bar{K}\pi$	small	501
$\Sigma\bar{K}\pi$	small	430

Ω BARYONS ($S = -3, I = 0$)

$$\Omega^- = sss$$

Ω^-

$$I(J^P) = 0(\frac{3}{2}^+)$$

J^P is not yet measured; $\frac{3}{2}^+$ is the quark model prediction.

Mass $m = 1672.45 \pm 0.29$ MeV

$$(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\Omega^-} = (-1 \pm 8) \times 10^{-5}$$

$$\text{Mean life } \tau = (0.821 \pm 0.011) \times 10^{-10} \text{ s}$$

$$c\tau = 2.461 \text{ cm}$$

$$(\tau_{\Omega^-} - \tau_{\bar{\Omega}^+}) / \tau_{\Omega^-} = -0.002 \pm 0.040$$

$$\text{Magnetic moment } \mu = -2.02 \pm 0.05 \mu_N$$

Decay parameters

$$\Lambda K^- \quad \alpha = -0.026 \pm 0.023$$

$$\frac{1}{2}[\alpha(\Lambda K^-) + \alpha(\bar{\Lambda} K^+)] = -0.004 \pm 0.040$$

$$\Xi^0 \pi^- \quad \alpha = 0.09 \pm 0.14$$

$$\Xi^- \pi^0 \quad \alpha = 0.05 \pm 0.21$$

Ω^- DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
ΛK^-	(67.8 ± 0.7) %		211
$\Xi^0 \pi^-$	(23.6 ± 0.7) %		294
$\Xi^- \pi^0$	(8.6 ± 0.4) %		290
$\Xi^- \pi^+ \pi^-$	($4.3^{+3.4}_{-1.3}$) $\times 10^{-4}$		190
$\Xi(1530)^0 \pi^-$	($6.4^{+5.1}_{-2.0}$) $\times 10^{-4}$		17
$\Xi^0 e^- \bar{\nu}_e$	(5.6 ± 2.8) $\times 10^{-3}$		319
$\Xi^- \gamma$	< 4.6 $\times 10^{-4}$	90%	314
$\Delta S = 2$ forbidden (S2) modes			
$\Lambda \pi^-$	$S2 < 1.9 \times 10^{-4}$	90%	449

$\Omega(2250)^-$

$I(J^P) = 0(?^?)$

Mass $m = 2252 \pm 9$ MeV

Full width $\Gamma = 55 \pm 18$ MeV

$\Omega(2250)^-$ DECAY MODES

	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi^- \pi^+ K^-$	seen	531
$\Xi(1530)^0 K^-$	seen	437

CHARMED BARYONS ($C=+1$)

$$\begin{aligned}\Lambda_c^+ &= u d c, \quad \Sigma_c^{++} = u u c, \quad \Sigma_c^+ = u d c, \quad \Sigma_c^0 = d d c, \\ \Xi_c^+ &= u s c, \quad \Xi_c^0 = d s c, \quad \Omega_c^0 = s s c\end{aligned}$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

J is not well measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2284.9 \pm 0.6$ MeV

Mean life $\tau = (0.206 \pm 0.012) \times 10^{-12}$ s

$c\tau = 61.8 \mu\text{m}$

Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.98 \pm 0.19$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.82^{+0.11}_{-0.07}$$

Nearly all branching fractions of the Λ_c^+ are measured relative to the $pK^-\pi^+$ mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$ in a Note at the beginning of the branching-ratio measurements in the Listings. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

Λ_c^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level (MeV/c)
Hadronic modes with a p and one \bar{K}		
$p\bar{K}^0$	(2.3 \pm 0.6) %	872
$pK^-\pi^+$	[<i>I</i>] (5.0 \pm 1.3) %	822
$p\bar{K}^*(892)^0$	[<i>m</i>] (1.6 \pm 0.5) %	681
$\Delta(1232)^{++}K^-$	(8.6 \pm 3.0) $\times 10^{-3}$	709
$\Lambda(1520)\pi^+$	[<i>m</i>] (5.9 \pm 2.1) $\times 10^{-3}$	626
$pK^-\pi^+$ nonresonant	(2.8 \pm 0.8) %	822
$p\bar{K}^0\pi^0$	(3.3 \pm 1.0) %	822
$p\bar{K}^0\eta$	(1.2 \pm 0.4) %	567

$p\bar{K}^0\pi^+\pi^-$	(2.6 ± 0.7) %	753
$pK^-\pi^+\pi^0$	(3.4 ± 1.0) %	758
$pK^*(892)^-\pi^+$	[m] (1.1 ± 0.5) %	579
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	(3.6 ± 1.2) %	758
$\Delta(1232)\bar{K}^*(892)$	seen	416
$pK^-\pi^+\pi^+\pi^-$	(1.1 ± 0.8) × 10 ⁻³	670
$pK^-\pi^+\pi^0\pi^0$	(8 ± 4) × 10 ⁻³	676
$pK^-\pi^+\pi^0\pi^0\pi^0$	(5.0 ± 3.4) × 10 ⁻³	573

Hadronic modes with a p and zero or two K 's

$p\pi^+\pi^-$	(3.5 ± 2.0) × 10 ⁻³	926
$p f_0(980)$	[m] (2.8 ± 1.9) × 10 ⁻³	621
$p\pi^+\pi^+\pi^-\pi^-$	(1.8 ± 1.2) × 10 ⁻³	851
pK^+K^-	(2.3 ± 0.9) × 10 ⁻³	615
$p\phi$	[m] (1.2 ± 0.5) × 10 ⁻³	589

Hadronic modes with a hyperon

$\Lambda\pi^+$	(9.0 ± 2.8) × 10 ⁻³	863
$\Lambda\pi^+\pi^0$	(3.6 ± 1.3) %	843
$\Lambda\rho^+$	< 5 %	CL=95% 638
$\Lambda\pi^+\pi^+\pi^-$	(3.3 ± 1.0) %	806
$\Lambda\pi^+\eta$	(1.8 ± 0.6) %	690
$\Sigma(1385)^+\eta$	[m] (8.5 ± 3.3) × 10 ⁻³	569
$\Lambda K^+\bar{K}^0$	(6.0 ± 2.1) × 10 ⁻³	441
$\Sigma^0\pi^+$	(9.9 ± 3.2) × 10 ⁻³	824
$\Sigma^+\pi^0$	(1.00 ± 0.34) %	826
$\Sigma^+\eta$	(5.5 ± 2.3) × 10 ⁻³	712
$\Sigma^+\pi^+\pi^-$	(3.4 ± 1.0) %	803
$\Sigma^+\rho^0$	< 1.4 %	CL=95% 578
$\Sigma^-\pi^+\pi^+$	(1.8 ± 0.8) %	798
$\Sigma^0\pi^+\pi^0$	(1.8 ± 0.8) %	802
$\Sigma^0\pi^+\pi^+\pi^-$	(1.1 ± 0.4) %	762
$\Sigma^+\pi^+\pi^-\pi^0$	—	766
$\Sigma^+\omega$	[m] (2.7 ± 1.0) %	568
$\Sigma^+\pi^+\pi^+\pi^-\pi^-$	(3.0 ± 4.1) × 10 ⁻³	707
$\Sigma^+K^+K^-$	(3.5 ± 1.2) × 10 ⁻³	346
$\Sigma^+\phi$	[m] (3.5 ± 1.7) × 10 ⁻³	292
$\Sigma^+K^+\pi^-$	(7 ± 6) × 10 ⁻³	668
Ξ^0K^+	(3.9 ± 1.4) × 10 ⁻³	652
$\Xi^-\pi^+\pi^+$	(4.9 ± 1.7) × 10 ⁻³	564
$\Xi(1530)^0K^+$	[m] (2.6 ± 1.0) × 10 ⁻³	471

Semileptonic modes			
$\Lambda \ell^+ \nu_\ell$	[n]	(2.0 \pm 0.6) %	—
$\Lambda e^+ \nu_e$		(2.1 \pm 0.6) %	870
$\Lambda \mu^+ \nu_\mu$		(2.0 \pm 0.7) %	866
Inclusive modes			
e^+ anything		(4.5 \pm 1.7) %	—
$p e^+$ anything		(1.8 \pm 0.9) %	—
p anything		(50 \pm 16) %	—
p anything (no Λ)		(12 \pm 19) %	—
n anything		(50 \pm 16) %	—
n anything (no Λ)		(29 \pm 17) %	—
Λ anything		(35 \pm 11) %	S=1.4
Σ^\pm anything	[o]	(10 \pm 5) %	—
$\Delta C = 1$ weak neutral current ($C1$) modes, or Lepton number (L) violating modes			
$p \mu^+ \mu^-$	$C1$	< 3.4	$\times 10^{-4}$
$\Sigma^- \mu^+ \mu^+$	L	< 7.0	$\times 10^{-4}$
			CL=90%
			936
			811

 $\Lambda_c(2593)^+$

$I(J^P) = 0(\frac{1}{2}^-)$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant.

Mass $m = 2593.9 \pm 0.8$ MeV $m - m_{\Lambda_c^+} = 308.9 \pm 0.6$ MeV (S = 1.1)Full width $\Gamma = 3.6^{+2.0}_{-1.3}$ MeV

$\Lambda_c^+ \pi\pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

$\Lambda_c(2593)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[p] ≈ 67 %	124
$\Sigma_c(2455)^{++} \pi^-$	24 ± 7 %	17
$\Sigma_c(2455)^0 \pi^+$	24 ± 7 %	23
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	18 ± 10 %	124
$\Lambda_c^+ \pi^0$	not seen	261
$\Lambda_c^+ \gamma$	not seen	290

$\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

Mass $m = 2626.6 \pm 0.8$ MeV ($S = 1.2$)

$m - m_{\Lambda_c^+} = 341.7 \pm 0.6$ MeV ($S = 1.6$)

Full width $\Gamma < 1.9$ MeV, CL = 90%

$\Lambda_c^+ \pi \pi$ and its submode $\Sigma(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[p] $\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	<5	90%	100
$\Sigma_c(2455)^0 \pi^+$	<5	90%	101
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	not seen		293
$\Lambda_c^+ \gamma$	not seen		319

$\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$\Sigma_c(2455)^{++}$ mass $m = 2452.8 \pm 0.6$ MeV

$\Sigma_c(2455)^+$ mass $m = 2453.6 \pm 0.9$ MeV

$\Sigma_c(2455)^0$ mass $m = 2452.2 \pm 0.6$ MeV

$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.87 \pm 0.19$ MeV

$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 168.7 \pm 0.6$ MeV

$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.30 \pm 0.20$ MeV

$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.57 \pm 0.23$ MeV

$m_{\Sigma_c^+} - m_{\Sigma_c^0} = 1.4 \pm 0.6$ MeV

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	90

$\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Sigma_c(2520)^{++}$ mass $m = 2519.4 \pm 1.5$ MeV

$\Sigma_c(2520)^0$ mass $m = 2517.5 \pm 1.4$ MeV

$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 234.5 \pm 1.4$ MeV

$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.6 \pm 1.3$ MeV

$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 1.9 \pm 1.7$ MeV

$\Sigma_c(2520)^{++}$ full width $\Gamma = 18 \pm 5$ MeV

$\Sigma_c(2520)^0$ full width $\Gamma = 13 \pm 5$ MeV

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

$\Sigma_c(2520)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$\Lambda_c^+ \pi$

≈ 100 %

180

Ξ_c^+

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2466.3 \pm 1.4$ MeV

Mean life $\tau = (0.33^{+0.06}_{-0.04}) \times 10^{-12}$ s

$c\tau = 98$ μ m

No absolute branching fractions have been measured. The following are branching ratios relative to $\Xi^- \pi^+ \pi^+$.

Ξ_c^+ DECAY MODES

Fraction (Γ_i/Γ)

p

Confidence level (MeV/c)

$\Lambda K^- \pi^+ \pi^+$

[q] 0.58 ± 0.18

785

$\Lambda \bar{K}^*(892)^0 \pi^+$

[m,q] < 0.29

90%

603

$\Sigma(1385)^+ K^- \pi^+$

[m,q] < 0.41

90%

677

$\Sigma^+ K^- \pi^+$

[q] 1.18 ± 0.31

809

$\Sigma^+ \bar{K}^*(892)^0$

[m,q] 0.92 ± 0.30

654

$\Sigma^0 K^- \pi^+ \pi^+$

[q] 0.49 ± 0.26

734

$\Xi^0 \pi^+$	[q]	0.55 ± 0.16	876
$\Xi^- \pi^+ \pi^+$	[q]	$\equiv 1.0$	850
$\Xi(1530)^0 \pi^+$	[m,q]	<0.2	90%
$\Xi^0 \pi^+ \pi^0$	[q]	2.34 ± 0.68	855
$\Xi^0 \pi^+ \pi^+ \pi^-$	[q]	1.74 ± 0.50	817
$\Xi^0 e^+ \nu_e$	[q]	$2.3 \begin{array}{l} +0.7 \\ -0.9 \end{array}$	883
$p K^- \pi^+$	[q]	0.20 ± 0.05	—

 Ξ_c^0

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

 J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.Mass $m = 2471.8 \pm 1.4$ MeV

$m_{\Xi_c^0} - m_{\Xi_c^+} = 5.5 \pm 1.8$ MeV

Mean life $\tau = (0.098^{+0.023}_{-0.015}) \times 10^{-12}$ s

$c\tau = 29 \mu\text{m}$

 Ξ_c^0 DECAY MODESFraction (Γ_i/Γ) p (MeV/c)

$\Lambda \bar{K}^0$	seen	907
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	788
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	704
$\Xi^- \pi^+$	seen	876
$\Xi^- \pi^+ \pi^+ \pi^-$	seen	817
$p K^- \bar{K}^*(892)^0$	seen	408
$\Omega^- K^+$	seen	523
$\Xi^- e^+ \nu_e$	seen	883
$\Xi^- \ell^+ \text{anything}$	seen	—

 $\Xi_c'^+$

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

 J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.Mass $m = 2574.1 \pm 3.3$ MeV

$m_{\Xi_c'^+} - m_{\Xi_c^+} = 107.8 \pm 3.0$ MeV

The $\Xi_c'^+ - \Xi_c^+$ mass difference is too small for any strong decay to occur. $\Xi_c'^+$ DECAY MODESFraction (Γ_i/Γ) p (MeV/c)

$\Xi_c^+ \gamma$	seen	106
------------------	------	-----

$\Xi_c^{\prime 0}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2578.8 \pm 3.2$ MeV

$$m_{\Xi_c^{\prime 0}} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The $\Xi_c^{\prime 0} - \Xi_c^0$ mass difference is too small for any strong decay to occur.

$\Xi_c^{\prime 0}$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$$\Xi_c^0 \gamma$$

seen

105

$\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Xi_c(2645)^+$ mass $m = 2647.4 \pm 2.0$ MeV ($S = 1.2$)

$\Xi_c(2645)^0$ mass $m = 2644.5 \pm 1.8$ MeV

$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.6 \pm 1.4$ MeV ($S = 1.7$)

$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.2 \pm 1.1$ MeV

$\Xi_c(2645)^+$ full width $\Gamma < 3.1$ MeV, CL = 90%

$\Xi_c(2645)^0$ full width $\Gamma < 5.5$ MeV, CL = 90%

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

$\Xi_c(2645)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$$\Xi_c^0 \pi^+$$

seen

103

$$\Xi_c^+ \pi^-$$

seen

107

$\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$\Xi_c(2815)^+$ mass $m = 2814.9 \pm 1.8$ MeV

$\Xi_c(2815)^0$ mass $m = 2819.0 \pm 2.5$ MeV

$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.6 \pm 1.2$ MeV

$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 347.2 \pm 2.1$ MeV

$\Xi_c(2815)^+$ full width $\Gamma < 3.5$ MeV, CL = 90%

$\Xi_c(2815)^0$ full width $\Gamma < 6.5$ MeV, CL = 90%

The $\Xi_c \pi\pi$ modes are consistent with being entirely via $\Xi_c(2645)\pi$.

$\Xi_c(2815)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi_c^+ \pi^+ \pi^-$	seen	196
$\Xi_c^0 \pi^+ \pi^-$	seen	193



$$I(J^P) = 0(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2704 \pm 4$ MeV ($S = 1.8$)

Mean life $\tau = (0.064 \pm 0.020) \times 10^{-12}$ s

$$c\tau = 19 \text{ } \mu\text{m}$$

Ω_c^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Sigma^+ K^- K^- \pi^+$	seen	697
$\Xi^- K^- \pi^+ \pi^+$	seen	838
$\Omega^- \pi^+$	seen	827
$\Omega^- \pi^- \pi^+ \pi^+$	seen	759

BOTTOM BARYONS ($B = -1$)

$$\Lambda_b^0 = u d b, \Xi_b^0 = u s b, \Xi_b^- = d s b$$

Λ_b^0

$$I(J^P) = 0(\frac{1}{2}^+)$$

$I(J^P)$ not yet measured; $0(\frac{1}{2}^+)$ is the quark model prediction.

Mass $m = 5624 \pm 9$ MeV ($S = 1.8$)

Mean life $\tau = (1.229 \pm 0.080) \times 10^{-12}$ s

$$c\tau = 368 \mu\text{m}$$

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates in Z decay (or high-energy $p\bar{p}$), branching ratios, and detection efficiencies. They scale with the LEP b -baryon production fraction $B(b \rightarrow b\text{-baryon})$ and are evaluated for our value $B(b \rightarrow b\text{-baryon}) = (11.6 \pm 2.0)\%$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note “Production and Decay of b -Flavored Hadrons.”

Λ_b^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$J/\psi(1S)\Lambda$	$(4.7 \pm 2.8) \times 10^{-4}$	1744	
$\Lambda_c^+ \pi^-$	seen	2345	
$\Lambda_c^+ a_1(1260)^-$	seen	2156	
$\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything}$	[r] $(7.9 \pm 1.9) \%$	–	
$p\pi^-$	$< 5.0 \times 10^{-5}$	90%	2732
pK^-	$< 5.0 \times 10^{-5}$	90%	2711

b -baryon ADMIXTURE ($\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$)

$$\text{Mean life } \tau = (1.208 \pm 0.051) \times 10^{-12} \text{ s}$$

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates in Z decay (or high-energy $p\bar{p}$), branching ratios, and detection efficiencies. They scale with the LEP b -baryon production fraction $B(b \rightarrow b\text{-baryon})$ and are evaluated for our value $B(b \rightarrow b\text{-baryon}) = (11.6 \pm 2.0)\%$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note “Production and Decay of *b*-Flavored Hadrons.”

<i>b</i>-baryon ADMIXTURE (Λ_b,Ξ_b,Σ_b,Ω_b)	Fraction (Γ_i/Γ)	<i>p</i> (MeV/c)
$p \mu^- \bar{\nu}$ anything	(4.2 $^{+ 1.8}_{- 1.5}$) %	—
$p \ell \bar{\nu}_\ell$ anything	(4.1 ± 1.0) %	—
p anything	(51 ± 17) %	—
$\Lambda \ell^- \bar{\nu}_\ell$ anything	(2.7 ± 0.8) %	—
$\Lambda/\bar{\Lambda}$ anything	(28 ± 7) %	—
$\Xi^- \ell^- \bar{\nu}_\ell$ anything	(4.8 ± 1.3) $\times 10^{-3}$	—

NOTES

- [a] The masses of the p and n are most precisely known in u (unified atomic mass units). The conversion factor to MeV, $1\text{ u} = 931.494013 \pm 0.000037$ MeV, is less well known than are the masses in u.
- [b] These two results are not independent, and both use the more precise measurement of $|q_{\bar{p}}/m_{\bar{p}}|/(q_p/m_p)$.
- [c] The limit is from neutrality-of-matter experiments; it assumes $q_n = q_p + q_e$. See also the charge of the neutron.
- [d] The first limit is geochemical and independent of decay mode. The second entry, a rough range of limits, assumes the dominant decay modes are among those investigated. For antiprotons the best limit, inferred from the observation of cosmic ray \bar{p} 's is $\tau_{\bar{p}} > 10^7$ yr, the cosmic-ray storage time, but this limit depends on a number of assumptions. The best direct observation of stored antiprotons gives $\tau_{\bar{p}}/\mathcal{B}(\bar{p} \rightarrow e^- \gamma) > 7 \times 10^5$ yr.
- [e] There is some controversy about whether nuclear physics and model dependence complicate the analysis for bound neutrons (from which the best limit comes). The first limit here is from reactor experiments with free neutrons.
- [f] The parameters g_A , g_V , and g_{WM} for semileptonic modes are defined by $\overline{B}_f[\gamma_\lambda(g_V + g_A\gamma_5) + i(g_{WM}/m_{B_i}) \sigma_{\lambda\nu} q^\nu]B_i$, and ϕ_{AV} is defined by $g_A/g_V = |g_A/g_V|e^{i\phi_{AV}}$. See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.
- [g] Time-reversal invariance requires this to be 0° or 180° .
- [h] The decay parameters γ and Δ are calculated from α and ϕ using

$$\gamma = \sqrt{1-\alpha^2} \cos\phi, \quad \tan\Delta = -\frac{1}{\alpha} \sqrt{1-\alpha^2} \sin\phi.$$

See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.

- [i] See the Listings for the pion momentum range used in this measurement.
- [j] The error given here is only an educated guess. It is larger than the error on the weighted average of the published values.
- [k] A theoretical value using QED.
- [l] See the “Note on Λ_c^+ Branching Fractions” in the Branching Fractions of the Λ_c^+ Particle Listings.
- [m] This branching fraction includes all the decay modes of the final-state resonance.
- [n] An ℓ indicates an e or a μ mode, not a sum over these modes.

- [o] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [p] Assuming isospin conservation, so that the other third is $\Lambda_c^+ \pi^0 \pi^0$.
- [q] No absolute branching fractions have been measured. The following are branching *ratios* relative to $\Xi^- \pi^+ \pi^+$.
- [r] Not a pure measurement. See note at head of Λ_b^0 Decay Modes.