

GAUGE AND HIGGS BOSONS

γ

$$I(J^{PC}) = 0,1(1^{--})$$

Mass $m < 2 \times 10^{-16}$ eV
 Charge $q < 5 \times 10^{-30}$ e
 Mean life $\tau = \text{Stable}$

**g
or gluon**

$$I(J^P) = 0(1^-)$$

Mass $m = 0$ [a]
 SU(3) color octet

W

$$J = 1$$

Charge = ± 1 e
 Mass $m = 80.419 \pm 0.056$ GeV
 $m_Z - m_W = 10.76 \pm 0.05$ GeV
 $m_{W^+} - m_{W^-} = -0.2 \pm 0.6$ GeV
 Full width $\Gamma = 2.12 \pm 0.05$ GeV
 $\langle N_{\text{charged}} \rangle = 19.3 \pm 0.4$

W^- modes are charge conjugates of the modes below.

W^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	$\frac{p}{\text{MeV}/c}$
$\ell^+ \nu$	[b] (10.56 \pm 0.14) %		—
$e^+ \nu$	(10.66 \pm 0.20) %	95%	40205
$\mu^+ \nu$	(10.49 \pm 0.29) %	95%	40205
$\tau^+ \nu$	(10.4 \pm 0.4) %		40185
hadrons	(68.5 \pm 0.6) %		—
$\pi^+ \gamma$	< 7	$\times 10^{-5}$	95% 40205
$D_s^+ \gamma$	< 1.3	$\times 10^{-3}$	95% —
cX	(35 \pm 4) %		—
$c\bar{s}$	(32 $^{+13}_{-11}$) %		—
invisible	[c] (1.4 \pm 2.8) %		—

Z

$$J = 1$$

Charge = 0
 Mass $m = 91.1882 \pm 0.0022$ GeV [d]
 Full width $\Gamma = 2.4952 \pm 0.0026$ GeV
 $\Gamma(\ell^+ \ell^-) = 84.057 \pm 0.099$ MeV [b]
 $\Gamma(\text{invisible}) = 499.4 \pm 1.7$ MeV [e]
 $\Gamma(\text{hadrons}) = 1743.8 \pm 2.2$ MeV
 $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-) = 0.9999 \pm 0.0032$
 $\Gamma(\tau^+ \tau^-)/\Gamma(e^+ e^-) = 1.0012 \pm 0.0036$ [f]

Average charged multiplicity

$$\langle N_{\text{charged}} \rangle = 21.07 \pm 0.11$$

Couplings to leptons

$$g_V^\ell = -0.03795 \pm 0.00071$$

$$g_A^\ell = -0.50145 \pm 0.00030$$

$$g^{V_e} = 0.53 \pm 0.09$$

$$g^{V_\mu} = 0.502 \pm 0.017$$

Asymmetry parameters [g]

$$A_e = 0.152 \pm 0.004 \quad (S = 1.2)$$

$$A_\mu = 0.102 \pm 0.034$$

$$A_\tau = 0.141 \pm 0.006$$

$$A_c = 0.66 \pm 0.11$$

$$A_b = 0.91 \pm 0.05$$

Charge asymmetry (%) at Z pole

$$A_{FB}^{(0\ell)} = 1.82 \pm 0.11$$

$$A_{FB}^{(0u)} = 4 \pm 7$$

$$A_{FB}^{(0s)} = 9.8 \pm 1.1$$

$$A_{FB}^{(0c)} = 7.01 \pm 0.45$$

$$A_{FB}^{(0b)} = 10.03 \pm 0.22$$

Z DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	$\frac{p}{\text{MeV}/c}$
$e^+ e^-$	(3.367 \pm 0.005) %		45594
$\mu^+ \mu^-$	(3.367 \pm 0.008) %		45593
$\tau^+ \tau^-$	(3.371 \pm 0.009) %		45559
$\ell^+ \ell^-$	[b] (3.3688 \pm 0.0026) %		—
invisible	(20.02 \pm 0.06) %		—
hadrons	(69.89 \pm 0.07) %		—
$(u\bar{u} + c\bar{c})/2$	(10.1 \pm 1.1) %		—

$(d\bar{d} + s\bar{s} + b\bar{b})/3$	(16.6 ± 0.6) %	—
$c\bar{c}$	(11.68 ± 0.34) %	—
$b\bar{b}$	(15.13 ± 0.05) %	—
$b\bar{b}b\bar{b}$	(4.2 ± 1.6) × 10 ⁻⁴	—
$g g g$	< 1.1 % CL=95%	—
$\pi^0 \gamma$	< 5.2 × 10 ⁻⁵ CL=95%	45593
$\eta \gamma$	< 5.1 × 10 ⁻⁵ CL=95%	45592
$\omega \gamma$	< 6.5 × 10 ⁻⁴ CL=95%	45590
$\eta'(958) \gamma$	< 4.2 × 10 ⁻⁵ CL=95%	45588
$\gamma \gamma$	< 5.2 × 10 ⁻⁵ CL=95%	45594
$\gamma \gamma \gamma$	< 1.0 × 10 ⁻⁵ CL=95%	45594
$\pi^\pm W^\mp$	[h] < 7 × 10 ⁻⁵ CL=95%	10139
$\rho^\pm W^\mp$	[h] < 8.3 × 10 ⁻⁵ CL=95%	10114
$J/\psi(1S) X$	(3.51 ^{+0.23} _{-0.25}) × 10 ⁻³ S=1.1	—
$\psi(2S) X$	(1.60 ± 0.29) × 10 ⁻³	—
$\chi_{c1}(1P) X$	(2.9 ± 0.7) × 10 ⁻³	—
$\chi_{c2}(1P) X$	< 3.2 × 10 ⁻³ CL=90%	—
$\Upsilon(1S) X + \Upsilon(2S) X + \Upsilon(3S) X$	(1.0 ± 0.5) × 10 ⁻⁴	—
$\Upsilon(1S) X$	< 4.4 × 10 ⁻⁵ CL=95%	—
$\Upsilon(2S) X$	< 1.39 × 10 ⁻⁴ CL=95%	—
$\Upsilon(3S) X$	< 9.4 × 10 ⁻⁵ CL=95%	—
$(D^0/\bar{D}^0) X$	(20.7 ± 2.0) %	—
$D^\pm X$	(12.2 ± 1.7) %	—
$D^*(2010)^\pm X$	[h] (11.4 ± 1.3) %	—
$B_s^0 X$	seen	—
$B_c^+ X$	searched for	—
anomalous γ + hadrons	[i] < 3.2 × 10 ⁻³ CL=95%	—
$e^+ e^- \gamma$	[i] < 5.2 × 10 ⁻⁴ CL=95%	45594
$\mu^+ \mu^- \gamma$	[i] < 5.6 × 10 ⁻⁴ CL=95%	45593
$\tau^+ \tau^- \gamma$	[i] < 7.3 × 10 ⁻⁴ CL=95%	45559
$\ell^+ \ell^- \gamma \gamma$	[j] < 6.8 × 10 ⁻⁶ CL=95%	—
$q\bar{q} \gamma \gamma$	[j] < 5.5 × 10 ⁻⁶ CL=95%	—
$\nu\bar{\nu} \gamma \gamma$	[j] < 3.1 × 10 ⁻⁶ CL=95%	45594
$e^\pm \mu^\mp$	LF [h] < 1.7 × 10 ⁻⁶ CL=95%	45593
$e^\pm \tau^\mp$	LF [h] < 9.8 × 10 ⁻⁶ CL=95%	45576
$\mu^\pm \tau^\mp$	LF [h] < 1.2 × 10 ⁻⁵ CL=95%	45576
$p e$	L,B < 1.8 × 10 ⁻⁶ CL=95%	—
$p \mu$	L,B < 1.8 × 10 ⁻⁶ CL=95%	—

Higgs Bosons — H^0 and H^\pm , Searches for

H^0 Mass $m > 95.3$ GeV, CL = 95%

H^\pm in Supersymmetric Models ($m_{H^\pm_1} < m_{H^\pm_2}$)

Mass $m > 82.6$ GeV, CL = 95%

A^0 Pseudoscalar Higgs Boson in Supersymmetric Models [K]

Mass $m > 84.1$ GeV, CL = 95% $\tan\beta > 1$

H^\pm Mass $m > 69.0$ GeV, CL = 95%

See the Particle Listings for a Note giving details of Higgs Bosons.

Heavy Bosons Other Than Higgs Bosons, Searches for

Additional W Bosons

W_R — right-handed W

Mass $m > 715$ GeV, CL = 90% (electroweak fit)

W' with standard couplings decaying to $e\nu, \mu\nu$

Mass $m > 720$ GeV, CL = 95%

Additional Z Bosons

Z'_{SM} with standard couplings

Mass $m > 690$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 898$ GeV, CL = 95% (electroweak fit)

Z'_{LR} of $SU(2)_L \times SU(2)_R \times U(1)$

(with $g_L = g_R$)

Mass $m > 630$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 564$ GeV, CL = 95% (electroweak fit)

Z'_χ of $SO(10) \rightarrow SU(5) \times U(1)_\chi$ (with $g_\chi = e/\cos\theta_W$)

Mass $m > 595$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 545$ GeV, CL = 95% (electroweak fit)

Z'_ψ of $E_6 \rightarrow SO(10) \times U(1)_\psi$ (with $g_\psi = e/\cos\theta_W$)

Mass $m > 590$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 294$ GeV, CL = 95% (electroweak fit)

Z'_η of $E_6 \rightarrow SU(3) \times SU(2) \times U(1) \times U(1)_\eta$ (with $g_\eta = e/\cos\theta_W$)

Mass $m > 620$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 365$ GeV, CL = 95% (electroweak fit)

Scalar Leptoquarks

Mass $m > 225$ GeV, CL = 95% (1st generation, pair prod.)

Mass $m > 200$ GeV, CL = 95% (1st gener., single prod.)

Mass $m > 202$ GeV, CL = 95% (2nd gener., pair prod.)

Mass $m > 73$ GeV, CL = 95% (2nd gener., single prod.)

Mass $m > 99$ GeV, CL = 95% (3rd gener., pair prod.)

(See the Particle Listings for assumptions on leptoquark quantum numbers and branching fractions.)

Axions (A^0) and Other Very Light Bosons, Searches for

The standard Peccei-Quinn axion is ruled out. Variants with reduced couplings or much smaller masses are constrained by various data. The Particle Listings in the full *Review* contain a Note discussing axion searches.

The best limit for the half-life of neutrinoless double beta decay with Majoron emission is $> 7.2 \times 10^{24}$ years (CL = 90%).

NOTES

- [a] Theoretical value. A mass as large as a few MeV may not be precluded.
- [b] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.
- [c] This represents the width for the decay of the W boson into a charged particle with momentum below detectability, $p < 200$ MeV.
- [d] The Z -boson mass listed here corresponds to a Breit-Wigner resonance parameter. It lies approximately 34 MeV above the real part of the position of the pole (in the energy-squared plane) in the Z -boson propagator.
- [e] This partial width takes into account Z decays into $\nu\bar{\nu}$ and any other possible undetected modes.
- [f] This ratio has not been corrected for the τ mass.
- [g] Here $A \equiv 2g_V g_A / (g_V^2 + g_A^2)$.
- [h] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [i] See the Z Particle Listings for the γ energy range used in this measurement.
- [j] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.
- [k] The limits assume no invisible decays.