


$\pi_2(1670)$ $I^G(J^{PC}) = 1^-(2^-+)$ **$\pi_2(1670)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1670 ± 20	OUR ESTIMATE				This is only an educated guess; the error given is larger than the error on the average of the published values.
1672.0 ± 3.5	OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
1687 ± 9	± 15	AMELIN	99 VES	37	$\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1669 ± 4		BARBERIS	98B	450	$p p \rightarrow p_f \rho \pi p_s$
1670 ± 4		BARBERIS	98B	450	$p p \rightarrow p_f f_2(1270) \pi p_s$
1730 ± 20		¹ AMELIN	95B VES	36	$\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1690 ± 14		² BERDNIKOV	94 VES	37	$\pi^- A \rightarrow K^+ K^- \pi^- A$
1710 ± 20	700	ANTIPOV	87 SIGM	50	$\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
1676 ± 6		² EVANGELISTA	81 OMEG	12	$\pi^- p \rightarrow 3\pi p$
1657 ± 14		^{2,3} DAUM	80D SPEC	63–94	$\pi p \rightarrow 3\pi X$
1662 ± 10	2000	² BALTAY	77 HBC	15	$\pi^+ p \rightarrow p 3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1742 ± 31	± 49	ANTREASYAN	90 CBAL		$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
1624 ± 21		⁴ BELLINI	85 SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1622 ± 35		⁵ BELLINI	85 SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1693 ± 28		⁶ BELLINI	85 SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1710 ± 20		⁷ DAUM	81B SPEC	63,94	$\pi^- p$
1660 ± 10		² ASCOLI	73 HBC	5–25	$\pi^- p \rightarrow p \pi_2$

¹ From a fit to $J^{PC} = 2^-+ f_2(1270) \pi, f_0(1370) \pi$ waves.² From a fit to $J^P = 2^- S$ -wave $f_2(1270) \pi$ partial wave.³ Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.⁴ From $f_2(1270) \pi$ decay.⁵ From $\rho \pi$ decay.⁶ From $\sigma \pi$ decay.⁷ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.



$\pi_2(1670)$ mass (MeV)

$\pi_2(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
259 ± 11 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.				
168 ± 43 ± 53		AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268 ± 15		BARBERIS	98B		$450 p p \rightarrow p_f \rho \pi p_s$
256 ± 15		BARBERIS	98B		$450 p p \rightarrow p_f f_2(1270) \pi p_s$
310 ± 20	8	AMELIN	95B	VES	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
190 ± 50	9	BERDNIKOV	94	VES	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$
170 ± 80	700	ANTIPOV	87	SIGM	$50 \pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260 ± 20	9	EVANGELISTA	81	OMEG	$12 \pi^- p \rightarrow 3\pi p$
219 ± 20	9,10	DAUM	80D	SPEC	$63-94 \pi^- p \rightarrow 3\pi X$
285 ± 60	2000	9 BALTAY	77	HBC	$15 \pi^+ p \rightarrow p 3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
236 ± 49 ± 36		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
304 ± 22	11	BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
404 ± 108	12	BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
330 ± 90	13	BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
312 ± 50	14	DAUM	81B	SPEC	$63,94 \pi^- p$
270 ± 60	9	ASCOLI	73	HBC	$5-25 \pi^- p \rightarrow p \pi_2$

⁸ From a fit to $J^{PC} = 2^- + f_2(1270)\pi$, $f_0(1370)\pi$ waves.

⁹ From a fit to $J^P = 2^- f_2(1270)\pi$ partial wave.


¹⁰ Clear phase rotation seen in $2^- S$, $2^- P$, $2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.

¹¹ From $f_2(1270)\pi$ decay.

¹² From $\rho\pi$ decay.

¹³ From $\sigma\pi$ decay.

¹⁴ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.



$\pi_2(1670)$ width (MeV)

$\pi_2(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 3π	$(95.8 \pm 1.4) \%$	
Γ_2 $f_2(1270)\pi$	$(56.2 \pm 3.2) \%$	
Γ_3 $\rho\pi$	$(31 \pm 4) \%$	
Γ_4 $\sigma\pi$	$(13 \pm 6) \%$	
Γ_5 $f_0(1370)\pi$	$(8.7 \pm 3.4) \%$	
Γ_6 $K\bar{K}^*(892) + \text{c.c.}$	$(4.2 \pm 1.4) \%$	

Γ_7	$\omega\rho$	(2.7±1.1) %		
Γ_8	$\gamma\gamma$			
Γ_9	$\eta\pi$			
Γ_{10}	$\pi^\pm 2\pi^+ 2\pi^-$			
Γ_{11}	$\rho(1450)\pi$	< 3.6	$\times 10^{-3}$	97.7%
Γ_{12}	$b_1(1235)\pi$	< 1.9	$\times 10^{-3}$	97.7%

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-53			
x_5	-29	-59		
x_6	-8	-21	-9	
	x_2	x_3	x_5	

$\pi_2(1670)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$				Γ_8
VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG COMMENT
<0.072	90	15 ACCIARRI	97T L3	$e^+ e^- \rightarrow$
<0.19	90	15 ALBRECHT	97B ARG	$e^+ e^- \pi^+ \pi^- \pi^0$ $e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.41 ±0.23±0.28		ANTREASYAN 90	CBAL 0	$e^+ e^- \rightarrow$ $e^+ e^- \pi^0 \pi^0 \pi^0$
0.8 ±0.3 ±0.12		16 BEHREND	90C CELL 0	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^- \pi^0$
1.3 ±0.3 ±0.2		17 BEHREND	90C CELL 0	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^- \pi^0$

¹⁵ Decaying into $f_2(1270)\pi$ and $\rho\pi$.

¹⁶ Constructive interference between $f_2(1270)\pi, \rho\pi$ and background.

¹⁷ Incoherent Ansatz.

$\pi_2(1670)$ BRANCHING RATIOS

$\Gamma(3\pi)/\Gamma_{\text{total}}$		$\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3 + \Gamma_5)/\Gamma$
0.958±0.014 OUR FIT		

$\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$

VALUE

0.29±0.04 OUR FIT

0.29±0.05

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.3

18 DAUM 81B SPEC 63,94 $\pi^- p$

18 From a two-resonance fit to four 2^-0^+ waves.

$\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$

(With $f_2(1270) \rightarrow \pi^+\pi^-$)

VALUE

0.604±0.035 OUR FIT

0.60 ±0.05 OUR AVERAGE

Error includes scale factor of 1.3. See the ideogram below.

0.61 ±0.04

19 DAUM 81B SPEC 63,94 $\pi^- p$

0.76 +0.24
-0.34

ARMENISE 69 DBC + 5.1 $\pi^+ d \rightarrow d 3\pi$

0.35 ±0.20

BALTAY 68 HBC + 7–8.5 $\pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.59

BARTSCH 68 HBC + 8 $\pi^+ p \rightarrow 3\pi p$


19 From a two-resonance fit to four 2^-0^+ waves.

$\frac{1}{2}\Gamma_3/(0.567\Gamma_2 + \frac{1}{2}\Gamma_3 + 0.624\Gamma_5)$

DOCUMENT ID TECN CHG COMMENT

WEIGHTED AVERAGE
0.60±0.05 (Error scaled by 1.3)

Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.



$\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$

(With $f_2(1270) \rightarrow \pi^+\pi^-$)

$\Gamma_3/0.564\Gamma_2$

VALUE

1.01±0.05

DOCUMENT ID COMMENT

BARBERIS 98B 450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$
(All η decays.)

VALUE

<0.09

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10

DOCUMENT ID

BALTAY

TECN

HBC

CHG

+

COMMENT

7–8.5 $\pi^+ p$

CRENNELL 70 HBC – 6 $\pi^- p \rightarrow$
 $f_2 \pi^- N$

$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$

VALUE

<0.10

<0.1

$\Gamma_9/(0.567\Gamma_2 + \frac{1}{2}\Gamma_3 + 0.624\Gamma_5)$

DOCUMENT ID

CRENNELL 70 HBC

TECN

HBC

CHG

–

COMMENT

6 $\pi^- p \rightarrow$
 $f_2 \pi^- N$

BALTAY 68 HBC

+

7.8.5 $\pi^+ p$

$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$

VALUE

<0.0036

97.7

Γ_{11}/Γ

DOCUMENT ID

AMELIN 99 VES

TECN

VES

COMMENT

37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

|

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$

VALUE

<0.0019

97.7

Γ_{12}/Γ

DOCUMENT ID

AMELIN 99 VES

TECN

VES

COMMENT

37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

|

$\Gamma(f_0(1370)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$

(With $f_0(1370) \rightarrow \pi^+ \pi^-$.)

VALUE

0.10±0.04 OUR FIT

0.10±0.05

DOCUMENT ID

20 DAUM

TECN

COMMENT

81B SPEC 63,94 $\pi^- p$

20 From a two-resonance fit to four $2^- 0^+$ waves.

$0.624\Gamma_5/(0.567\Gamma_2 + \frac{1}{2}\Gamma_3 + 0.624\Gamma_5)$

$\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(f_2(1270)\pi)$

VALUE

0.075±0.025 OUR FIT

0.075±0.025

DOCUMENT ID

21 ARMSTRONG 82B OMEG

TECN

COMMENT

– 16 $\pi^- p \rightarrow K^+ K^- \pi^- p$

21 From a partial-wave analysis of $K^+ K^- \pi^-$ system.

Γ_6/Γ_2

$\Gamma(\omega\rho)/\Gamma_{\text{total}}$

VALUE

0.027±0.004±0.010

23 AMELIN

TECN

COMMENT

37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

|

$\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi)$

VALUE

0.24±0.10

DOCUMENT ID

24,25 BAKER

TECN

COMMENT

99 SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

Γ_7/Γ

D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.18±0.06	24 BAKER	99 SPEC	$1.94 \bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.22±0.10	22 DAUM	81B SPEC	$63.94 \pi^- p$
22 From a two-resonance fit to four 2^-0^+ waves.			
23 Normalized to the $B(\pi_2(1670) \rightarrow f_2\pi)$.			
24 Using preliminary CBAR data.			
25 With the $\sigma\pi$ in $L=2$ and the $f_2(1270)\pi$ in $L=0$.			

 $\pi_2(1670)$ REFERENCES

AMELIN	99	PAN 62 445 Translated from YAF 62	D.V. Amelin <i>et al.</i> 487.	(VES Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA102 Collab.)
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
		Translated from YAF 41	1223.	
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
EVANGELISTA	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
Also	81B	NP B186 594	C. Evangelista	
DAUM	80D	PL 89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TNTO, GENO, HAMB, MILA+) JP
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)
ARMENISE	69	LNC 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP

OTHER RELATED PAPERS

ZAIMIDOROGA	99	PAN 30 1 Translated from SJPN 30 5.	O.A. Zaimidoroga	
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
CHEN	83B	PR D28 2304	T.Y. Chen <i>et al.</i>	(ARIZ, FNAL, FLOR, NDAM+)
LEEDOM	83	PR D27 1426	I.D. Leedom <i>et al.</i>	(PURD, TNTO)
BELLINI	82B	NP B199 1	G. Bellini <i>et al.</i>	(CERN, MILA, JINR+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
PERNEGR	78	NP B134 436	J. Pernegr <i>et al.</i>	(ETH, CERN, LOIC+)
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)
LEV RAT	66	PL 22 714	B. Levrat <i>et al.</i>	
VETLITSKY	66	PL 21 579	I.A. Veltlitsky <i>et al.</i>	(ITEP)
FORINO	65B	PL 19 68	A. Forino <i>et al.</i>	(BGNA, BARI, FIRZ, ORSAY+)