

$K_2(2250)$ $I(J^P) = \frac{1}{2}(2^-)$

OMMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave.

 $K_2(2250)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2247 ± 17 OUR AVERAGE					
2200 \pm 40		1 ARMSTRONG 83C OMEG	—	18	$K^- p \rightarrow \Lambda \bar{p} X$
2235 \pm 50		1 BAUBILLIER 81 HBC	—	8	$K^- p \rightarrow \Lambda \bar{p} X$
2260 \pm 20		1 CLELAND 81 SPEC	\pm	50	$K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2147 \pm 4	37	CHLIAPNIK... 79 HBC	+	32	$K^+ p \rightarrow \bar{\Lambda} p X$
2240 \pm 20	20	LISSAUER 70 HBC		9	$K^+ p$
$^1 J^P = 2^-$ from moments analysis.					

 $K_2(2250)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
180 ± 30 OUR AVERAGE					
Error includes scale factor of 1.4.					
150 \pm 30		2 ARMSTRONG 83C OMEG	—	18	$K^- p \rightarrow \Lambda \bar{p} X$
210 \pm 30		2 CLELAND 81 SPEC	\pm	50	$K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
\sim 200		2 BAUBILLIER 81 HBC	—	8	$K^- p \rightarrow \Lambda \bar{p} X$
\sim 40	37	CHLIAPNIK... 79 HBC	+	32	$K^+ p \rightarrow \bar{\Lambda} p X$
80 \pm 20	20	LISSAUER 70 HBC		9	$K^+ p$
$^2 J^P = 2^-$ from moments analysis.					

 $K_2(2250)$ DECAY MODES

Mode

Γ_1	$K \pi \pi$
Γ_2	$p \bar{\Lambda}$

 $K_2(2250)$ REFERENCES

ARMSTRONG 83C NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BAUBILLIER 81 NP B183 1	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+) JP
CLELAND 81 NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+) JP
CHLIAPNIK... 79 NP B158 253	P.V. Chliapnikov <i>et al.</i>	(CERN, BELG, MONS)
LISSAUER 70 NP B18 491	D. Lissauer <i>et al.</i>	(LBL)

OTHER RELATED PAPERS

ALEXANDER 68B PRL 20 755

G. Alexander *et al.*

(LRL)